

the desired product (99.8 mg, 79%) as a colorless oil. Proton NMR indicated a 3:2 mixture of isomers due to partial epimerization at alanine chiral center. MS found: $(M+H)^+ = 369$.

- 5 (18b) Following a procedure analogous to (1e), the ester from (18a) (94.5 mg, 0.256 mmol) was reacted with hydroxylamine to give the hydroxamic acid (90.1 mg, 95%) as a white solid. MS found: $(M-H)^- = 368$.

Example 19

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-[(4-pyridinyl)methoxy]phenyl]-1-pyrrolidineacetamide

- 10 (19a) Cesium carbonate (331 mg, 2.8 eq) was added to the phenol from (7a) (100.7 mg, 0.363 mmol), and 4-picolyl chloride hydrochloride (119 mg, 2 eq) in methyl sulfoxide (2 mL). After 20 h at rt, same portions of cesium carbonate and 4-picolyl chloride hydrochloride were added. After 30 min at 75 °C, saturated ammonium chloride (6 mL) and ethyl acetate (100 mL) were added. The mixture was washed with water (6 mL), brine (6 mL), dried (MgSO₄) and concentrated. Silica gel chromatography (ethyl acetate) gave the desired product (106.7 mg, 80%) as a colorless oil. Proton NMR indicated a 4.5:1 mixture of isomers due to partial epimerization at alanine chiral center. MS found: $(M+H)^+ = 369$.
- 20 (19b) Following a procedure analogous to (1e), the ester from (19a) (99.8 mg, 0.271 mmol) was reacted with hydroxylamine to give the hydroxamic acid (81.2 mg, 81%) as a white solid. MS found: $(M-H)^- = 368$.

Example 20

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-(2-methylpropyl)phenyl]-2-oxo-1-pyrrolidineacetamide

- 30 (20a) Iodomethane (3.82 mL, 2.5 eq.) was added to a mixture of ibuprofen (4.97 g, 24.1 mmol), 1,8-diazabicyclo[4.3.0]non-5-ene (4.32 mL, 1.2 eq.) and benzene (100 mL) and the mixture was heated to reflux for 1 h. Following addition of hexane (100 mL), the mixture was filtered through a silica gel pad and the filter cake washed with ether-hexane (1:1, v/v) until
- 35

free of product. The filtrate was concentrated in vacuo to give the methyl ester as a colorless liquid (5.12 g, 96%).

(20b) Following a procedure analogous to (1a), ibuprofen methyl ester from (20a) (4.655 g) was reacted with sodium bis(trimethylsilyl)amide and allyl bromide to yield crude product (6.39 g) as a yellow liquid. This material was used in the subsequent reaction without purification.

(20c) Following a procedure analogous to (1c), the crude material from (20b) (6.19 g) was ozonolyzed to give crude aldehyde (6.53 g) as a yellow oil. This material was used in the subsequent reaction without purification.

(20d) Following a procedure analogous to (1d), crude aldehyde from (20c) (2.05 g) was reacted with D-alanine methyl ester hydrochloride. Silica gel chromatography (ethyl acetate-hexane, 20:80 then 30:70) gave less polar isomer (371.8 mg), more polar isomer (289.6 mg), and a 1:3 mixture of the two isomers (337.8 mg). The total yield is 999.2 mg (49% for three steps). MS found: $(M+H)^+ = 318$.

(20e) Following a procedure analogous to (1e), the less polar isomer from (20d) (210 mg, 0.660 mmol) was reacted with hydroxylamine to give the hydroxamic acid (186.7 mg, 89%). MS found: $(M-H)^- = 317$.

(20f) Following a procedure analogous to (1e), the more polar isomer from (20d) (200 mg, 0.630 mmol) was reacted with hydroxylamine to give the hydroxamic acid (167.2 mg, 83%) as a white solid. MS found: $(M-H)^- = 317$.

Example 21

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-phenyl-1-pyrrolidineacetamide

(21a) Following a procedure analogous to (20a), 2-phenylpropionic acid (10.0 g, 66.5 mmol) was reacted with iodomethane and 1,8-diazabicyclo[4.3.0]non-5-ene to give the ester (9.57 g, 88%) as a colorless liquid.

(21b) Following a procedure analogous to (1a), the methyl ester from (21a) (9.28 g, 56.5 mmol) was reacted with sodium bis(trimethylsilyl)amide and allyl bromide to yield crude

product (11.96 g) as a yellow liquid. This material was used in the subsequent reaction without purification.

(21c) Following a procedure analogous to (1c), the crude material from (21b) (6.76 g) was ozonolyzed to give crude aldehyde (8.53 g) as a yellow oil. This material was used in the subsequent reaction without purification.

(21d) Following a procedure analogous to (1d), the crude aldehyde from (21c) (1.93 g) was reacted with D-alanine methyl ester hydrochloride. Silica gel chromatography (ethyl acetate-hexane, 30:70 then 40:60) gave less polar isomer (230 mg), more polar isomer (270 mg), and a 3:2 mixture of the two isomers (380 mg). The total yield is 880 mg (47% for three steps). MS found: $(M+H)^+ = 262$.

(21e) Following a procedure analogous to (1e), the less polar isomer from (21d) (141.1 mg, 0.540 mmol) was reacted with hydroxylamine to give the hydroxamic acid (141.5 mg, 100%) as a solid. MS found: $(M-H)^- = 261$.

(21f) Following a procedure analogous to (1e), the more polar isomer from (21d) (165.2 mg, 0.632 mmol) was reacted with hydroxylamine to give the hydroxamic acid (149.6 mg, 90%) as a solid. MS found: $(M-H)^- = 261$.

Example 22

N-hydroxy-2-oxo-3-phenyl-1-pyrrolidineacetamide

(22a) Following a procedure analogous to (1a), methyl phenylacetate (10.0 mL, 69.2 mmol) was reacted with sodium bis(trimethylsilyl)amide and allyl bromide to yield the desired (13.10 g, 100%) as a colorless liquid.

(22b) Following a procedure analogous to (1c), the material from (22a) (7.06 g, 36.8 mmol) was ozonolyzed to give crude aldehyde (9.00 g) as a yellow oil. This material was used in the subsequent reaction without purification.

(22c) Following a procedure analogous to (1d), the crude aldehyde from (22b) (2.00 g) was reacted with glycine methyl ester hydrochloride. Silica gel chromatography (ethyl acetate-hexane, 50:50) gave the desired lactam (1.05 g, 55% for two steps).

(22d) Following a procedure analogous to (1e), the lactam from (22c) (433.8 mg, 1.86 mmol) was reacted with hydroxylamine to give the hydroxamic acid (261 mg, 60%) as a yellow powder. MS found: $(M-H)^- = 233$.

5

Example 23

(+/-)-N-hydroxy-3-methyl-2-oxo-3-phenyl-1-pyrrolidineacetamide

(23a) Following a procedure analogous to (1d), the crude aldehyde from (21c) (2.19 g) was reacted with glycine methyl ester hydrochloride. Silica gel chromatography (ethyl acetate-hexane, 35:65) gave the desired lactam (650 mg, 32% for three steps) as a colorless oil. MS found: $(M+H)^+ = 248$.
(23b) Following a procedure analogous to (1e), the lactam from (23a) (433.8 mg, 1.86 mmol) was reacted with hydroxylamine to give the hydroxamic acid (261 mg, 90%) as a white powder. MS found: $(M-H)^- = 247$.

Example 24

[1(R)]-N-hydroxy- α -methyl-2-oxo-3-phenyl-1-pyrrolidineacetamide

(24a) Following a procedure analogous to (1d), the crude aldehyde from (22b) (2.00 g) was reacted with D-alanine methyl ester hydrochloride. Silica gel chromatography (ethyl acetate-hexane, 30:70 then 40:60 then 50:50) gave less polar isomer (309.3 mg), more polar isomer (347.2 mg), and a 1:1 mixture of the two isomers (163.4 mg). The total yield is 819.9 mg (41% for two steps). MS found: $(M+H)^+ = 248$.
(24b) Following a procedure analogous to (1e), the less polar isomer from (24a) (243.7 mg, 0.985 mmol) was reacted with hydroxylamine to give the hydroxamic acid (210 mg, 86%) as a white solid. MS found: $(M-H)^- = 247$.
(24c) Following a procedure analogous to (1e), the more polar isomer from (24a) (202.8 mg, 0.820 mmol) was reacted with hydroxylamine to give the hydroxamic acid (180 mg, 88%) as a white solid. MS found: $(M-H)^- = 247$.

Example 25**[1(R)]-N-hydroxy-3-(4-methoxyphenyl)- α -methyl-2-oxo-1-pyrrolidineacetamide**

(25a) Following a procedure analogous to (1c), the crude material from (2a) (8.22 g) was ozonolyzed to give crude aldehyde (8.22 g) as a yellow oil. This material was used in the subsequent reaction without purification.

(25b) Following a procedure analogous to (1d), the crude aldehyde from (25a) (2.21 g) was reacted with D-alanine methyl ester hydrochloride. Silica gel chromatography (ethyl acetate-hexane, 45:55 then 50:50) gave less polar isomer (215.8 mg), more polar isomer (181.1 mg), and a 1:1 mixture of the two isomers (623 mg). The total yield is 1.020 g (49% for three steps). MS found: $(M+H)^+ = 278$.

(25c) Following a procedure analogous to (1e), the less polar isomer from (25b) (154.6 mg, 0.557 mmol) was reacted with hydroxylamine to give the hydroxamic acid (120.4 mg, 78%) as a viscous oil. MS found: $(M-H)^- = 277$.

(25d) Following a procedure analogous to (1e), the more polar isomer from (25b) (130.3 mg, 0.470 mmol) was reacted with hydroxylamine to give the hydroxamic acid (117.9 mg, 90%) as a solid. MS found: $(M-H)^- = 277$.

Example 26**[1(R)]-3-cyclohexyl-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide**

(26a) A mixture of the more polar isomer from (24a) (36.5 mg, 0.14 mmol), rhodium on alumina (17 mg), 4 N dioxane solution of hydrogen chloride (2 drops) and methanol (2 mL) was hydrogenated under 45 psi overnight. The mixture was filtered through a celite pad and the filter cake washed with ethyl acetate-hexane (40:60). The filtrate was concentrated to give the desired product (37.4 mg, 100%) as a colorless liquid. MS found: $(M+H)^+ = 268$.

(26b) Following a procedure analogous to (1e), the ester from (26a) (52.4 mg, 0.196 mmol) was reacted with hydroxylamine to give the hydroxamic acid (25.2 mg, 48%) as a solid. MS found: $(M-H)^- = 267$.

Example 27**[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-(2-phenylethyl)-1-pyrrolidineacetamide**

- 5 (27a) A 2.5 M hexane solution of n-butyllithium (5.12 mL, 1.1 eq) was added dropwise to diisopropylamine (1.80 mL, 1.1 eq) in tetrahydrofuran (50 mL) at 0 °C. The resultant mixture was stirred for 20 min at 0 °C and cooled to -78 °C. A solution of ethyl 2-methyl-4-pentenoate (1.90 mL, 11.7 mmol) in
- 10 tetrahydrofuran (25 mL) was added. The mixture was stirred at -78 °C for 30 min and warmed to 0 °C. 2-Phenylethyl bromide (1.71 mL, 1.05 eq) in tetrahydrofuran (25 mL) was added dropwise. After additional 2 h at 0 °C, saturated ammonium chloride (50 mL) was added and the mixture extracted with
- 15 ethyl acetate (3 x). The combined extracts were washed with brine, dried (MgSO₄) and concentrated. Silica gel chromatography (ethyl acetate-hexane, 0:100 then 5:95) gave the desired product (1.95 g, 68%) as a liquid. MS found: (M+H)⁺ = 247.
- 20 (27b) Following a procedure analogous to (1c), the olefin from (27a) (1.86 g, 7.55 mmol) was ozonolyzed. Silica gel chromatography (ethyl acetate-hexane, 10:90) gave the desired aldehyde (1.67 g, 89%) as a colorless oil. MS found: (M+H)⁺ = 249.
- 25 (27c) Following a procedure analogous to (1d), the aldehyde from (27b) (1.66 g, 6.68 mmol) was reacted with D-alanine methyl ester hydrochloride. Silica gel chromatography (ethyl acetate-hexane, 35:65 then 40:60) gave the lactam (1.32 g, 68%) as a 1:1 mixture of two diastereomers. MS found: (M+H)⁺
- 30 = 290.
- (27d) Following a procedure analogous to (1e), the ester from (27c) (52.4 mg, 0.196 mmol) was reacted with hydroxylamine to give the hydroxamic acid (226.6 mg, 96%) as a 1:1 mixture of two isomers. MS found: (M-H)⁻ = 289.

35

Example 28**[1(R)]-3-(2-cyclohexylethyl)-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide**

(28a) Following a procedure analogous to (26a), the ester from (27c) (180 mg, 0.622 mmol) was hydrogenated to give the desired product (184 mg, 100%) as a colorless oil. MS found: $(M+H)^+ = 296$.

- 5 (28b) Following a procedure analogous to (1e), the ester from (28a) (160 mg, 0.542 mmol) was reacted with hydroxylamine to give the hydroxamic acid (158 mg, 98%) as a 1:1 mixture of two isomers. MS found: $(M-H)^- = 295$.

10

Example 29

[1(R)]-N-hydroxy- α -methyl-2-oxo-3-phenyl-3-(phenylmethyl)-2-oxo-1-pyrrolidineacetamide

(29a) Following a procedure analogous to (20a), 2,3-diphenylacetic acid (10.26 g, 45.34 mmol) was reacted with iodomethane and 1,8-diazabicyclo[4.3.0]non-5-ene to give the ester (10.86 g, 100%) as a colorless liquid. MS found: $(M+H)^+ = 241$.

15

(29b) Following a procedure analogous to (1a), the ester from (29a) (10.56 g, 43.9 mmol) was reacted with sodium bis(trimethylsilyl)amide and allyl bromide to yield crude product (13.13 g) as a pale yellow oil. This material was used in the subsequent reaction without purification.

20

(29c) Following a procedure analogous to (1c), the crude material from (29b) (6.07 g) was ozonolyzed to give the crude aldehyde (7.10 g) as a yellow oil. This material was used in the subsequent reaction without purification.

25

(29d) Following a procedure analogous to (1d), the crude aldehyde from (29c) (2.08 g) was reacted with D-alanine methyl ester. Silica gel chromatography (ethyl acetate-hexane, 20:80 then 30:70) gave a 1:1 mixture of lactams (1.07 g, 53% for three steps) as a colorless viscous oil. MS found: $(M+H)^+ = 338$.

30

(29e) Following a procedure analogous to (1e), the ester from (29d) (980 mg, 2.90 mmol) was reacted with hydroxylamine to give the hydroxamic acid as a 1:1 mixture of two isomers. MS found: $(M-H)^- = 337$.

35

Example 30

[1(R)]-3,4,4',5'-tetrahydro-N-hydroxy- α -methyl-2-oxospiro[naphthalene-2(1H),3'-[3H]pyrrole]-1'(2'H)-acetamide

5

(30a) Following a procedure analogous to (20a), 1,2,3,4-tetrahydro-2-naphthoic acid (4.50 g, 25.5 mmol) was reacted with iodomethane and 1,8-diazabicyclo[4.3.0]non-5-ene to give the ester (4.62 g, 95%) as a pale yellow liquid. MS found:

10 (M+H)⁺ = 191.

(30b) Following a procedure analogous to (1a), the ester from (30a) (4.52 g) was reacted with sodium bis(trimethylsilyl)amide and allyl bromide to yield crude product (5.20 g) as a yellow oil. This material was used in the subsequent reaction without purification.

15

(30c) Following a procedure analogous to (1c), the crude olefin from (30b) (5.00 g) was ozonolyzed to give crude aldehyde (5.83 g) as a yellow oil. This material was used in the subsequent reaction without purification.

20

(30d) Following a procedure analogous to (1d), the crude aldehyde from (30c) (2.03 g) was reacted with D-alanine methyl ester hydrochloride. Silica gel chromatography (ethyl acetate-hexane, 30:70 then 40:60) gave a 1:1 mixture of lactams (732.1 mg, 34% for three steps). MS found: (M+H)⁺ = 288.

25

(30e) Following a procedure analogous to (1e), the ester from (30d) (510.7 mg, 1.788 mmol) was reacted with hydroxylamine to give the hydroxamic acid (431 mg, 84%) as a 1:1 mixture of two isomers. MS found: (M-H)⁻ = 287.

30

Example 31

[1(R)]-3-[4-[(3,5-dibromophenyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3,5-dibromobenzyl bromide, example 31 was prepared in an analogous series of reactions to (6b) and (6c). MS found: (M-H)⁻ = 523.

35

Example 32

[1(R)]-3-[4-[[3,5-bis(trifluoromethyl)phenyl]methoxy]phenyl]-N-hydroxy-
α,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3,5-bis(trifluoromethyl)benzyl bromide, example 32 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M-H)^- = 503$.

Example 33

[1(R)]-3-[4-[(3,5-dichlorophenyl)methoxy]phenyl]-N-hydroxy-
α,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3,5-dichlorobenzyl chloride, example 33 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M-H)^- = 435$.

Example 34

[1(R)]-N-hydroxy-α,3-dimethyl-3-[4-[(2-methyl-1-naphthalenyl)methoxy]phenyl]-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 1-chloromethyl-2-methylnaphthalene, example 34 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M+Na)^+ = 455$.

Example 35

[1(R)]-3-[4-[(3,5-dimethoxyphenyl)methoxy]phenyl]-N-hydroxy-
α,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3,5-dimethoxybenzyl chloride, example 35 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M-H)^- = 427$.

Example 36

[1(R)]-3-[4-[[4-chloro-2-(trifluoromethyl)-6-quinolinyl]methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 6-bromomethyl-4-chloro-2-trifluoromethylquinoline, example 36 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M-H)^- = 520$.

Example 37

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-[[4-(1,2,3-thiadiazol-4-yl)phenyl]methoxy]phenyl]-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 4-(4-bromomethylphenyl)-1,2,3-thiadiazole, example 37 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M-H)^- = 451$.

Example 38

[1(R)]-3-[4-([1,1'-biphenyl]-2-ylmethoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 2-phenylbenzyl bromide, example 38 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M-H)^- = 443$.

Example 39

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 4-bromomethyl-2,6-dichloropyridine, example 39 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M-H)^- = 436$.

Example 40

[1(R)]-3-[4-(1H-benzotriazol-1-ylmethoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 1-chloromethylbenzotriazole, example 40 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M-H)^- = 408$.

5

Example 41

[1(R)]-3-[4-[(4,6-dimethyl-2-pyrimidinyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

10 Beginning with the phenol from (6a) and 2-chloromethyl-4,6-dimethylpyrimidine (Sakamoto et al, *Heterocycles* **1997**, 6, 525), example 41 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M-H)^- = 397$.

15

Example 42

[1(R)]-3-[4-(1,3-benzodioxol-5-ylmethoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

20 Beginning with the phenol from (6a) and 3,4-methylenedioxybenzyl chloride, example 42 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M-H)^- = 411$.

Example 43

25 [1(R)]-3-[4-[(2-chloro-6-ethoxy-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

30 Beginning with the phenol from (6a) and 4-bromomethyl-2-chloro-6-ethoxypyridine, example 43 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M-H)^- = 446$.

Example 44

35 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-(4-guininylmethoxy)phenyl]-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 4-chloromethylquinoline, example 44 was prepared in an analogous

series of reactions to (6b) and (6c). MS found: $(M+H)^+ = 420$.

Example 45

5 [1(R)]-3-[4-[(4,5-dimethyl-2-thiazolyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 2-bromomethyl-4,5-dimethylthiazole, example 45 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M-H)^- = 402$.

Example 46

15 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

Beginning with the phenol from (6a) and 4-chloromethyl-2,6-dimethylpyridine, example 46 was prepared in an analogous series of reactions to (6b) and (6c). MS found: $(M+H)^+ = 398$.

Example 47

25 [1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-[(3-methyl-5-nitrophenyl)methoxy]phenyl]-2-oxo-1-pyrrolidineacetamide

(47a) Following a procedure analogous to (6b), the phenol from (6a) (500 mg, 1.80 mmol) was reacted with 5-methyl-3-nitrobenzyl bromide to give the desired ether (690 mg, 90%). MS found: $(M+Na)^+ = 449$.

(47b) Following a procedure analogous to step (1f), the ester from (47a) (67.4 mg, 0.158 mmol) was reacted with hydroxylamine to give the hydroxamic acid (48.7 mg, 72%). MS found: $(M-H)^- = 426$.

Example 48

35 [1(R)]-3-[4-[(3-amino-5-methylphenyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

(48a) Zinc powder (2.5 g) was added to the ester from (47a) (670 mg, 1.57 mmol) in acetic acid (10 mL) and the mixture was stirred at 50 °C for 2 h. The solid was removed by filtration and washed with ethyl acetate. The filtrate was concentrated, treated with brine (15 mL) and 1 N NaOH (15 mL), and extracted with ethyl acetate (3 x). The combined extracts were dried (MgSO₄) and concentrated. Silica gel chromatography (ethyl acetate-hexane, 45:55 then 55:45) gave the desired aniline (610 mg, 98%). MS found: $(M+H)^+ = 397$.

(48b) Following a procedure analogous to step (1f), the ester from (48a) (80 mg, 0.202 mmol) was reacted with hydroxylamine to give the hydroxamic acid (63 mg, 79%). MS found: $(M-H)^- = 396$.

Example 49

[1(R)]-3-[4-[[3-(acetylamino)-5-methylphenyl]methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

(49a) Hunig's base (74 mg, 5 eq) and acetyl chloride (23 mg, 2 eq) were added sequentially to the aniline from (48a) (58 mg, 0.146 mmol) in dichloromethane (2.5 mL) at 0 °C. After 30 min at this temperature, saturated NaHCO₃ (5 mL) and ethyl acetate (100 mL) were added. The organic phase was separated, washed with brine (5 mL), dried (MgSO₄) and concentrated. Silica gel chromatography (ethyl acetate-hexane, 70:30) gave the acetamide (45 mg, 78%). MS found: $(M+Na)^+ = 461$.

(49b) Following a procedure analogous to step (1f), the ester from (49a) (40 mg, 0.091 mmol) was reacted with hydroxylamine to give the hydroxamic acid (27 mg, 67%). MS found: $(M-H)^- = 438$.

Example 50

[1(R)]-1,1-dimethylethyl [2-[[3-[[4-[1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-3-methyl-2-oxo-3-pyrrolidinyl]phenoxy]methyl]-5-methylphenyl]amino]-2-oxoethyl]carbamate

(50a) A mixture of the aniline from (48a) (100 mg, 0.252 mmol), N-(t-butoxycarbonyl)glycine (53 mg, 1.2 eq), BOP-Cl

(70.6 mg, 1.1 eq), NMM (76.5 mg, 3 eq) and THF (10 mL) were heated to reflux for 30 min. Following addition of water (15 mL) and sat K₂CO₃, THF was removed *in vacuo*. The aqueous residue was extracted with ethyl acetate (3 x 40 mL). The combined organic extracts were dried (MgSO₄) and concentrated. Silica gel chromatography (MeOH-CH₂Cl₂, 5:95) gave the desired amide (130 mg, 93%). MS found: (M+Na)⁺ = 576.

(50b) Following a procedure analogous to step (1f), the ester from (50a) (120 mg, 0.217 mmol) was reacted with hydroxylamine to give the hydroxamic acid (100 mg, 83%). MS found: (M-H)⁻ = 553.

Example 51

[1(R)]-3-[4-[[3-[(aminoacetyl)amino]-5-methylphenyl]methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

The hydroxamic acid from (50b) (60 mg, 0.108 mmol) was stirred with trifluoroacetic acid (1 mL) and CH₂Cl₂ (1 mL) for 2 h at rt and concentrated to give the TFA salt (58 mg, 94%). MS found: (M+H)⁺ = 455.

Example 52

[1(R)]-1,1-dimethylethyl [2-[[2-[[3-[[4-[1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-3-methyl-2-oxo-3-pyrrolidinyl]phenoxy]methyl]-5-methylphenyl]amino]-2-oxoethyl]amino]-2-oxoethyl]carbamate

Beginning with the aniline from (48a) and BOC-Gly-Gly-OH, example 52 was prepared in an analogous series of reactions to (50a) and (50b). MS found: (M+Na)⁺ = 634.

Example 53

[1(R)]-3-[4-[[3-[[[(aminoacetyl)amino]acetyl]amino]-5-methylphenyl]methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

Beginning with the hydroxamic acid from example 52, example 53 was prepared following a procedure analogous to example 51. MS found: (M+H)⁺ = 512.

Example 54

[1(R)]-N-[3-[[4-[1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-3-methyl-2-oxo-3-pyrrolidinyl]phenoxy]methyl]-5-methylphenyl]-4-morpholinecarboxamide

Beginning with the aniline from (48a) and 4-morpholinecarbonyl chloride, example 54 was prepared in an analogous series of reactions to example 49. MS found: $(M-H)^- = 509$.

Example 55

3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α,α ,3-trimethyl-2-oxo-1-pyrrolidineacetamide

(55a) Following a procedure analogous to step (1d), the

aldehyde from (1c) (1.50 g, 4.81 mmol) was reacted with α -aminoisobutyric acid methyl ester hydrochloride to give the lactam (396 mg, 22%). MS found: $(M+H)^+ = 382$.

(55b) Following a procedure analogous to step (3a), the lactam from (55a) (378 mg, 992 μ mol) was hydrogenolized to give the phenol (270 mg, 93%). MS found: $(M-H)^- = 290$.

(55c) Following a procedure analogous to step (6b), the phenol from (55b) (128 mg, 0.440 mmol) was reacted with 4-bromomethyl-2,6-dichloropyridine to give the picolyl ether (153 mg, 77%). MS found: $(M+Na)^+ = 473$.

(55d) The ester from (55c) was stirred in THF (3 mL) and 1 N NaOH (10 mL) at rt overnight. The mixture was acidified to pH 4 with 1 N HCl and THF removed in vacuo. The aqueous residue was extracted with ethyl acetate. The combined extracts were washed with brine, dried (MgSO₄) and concentrated to give the carboxylic acid (137 mg, 94%). MS found: $(M-H)^- = 435$.

(55e) Hunig's base (148 mg, 4 eq), hydroxylamine hydrochloride (40 mg, 2 eq) and BOP (152 mg, 1.2 eq) were added to the acid from (55d) (125 mg, 0.286 mmol) in DMF (5 mL) at 0 °C. the mixture was stirred at rt for 24 h and at 60 °C for 3 h. Sat ammonium chloride was added and the mixture extracted with ethyl acetate (2 x). The extracts were washed with sat NaHCO₃, water and brine, dried (MgSO₄) and concentrated.

Silica gel chromatography (methanol-chloroform, 8:92) provided the hydroxamic acid (50 mg, 39%). MS found: $(M+Na)^+ = 479$.

Example 56

5 [1(R)]-3-[1,1'-biphenyl]-4-yl-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

(56a) Triflic anhydride (1.45 mL, 2.2 eq) was added dropwise to a solution of the phenol from (6a) (1.09 g, 3.93 mmol) and 2,6-lutidine (1.01 mL, 2.2 eq) in CH_2Cl_2 (50 mL) at 0 °C.

10 After 10 min at this temperature, hexane (200 mL) was added. The mixture was filtered through a silica gel pad and the filter cake washed with ethyl acetate-hexane (1:1) until free of product. The filtrate was concentrated to give the triflate (1.49 g, 93%). MS found: $(M-H)^- = 408$.

15 (56b) A mixture of the triflate from (56a) (150 mg, 0.366 mmol), benzeneboronic acid (89.3 mg, 2 eq), triphenylphosphine (96 mg, 1 eq), potassium carbonate (202 mg, 4 eq) and anhydrous toluene (10 mL) was pumped then filled with nitrogen for 10 cycles to remove oxygen. Palladium(II) acetate (16.4 mg, 0.2 eq) was then quickly added and the flask was again deoxygenated for 10 cycles. This mixture was heated to reflux for 18 h. Following addition of ethyl acetate, the mixture was washed with water (2 x), brine, dried ($MgSO_4$) and concentrated. Silica gel chromatography (ethyl acetate-hexane, 25:75 then 50:50) give the biphenyl (118 mg, 96%). MS found: $(M+Na)^+ = 360$.

20 (56c) Following a procedure analogous to step (1f), the ester from (56b) (100 mg, 0.297 mmol) was reacted with hydroxylamine to give the hydroxamic acid (52 mg, 52%). MS found: $(M+H)^+ =$
30 339.

Example 57

[1(R)]-N-hydroxy- α ,3-dimethyl-3-(2'-methyl[1,1'-biphenyl]-4-yl)-2-oxo-1-pyrrolidineacetamide

35 Beginning with the triflate from (56a) and 2-methylbenzeneboronic acid, example 57 was prepared in an analogous series of reactions to (56b) and (56c). MS found: $(M+H)^+ = 353$.

Example 58**[1(R)]-N-hydroxy- α ,3-dimethyl-3-(4'-methyl[1,1'-biphenyl]-4-yl)-2-oxo-1-pyrrolidineacetamide**

Beginning with the triflate from (56a) and 4-methylbenzeneboronic acid, example 58 was prepared in an analogous series of reactions to (56b) and (56c). MS found: $(M+H)^+ = 353$.

Example 59**[1(R)-3-(3',4'-dimethoxy[1,1'-biphenyl]-4-yl)-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide**

Beginning with the triflate from (56a) and 3,4-dimethoxybenzeneboronic acid, example 59 was prepared in an analogous series of reactions to (56b) and (56c). MS found: $(M-H)^- = 397$.

Example 60**[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[2'-(trifluoromethyl)[1,1'-biphenyl]-4-yl]-1-pyrrolidineacetamide**

Beginning with the triflate from (56a) and 2-trifluoromethylbenzeneboronic acid, example 60 was prepared in an analogous series of reactions to (56b) and (56c). MS found: $(M-H)^- = 405$.

Example 61**[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-(4-methylphenoxy)phenyl]-2-oxo-1-pyrrolidineacetamide**

(61a) Copper(II) acetate monohydrate (108 mg, 1 eq), p-tolueneboronic acid (147 mg, 1 eq), and 4 Å molecular sieve (400 mg) were added sequentially to the phenol from (6a) (150 mg, 0.541 mmol) and pyridine (0.219 mL, 5 eq) in dichloromethane. The resultant mixture was stirred at rt open to atmosphere for 20 h. The mixture was filtered through a silica gel pad and the filter cake washed with ethyl acetate until free of product. The filtrate was concentrated and purified by silica gel chromatography (ethyl acetate-hexane,

30:70 then 40:60) to give the phenyl ether (167.4 mg, 84%).

MS found: $(M+Na)^+ = 390$.

(61b) Following a procedure analogous to step (1f), the ester from (61a) (154 mg, 0.419 mmol) was reacted with hydroxylamine to give the hydroxamic acid (144 mg, 93%). MS found: $(M-H)^- = 367$.

Example 62

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-(4-phenoxyphenyl)-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and benzeneboronic acid, example 62 was prepared in an analogous series of reactions to (61a) and (61b). MS found: $(M-H)^- = 353$.

Example 63

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-(2-methylphenoxy)phenyl]-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 2-methylbenzeneboronic acid, example 63 was prepared in an analogous series of reactions to (61a) and (61b). MS found: $(M-H)^- = 367$.

Example 64

[1(R)]-3-[4-(3,5-dichlorophenoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3,5-dichlorobenzeneboronic acid, example 64 was prepared in an analogous series of reactions to (61a) and (61b). MS found: $(M-H)^- = 421$.

Example 65

[1(R)]-3-[4-(3,4-dimethoxyphenoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3,4-dimethoxybenzeneboronic acid, example 65 was prepared in an analogous series of reactions to (61a) and (61b). MS found: $(M-H)^- = 413$.

Example 66

[1(R)]-3-[4-(1,3-benzodioxol-5-yloxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3,4-methylenedioxybenzeneboronic acid, example 66 was prepared in an analogous series of reactions to (61a) and (61b). MS found: $(M-H)^- = 397$.

Example 67

10 [1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-[3-(1-methylethyl)phenoxy]phenyl]-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3-isopropylbenzeneboronic acid, example 67 was prepared in an analogous series of reactions to (61a) and (61b). MS found: $(M-H)^- = 395$.

Example 68

20 [1(R)]-N-hydroxy-3-[4-(3-methoxyphenoxy)phenyl]- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3-methoxybenzeneboronic acid, example 68 was prepared in an analogous series of reactions to (61a) and (61b). MS found: $(M-H)^- = 383$.

Example 69

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-(3-thienyloxy)phenyl]-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and thiophene-3-boronic acid, example 69 was prepared in an analogous series of reactions to (61a) and (61b). MS found: $(M-H)^- = 359$.

Example 70

35 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-(3,4,5-trimethoxyphenoxy)phenyl]-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3,4,5-trimethoxybenzeneboronic acid, example 70 was prepared in an

analogous series of reactions to (61a) and (61b). MS found:
(M-H)⁻ = 443.

Example 71

5 [1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-
N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3,5-bis(trifluoromethyl)benzeneboronic acid, example 71 was
10 prepared in an analogous series of reactions to (61a) and (61b). MS found: (M+H)⁺ = 491.

Example 72

15 [1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-(1-
naphthalenyloxy)phenyl]-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 1-naphthaleneboronic acid, example 72 was prepared in an
analogous series of reactions to (61a) and (61b). MS found:
20 (M+H)⁺ = 405.

Example 73

[1(R)]-N-hydroxy-3-[4-[3-
[(hydroxyimino)methyl]phenoxy]phenyl]- α ,3-dimethyl-2-
25 oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3-formylbenzeneboronic acid, example 73 was prepared in an
analogous series of reactions to (61a) and (61b). MS found:
30 (M+H)⁺ = 398.

Example 74

[1(R)]-N-hydroxy-3-[4-[4-[1-
(hydroxyimino)ethyl]phenoxy]phenyl]- α ,3-dimethyl-2-
35 oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 4-acetylbenzeneboronic acid, example 74 was prepared in an

analogous series of reactions to (61a) and (61b). MS found:
 $(M-H)^- = 410$.

Example 75

[1(R)]-3-[4-([1,1'-biphenyl]-4-yloxy)phenyl]-N-
hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 4-biphenylboronic
acid, example 75 was prepared in an analogous series of
10 reactions to (61a) and (61b). MS found: $(M+H)^+ = 431$.

Example 76

[1(R)]-3-[4-(3,5-dibromophenoxy)phenyl]-N-hydroxy- α ,3-
dimethyl-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (6a) and 3,5-dibromobenzeneboronic acid, example 76 was prepared in an analogous series of reactions to (61a) and (61b). MS found: $(M+H)^+ = 510$.

Example 77

[1(R)]-3-[4-[3-(acetylamino)phenoxy]phenyl]-N-hydroxy-
α,3-dimethyl-2-oxo-1-pyrrolidineacetamide

25 Beginning with the phenol from (6a) and 3-acetamidobenzeneboronic acid, example 77 was prepared in an analogous series of reactions to (61a) and (61b). MS found: $(M+H)^+ = 412$.

Example 78

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-(4-nitrophenoxy)phenyl]-2-oxo-1-pyrrolidineacetamide
(78a) Cesium carbonate (254 mg, 1.8 eq) was added to the phenol from (6a) (120 mg, 0.433 mmol) and 1-fluoro-4-nitrobenzene (122 mg, 2 eq) in DMSO (2 mL). After 1 h at rt, sat ammonium chloride (3 mL) and ethyl acetate (100 mL) were added. The mixture was washed with water (2x5 mL), brine (5 mL), dried (MgSO₄) and concentrated. Silica gel

chromatography (ethyl acetate-hexane, 50:50) gave the phenyl ether (139.7 mg, 81%). MS found: $(M+H)^+ = 399$.

(78b) Following a procedure analogous to step (1f), the ester from (78a) (125 mg, 0.314 mmol) was reacted with hydroxylamine to give the hydroxamic acid (80.6 mg, 64%). MS found: $(M-H)^- = 398$.

Example 79

[1(R)]-N-hydroxy- α ,3-dimethyl-3-(4-methylphenyl)-2-oxo-1-pyrrolidineacetamide

Beginning with methyl (4-methylphenyl)acetate, example 79 was prepared in an analogous series of reactions to example 1. MS found: $(M-H)^- = 275$.

Example 80

[1(R)]-3-[4-[[2,6-dimethyl-4-pyridinyl]oxylmethyl]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(80a-d) Beginning with methyl (4-methylphenyl)acetate, methyl (R)- α ,3-dimethyl-2-oxo-3-(4-methyl phenyl)-1-pyrrolidineacetate was prepared in an analogous series of reactions to (1a-d). The two isomers were separated by silica gel chromatography (ethyl acetate-hexane, 20:80 then 25:75). The more polar isomer was used for subsequent reactions. MS found: $(M+H)^+ = 276$.

(80e) N-bromosuccinimide (1.45 g, 1.05 eq) and benzoyl peroxide (28.2 mg, 0.015 eq) were added to the more polar ester from (80d) (2.14 g, 7.77 mmol) in carbon tetrachloride (50 mL). The suspension was stirred under two 250 W sun lamp radiation for 2 h. The mixture was concentrated and purified by silica gel chromatography (ethyl acetate-hexane, 20:80 then 30:70) to give the bromide (1.784 g, 65%). MS found: $(M+H)^+ = 354$.

(80f) Cesium carbonate (199 mg, 1.8 eq) was added to the bromide from (80e) (120 mg, 0.339 mmol) and 2,6-dimethyl-4-phenol (83 mg, 2 eq) in DMSO (4 mL). After 3 h at rt, sat ammonium chloride was added. The mixture was extracted with ethyl acetate (3 x). The combined extracts were washed with

brine, dried (MgSO₄) and concentrated. Silica gel chromatography (methanol-chloroform, 7:93) gave the pyridinyl ether (35 mg, 26%). MS found: (M+H)⁺ = 397.

(80g) Following a procedure analogous to step (1f), the ester from (80f) (30 mg, 0.0758 mmol) was reacted with hydroxylamine. The hydroxamic acid was isolated as a TFA salt (15 mg, 39%). MS found: (M+H)⁺ = 398.

Example 81

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-[(4-quinolinylloxy)methyl]phenyl]-1-pyrrolidineacetamide mono(trifluoroacetate)

Beginning with the bromide from (80e) and 4-hydroxyquinoline, example 81 was prepared in an analogous series of reactions to (80f) and (80g). MS found: (M+H)⁺ = 420.

Example 82

[1(R)]-N-hydroxy- α ,3-dimethyl-3-(4-nitrophenyl)-2-oxo-1-pyrrolidineacetamide

(82a) DBU (25.33 mL, 1.1 eq) was added dropwise to a mixture of 2-(4-nitrophenyl)propionic acid (30.00 g, 154 mmol) and iodomethane (10.55 mL, 1.1 eq) in toluene (250 mL). After 30 min at rt, ether (200 mL) was added. The mixture was filtered through a silica gel pad and the filter cake washed with ethyl acetate-hexane (1:1) until free of solvent. The combined filtrate was concentrated to give the ester (25.85 g, 80%). MS found: M⁺ = 209.

(82b) Sodium hydride (2.76 g, 1.2 eq, 60% in mineral oil) was added to the ester from (82a) (12.00 g, 57.4 mmol) and allyl bromide (9.93 mL, 2 eq) in DMF (200 mL) at 0 °C. After 30 min at rt, sat NH₄Cl (200 mL) was added and the mixture was concentrated to dryness in vacuo. The solid was treated with water (200 mL) and extracted with ether (3x200 mL). The combined extracts were washed with water, brine, dried (MgSO₄) and concentrated. The crude material was used in the next step without purification.

(82c) A 1 N solution of NaOH (100 mL) was added to half of the crude material from (82b) in methanol (200 mL). The mixture was stirred at rt overnight and at reflux for 1 h. Following removal of methanol in vacuo, the aqueous residue was washed with hexane (2x100 mL) to remove mineral oil. The combined hexane washings were back extracted with 1 N NaOH (30 mL).

The combined aqueous layer was acidified with 1 N HCl (180 mL), saturated with solid NaCl, and extracted with ethyl acetate (3x250 mL). The combined organic extracts were washed with brine (30 mL), dried (MgSO₄) and concentrated to give the carboxylic acid (6.38 g, 94% for 2 steps).

(82d) HATU (11.17 g, 1.1 eq) and NMM (10.27 mL, 3.5 eq) were added to the acid from (82c) (6.28 g, 26.7 mmol) and D-alanine methyl ester hydrochloride (4.10 g, 1.1 eq) in DMF (50 mL).

After 2 h at rt, ethyl acetate (750 mL) was added. The mixture was washed with 1 N HCl (3x50 mL), water (50 mL), sat NaHCO₃ (2x50 mL), water (50 mL), and brine (50 mL), dried (MgSO₄) and concentrated. The crude material was used in the next step without purification. MS found: $(M+H)^+ = 321$.

(82e) Ozone was bubbled through a solution of the crude olefin from (82d) in dichloromethane (200 mL) and methanol (100 mL) at -78 °C until starting material consumed. the mixture was purged with oxygen and treated with triphenylphosphine (7.00 g, 1.0 eq). After 1 h at rt, the mixture was concentrated.

The crude material was used in the next step without purification.

(82f) Triethylsilane (42.6 mL, 10 eq) and trifluoroacetic acid (20.6 mL, 10 eq) were added successively to the crude aldehyde from (82e) in dichloromethane at 0 °C. After 2 h at rt, the mixture was concentrated and purified by silica gel chromatography (ethyl acetate-toluene-hexane, 20:10:70 then 25:10:65 then 30:10:60 then 35:10:55) to give less polar lactam (2.211 mg), more polar lactam (2.184 g), and a 1:1 mixture of the two isomers (0.44 g). The total yield of the two isomers is 4.835 g (59% for three steps). MS found: $(M+H)^+ = 307$.

(82g) Following a procedure analogous to step (1f), the more polar ester from (82f) (100 mg, 0.326 mmol) was reacted with

hydroxylamine to give the hydroxamic acid (93.8 mg, 94%). MS found: $(M-H)^- = 306$.

Example 83

5 **[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-
[(phenylcarbonyl)amino]phenyl]-1-pyrrolidineacetamide**

(83a) The more polar isomer from (82f) (1.97 g, 6.43 mmol) and 10% Pd on carbon (0.5 g) in methanol (50 mL) and chloroform
10 (50 mL) was stirred under balloon pressure hydrogen for 2 h. Following removal of catalyst by filtration, the filtrate was concentrated to give the aniline (1.83 g, 100%). MS found: $(M+H)^+ = 277$.

(83b) Following a procedure analogous to step (49a), the
15 aniline from (83a) (100 mg, 0.362 mmol) was reacted with benzoyl chloride to give the benzamide (124 mg, 90%). MS found: $(M+Na)^+ = 403$.

(83c) Following a procedure analogous to step (1f), the
20 benzamide from (83b) (110 mg, 0.289 mmol) was reacted with hydroxylamine to give the hydroxamic acid (100 mg, 91%). MS found: $(M-H)^- = 380$.

Example 84

25 **[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-
[(phenylsulfonyl)amino]phenyl]-1-pyrrolidineacetamide**

Beginning with the aniline from (83b) and benzenesulfonyl chloride, example 84 was prepared in an analogous series of reactions to (49a) and (1f). MS found: $(M+Na)^+ = 440$.

Example 85

30 **[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-
[(phenylamino)carbonyl]amino]phenyl]-1-
pyrrolidineacetamide**

Beginning with the aniline from (83b) and phenyl
35 isocyanate, example 85 was prepared in an analogous series of reactions to (49a) and (1f). MS found: $(M+Na)^+ = 419$.

Example 86**[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-[(1-naphthalenylmethyl)aminolphenyl]-2-oxo-1-pyrrolidineacetamide**

- 5 (86a) Hunig's base (0.13 mL, 2 eq), 1-naphthaldehyde (62.2 mg, 1.1 eq) and 4 A molecular sieves (300 mg) were added to the aniline from (83a) (100 mg, 0.362 mmol) in 1,2-dichloroethane (3 mL). After 30 min at rt, NaBH(OAc)₃ (230 mg, 3 eq) was added and the mixture was stirred for 36 h. The precipitate
10 was removed by filtration. The filtrate was concentrated and purified by silica gel chromatography (ethyl acetate-hexane, 50:50) to give the secondary amine (117 mg, 78%). MS found: (M+Na)⁺ = 439.
- (86b) Following a procedure analogous to step (1f), the ester
15 from (86a) (108 mg, 0.260 mmol) was reacted with hydroxylamine to give the hydroxamic acid (75.4 mg, 70%). MS found: (M+Na)⁺ = 440.

Example 87

- 20 **[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-[(4-quinolinylmethyl)aminolphenyl]-1-pyrrolidineacetamide**

- Beginning with the aniline from (83b) and quinoline-4-carboxaldehyde, example 87 was prepared in an analogous series
25 of reactions to (86a) and (1f). MS found: (M+H)⁺ = 419.

Example 88

- 30 **[1(R)]-3-[4-[(3,5-dimethoxyphenyl)methyl]aminolphenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide**

- Beginning with the aniline from (83b) and 3,5-dimethoxybenzaldehyde, example 88 was prepared in an analogous series of reactions to (86a) and (1f). MS found: (M-H)⁻ =
35 426.

Example 89**3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide**

Beginning with the aldehyde from (1c) and glycine methyl ester hydrochloride, example 89 was prepared in an analogous series of reactions to (1d), (3a), (6b) and (1f), but using 3,5-dimethylbenzyl bromide in step (6b). MS found: $(M+Na)^+ = 405$.

Example 90**3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide**

Beginning with the aldehyde from (1c) and glycine methyl ester hydrochloride, example 90 was prepared in an analogous series of reactions to (1d), (3a), (6b) and (1f), but using 4-bromomethyl-2,6-dichloropyridine in step (6b). MS found: $(M+H)^+ = 424$.

Example 91**3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide**

Beginning with the aldehyde from (1c) and glycine methyl ester hydrochloride, example 91 was prepared in an analogous series of reactions to (1d), (3a), (6b) and (1f), but using 4-bromomethyl-2,6-dimethylpyridine hydrochloride in step (6b). MS found: $(M+H)^+ = 424$.

Example 92**[1(R)]-N-hydroxy-3-methyl- α -(1-methylethyl)-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide mono(trifluoroacetate)**

(92a) Following a procedure analogous to step (1d), the aldehyde from (1c) (3.00 g, 9.61 mmol) was reacted with D-valine methyl ester hydrochloride to give the lactam as mixture of two isomers. Silica gel chromatography (ether-hexane, 50:50 then 85:15) provided the less polar isomer (1.25 g, 30%). MS found: $(M+Na)^+ = 418$.

(92b) Following a procedure analogous to step (3a), the less polar lactam from (92a) (1.25 g, 3.18 mmol) was hydrogenolized to give the phenol (0.915 g, 94%). MS found: $(M+H)^+ = 300$.

(92c) Following a procedure analogous to step (6b), the phenol from (92b) (106 mg, 0.348 mmol) was reacted with 4-chloromethylquinoline to give the phenyl ether (134 mg, 86%).

MS found: $(M+H)^+ = 447$.

(92d) The 1.76 M NH_2OH/KOH solution in methanol was prepared fresh following the procedure described in (1e). The ester from (92c) (134 mg, 0.300 mmol) was treated with the hydroxylamine solution (3.4 mL, 20 eq). Additional hydroxylamine (2 mL, 0.5 mL and 2 mL) were added after 20 min, 40 min and 1.5 h, respectively. After a total of 2 h, the mixture was neutralized to pH 7 with 1 N HCl and concentrated. Purification by HPLC (acetonitrile-water-TFA, 15:85:0.1 to 50:50:0.1) provided the hydroxamic acid as a TFA salt (69 mg, 41%). MS found: $(M-H)^- = 446$.

Example 93

[1(R)]-N-hydroxy-3-methyl- α -(1-methylethyl)-2-oxo-3-[4-(phenylmethoxy)phenyl]-1-pyrrolidineacetamide

Following a procedure analogous to step (1f), the less polar lactam from (92a) was reacted with hydroxylamine to give the hydroxamic acid. MS found: $(M-H)^- = 395$.

Example 94

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl- α -(1-methylethyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

Beginning with the phenol from (92b) and 4-chloromethyl-2,6-dimethylpyridine, example 94 was prepared in an analogous series of reactions to (6b) and (92d). MS found: $(M+H)^+ = 426$.

Example 95**[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide**

- 5 (95a) Following a procedure analogous to step (1d), the aldehyde from (1c) (3.00 g, 9.61 mmol) was reacted with D-leucine methyl ester hydrochloride to give the lactam as mixture of two isomers. Silica gel chromatography (ether-toluene, 10:90) provided the less polar isomer (1.20 g, 31%).
- 10 MS found: $(M+Na)^+ = 432$.
- (95b) Following a procedure analogous to step (3a), the less polar lactam from (95a) (1.20 g, 2.93 mmol) was hydrogenolized to give the phenol (0.94 g, 100%). MS found: $(M+H)^+ = 320$.
- (95c) Following a procedure analogous to step (6b), the phenol
- 15 from (95b) (155 mg, 0.486 mmol) was reacted with 4-chloromethyl-2,6-dimethylpyridine to give the phenyl ether (191 mg, 90%). MS found: $(M+H)^+ = 439$.
- (95d) Following a procedure analogous to step (1f), the ester from (95c) (140 mg, 0.320 mmol) was reacted with hydroxylamine
- 20 to give the hydroxamic acid (115 mg, 82%). MS found: $(M+H)^+ = 440$.

Example 96**[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide**

- 25 Beginning with the phenol from (95b) and 4-bromomethyl-2,6-dichloropyridine, example 96 was prepared in an analogous series of reactions to (6b) and (1f). MS found: $(M-H)^- =$
- 30 479.

Example 97**[1(R)]-3-[4-[[3,5-bis(trifluoromethyl)phenyl]methoxy]phenyl]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide**

- 35 Beginning with the phenol from (95b) and 3,5-bis(trifluoromethyl)benzyl bromide, example 97 was prepared in

an analogous series of reactions to (6b) and (1f). MS found:
(M-H)⁻ = 454.

Example 98

5 [1(R)]-3-[4-[(3,5-dichlorophenyl)methoxy]phenyl]-N-
 hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-1-
 pyrrolidineacetamide

Beginning with the phenol from (95b) and 3,5-dichlorobenzyl bromide, example 98 was prepared in an analogous series of reactions to (6b) and (1f). MS found:
10 (M+H)⁺ = 479.

Example 99

15 [1(R)]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-3-
 [3-(phenylmethoxy)propyl]-1-pyrrolidineacetamide

(99a) Following a procedure analogous to step (1a), ethyl 2-methyl-4-pentenoate (3.00 g, 21.1 mmol) was reacted with 3-benzyloxy-1-bromopropane to give the crude ester. MS found:
20 (M+NH₄)⁺ = 308.

(99b) Following a procedure analogous to step (1c), the crude ester from (99a) was ozonolized to give the aldehyde (5.19 g, 84% for 2 steps). MS found: (M+NH₄)⁺ = 310.

(99c) Following a procedure analogous to step (1d), the
25 aldehyde from (99b) (5.06 g, 17.3 mmol) was reacted with D-leucine methyl ester hydrochloride to give the lactam as mixture of two isomers. Silica gel chromatography (ethyl acetate-hexane, 20:80 then 25:75 then 30:70) provided the less polar isomer (1.94 g), the more polar isomer (1.66 g) and a
30 1:1.1 mixture of both isomers (1.86 g). The total yield of both isomers is 5.46 g (84%). MS found: (M+H)⁺ = 376.

(99d) Following a procedure analogous to step (1f), the less polar lactam from (99c) (100 mg, 0.266 mmol) was reacted with hydroxylamine to give the hydroxamic acid (80.6 mg, 80%). MS
35 found: (M-H)⁻ = 375.

(99e) Following a procedure analogous to step (1f), the more polar lactam from (99c) (100 mg, 0.266 mmol) was reacted with

hydroxylamine to give the hydroxamic acid (81.8 mg, 82%). MS found: $(M-H)^- = 375$.

Example 101

5 [1(R)]-N-hydroxy-3-methyl-3-[2-methyl-4-(phenylmethoxy)phenyl]- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide

10 (101a) Following a procedure analogous to step (1a), methyl (4-benzyloxy-2-methylphenyl)acetate (5.00 g, 18.5 mmol) was reacted with iodomethane to give the crude ester. MS found: $(M+NH_4)^+ = 302$.

15 (101b) Following a procedure analogous to step (1b), the crude material from (101a) was reacted with allyl bromide to give the crude ester. MS found: $(M+NH_4)^+ = 342$.

20 (101c) Following a procedure analogous to step (1c), the crude ester from (101b) was ozonolized to give the aldehyde (5.42 g, 90% for 3 steps). MS found: $(M+NH_4)^+ = 344$.

25 (101d) Following a procedure analogous to step (1d), the aldehyde from (101c) (5.28 g, 16.2 mmol) was reacted with D-leucine methyl ester hydrochloride to give the lactam as mixture of two isomers. Silica gel chromatography (ethyl acetate-hexane, 20:80) provided the less polar isomer (1.363 g) and the more polar isomer (1.412 g). MS found: $(M+Na)^+ = 446$.

30 (101e) Following a procedure analogous to step (1f), the less polar lactam from (101d) (100 mg, 0.262 mmol) was reacted with hydroxylamine to give the hydroxamic acid (65.2 mg, 65%). MS found: $(M-H)^- = 423$.

Example 102

35 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxyl]-2-methylphenyl]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide

(102a) Following a procedure analogous to step (3a), the less polar lactam from (101d) (1.05 g, 2.48 mmol) was

hydrogenolized to give the phenol (731 mg, 88%). MS found:
(M-H)⁻ = 332.

(102b) Following a procedure analogous to step (6b), the phenol from (102a) (100 mg, 0.300 mmol) was reacted with 4-bromomethyl-2,6-dichloropyridine to give the picolyl ether (116 mg, 78%). MS found: (M+Na)⁺ = 515.

(102c) Following a procedure analogous to step (1f), the ester from (102b) (105 mg, 0.213 mmol) was reacted with hydroxylamine to give the hydroxamic acid (70.2 mg, 67%). MS found: (M-H)⁻ = 492.

Example 103

[1(R)]-N-hydroxy-3-methyl-3-[2-methyl-4-(2-naphthalenylmethoxy)phenyl]- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (102a) and 1-bromomethylnaphthlene, the desired product was prepared in an analogous series of reactions to (6b) and (1f). MS found: (M+H)⁺ = 475.

Example 104

[1(R)]-N-hydroxy-3-methyl- α -(2-methylpropyl)-3-[2-methyl-4-(4-pyridinylmethoxy)phenyl]-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (102a) and 4-chloromethylpyridine, example 104 was prepared in an analogous series of reactions to (6b) and (1f). MS found: (M+H)⁺ = 426.

Example 105

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]-2-methylphenyl]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (102a) and 4-chloromethyl-2,6-dimethylpyridine, example 105 was prepared in an analogous series of reactions to (6b) and (1f). MS found: (M+H)⁺ = 454.

Example 106

[1(R)]-N-hydroxy-3-methyl- α -[2-(methylthio)ethyl]-2-oxo-3-[4-(phenylmethoxy)phenyl]-1-pyrrolidineacetamide

(106a) Following a procedure analogous to step (1d), the aldehyde from (1c) (4.19 g, 13.4 mmol) was reacted with D-methionine methyl ester hydrochloride to give the lactam as a 1:1 mixture of two isomers (4.39 g, 77%). MS found: $(M+H)^+ = 428$.

(106b) Following a procedure analogous to step (1f), the lactam from (106a) (144 mg, 0.337 mmol) was reacted with hydroxylamine to give the hydroxamic acid (90.7 mg, 63%). MS found: $(M-H)^- = 427$.

Example 107

[1(R)]-3-[4-(3,5-dibromophenoxy)phenyl]-3-methyl- α -[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetic acid

(107a) Oxone (19.0 g, 3 eq) in water (100 mL) was added to the lactam from (106a) (8.80 g, 20.6 mmol) in methanol (100 mL) at 0 °C. After 30 min at 0 °C and 4 h at rt, methanol was removed in vacuo. The aqueous residue was diluted with water (300 mL) and extracted with chloroform (3x400 mL). The combined organic extracts were washed with water (50 mL) and brine (50 mL), dried (MgSO₄) and concentrated. Silica gel chromatography (ethyl acetate-hexane, 60:40 then 70:30 then 100:0) provided the more polar sulfone (2.88 g, 30%). MS found: $(M+Na)^+ = 482$.

(107b) Following a procedure analogous to step (3a), the sulfone from (107a) (2.88 g, 6.27 mmol) was hydrogenolized to give the phenol (2.15 g, 93%). MS found: $(M+H)^+ = 370$.

(107c) Following a procedure analogous to step (61a), the phenol from (107b) (120 mg, 0.325 mmol) was reacted with 3,5-dibromobenzenboronic acid to give the phenyl ether (150 mg, 77%). MS found: $(M+H)^+ = 604$.

(107d) A 1 N solution of LiOH (0.28 mL, 1.3 eq) was added to the ester from (107c) (128 mg, 0.212 mmol) in THF (1.5 mL) at 0 °C. After 30 min at this temperature, the mixture was acidified to pH 2-3. The mixture was concentrated to dryness, treated with ethyl acetate (100 mL), and filtered. The

filtrate was concentrated to give the carboxylic acid (121 mg, 97%). MS found: $(M-H)^- = 492$.

Example 108

5 [1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-
N-hydroxy-3-methyl- α -[2-(methylsulfonyl)ethyl]-2-oxo-
1-pyrrolidineacetamide

Beginning with the phenol from (107b) and 3,5-bis(trifluoromethyl)benzene boronic acid, example 108 was
10 prepared in an analogous series of reactions to (61a) and (1f). MS found: $(M-H)^- = 581$.

Example 109

15 [1(R)]-3-[4-(3,5-dibromophenoxy)phenyl]-N-hydroxy-3-
methyl- α -[2-(methylsulfonyl)ethyl]-2-oxo-1-
pyrrolidineacetamide

Following a procedure analogous to step (1f), the lactam from (107c) (156 mg, 0.259 mmol) was reacted with
hydroxylamine to give the hydroxamic acid (110 mg, 70%). MS
20 found: $(M-H)^- = 603$.

Example 110

[1(R)]-3-[4-[(2,6-dichloro-4-
pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl- α -[2-
25 (methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (107b) and 4-bromomethyl-2,6-dichloropyridine, example 110 was prepared in an analogous series of reactions to (6b) and (1f). MS found: $(M-H)^- =$
528.

30

Example 111

[1(R)]-3-[4-[(2,6-dimethyl-4-
pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl- α -[2-
35 (methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide

Beginning with the phenol from (107b) and 4-chloromethyl-2,6-dimethylpyridine, example 111 was prepared in an analogous series of reactions to (6b) and (1f). MS found: $(M+H)^+ =$
490.

Example 112

[1(R)]-N-hydroxy-3-methyl- α -(2-(methylsulfonyl)ethyl)-
2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-
5 pyrrolidineacetamide mono(trifluoroacetate)

Beginning with the phenol from (107b) and 4-chloromethylquinoline hydrochloride, example 112 was prepared in an analogous series of reactions to (6b) and (1f). MS found: $(M+H)^+ = 512$.

Example 113

N-hydroxy-1-[3-methyl-2-oxo-3-[4-
(phenylmethoxy)phenyl]-1-
10 pyrrolidinyl]cyclopropanecarboxamide

15 (113a) Following a procedure analogous to step (1d), the aldehyde from (1c) (400 mg, 1.28 mmol) was reacted with 1-aminocyclopropane-1-carboxylic acid methyl ester hydrochloride to give the lactam (280 mg, 58%). MS found: $(M+H)^+ = 380$.

(113b) Following a procedure analogous to step (1f), the ester
20 from (113a) (100 mg, 0.264 mmol) was reacted with hydroxylamine to give the hydroxamic acid (76 mg, 76%). MS found: $(M-H)^- = 379$.

Example 114

25 [1(R)]-N-hydroxy- α -(4-hydroxyphenyl)methyl]-3-methyl-
2-oxo-3-[4-(phenylmethoxy)phenyl]-1-
pyrrolidineacetamide

Beginning with the aldehyde from (1c) and D-tyrosine methyl ester hydrochloride, example 114 was prepared in an
30 analogous series of reactions to (1d) and (1f). MS found: $(M-H)^- = 395$.

Example 115

[1(R)]-3-[4-[(2,6-dichloro-4-
35 pyridinyl)methoxy]phenyl]-N-hydroxy- α -(2-
hydroxyethyl)-3-methyl-2-oxo-1-pyrrolidineacetamide

(115a) A mixture of D-homoserine (25.00 g, 210 mmol), 35-37% hydrochloric acid (200 mL) and water (200 mL) was heated to

reflux for 3 h. Removal of solvent in vacuo provided the aminolactone hydrochloride (27.68 g, 96%). MS found: $(M+NH_4)^+ = 119$.

(115b) Following a procedure analogous to step (1d), the aldehyde from (1c) (3.00 g, 9.60 mmol) was reacted with the aminolactone hydrochloride from (115a) (1.45 g, 1.1 eq) to give the lactam as mixture of two isomers. Silica gel chromatography (ethyl acetate-hexane, 20:80) provided the less polar isomer (1.51 g) and the more polar isomer (1.45 g). MS found: $(M+NH_4)^+ = 383$.

(115c) Following a procedure analogous to step (3a), the more polar lactam from (115b) (1.40 g, 3.83 mmol) was hydrogenolized to give the phenol (1.06 g, 100%). MS found: $(M+H)^+ = 276$.

(115d) Following a procedure analogous to step (6b), the phenol from (115c) (1.03 g, 3.74 mmol) was reacted with 4-bromomethyl-2,6-dichloropyridine to give the picolyl ether (1.36 g, 84%). MS found: $(M+H)^+ = 435$.

(115e) Following a procedure analogous to step (1f), the ester from (115d) (71.0 mg, 0.163 mmol) was reacted with hydroxylamine to give the hydroxamic acid (59.1 mg, 77%) as a 85:15 mixture due to partial epimerization. MS found: $(M-H)^- = 466$.

Example 116

[1(R)]-1,1-dimethylethyl [5-[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]carbamate

(116a) Following a procedure analogous to step (1d), the aldehyde from (1c) (5.05 g, 16.2 mmol) was reacted with H-D-Lys(BOC)-OMe hydrochloride (5.28 g, 1.1 eq) to give the crude lactam as mixture of two isomers. The BOC protecting group came off during the cyclization.

(116b) The crude material from (116a) in methylene chloride (100 mL) and DMF (10 mL) was treated with Hunig's base (12.0 mL, 2 eq) and di-*t*-butyl dicarbonate (8.33 g, 1.2 eq) for 1 h at rt. Following addition of sat ammonium chloride (50 mL) and ethyl acetate (800 mL), the mixture was washed with water

(2x50 mL), brine (50 mL), dried (MgSO₄) and concentrated. Silica gel chromatography (ethyl acetate-hexane, 40:60 then 50:50) gave the BOC protected lactams (5.49 g, 65% for 2 steps) as a 1:1 mixture. MS found: $(M+Na)^+ = 547$.

5 (116c) Following a procedure analogous to step (3a), the lactam from (116b) (5.40 g, 10.3 mmol) was hydrogenolized. Silica gel chromatography (isopropanol-chloroform, 3:97 then 5:95) gave more polar phenol (1.29 g), a 1:1 mixture of both isomers (1.46 g), as well as the less polar isomer. MS found:
10 $(M+Na)^+ = 457$.

(116d) Following a procedure analogous to step (6b), the more polar phenol from (116c) (300 mg, 0.690 mmol) was reacted with 4-bromomethyl-2,6-dichloropyridine to give the picolyl ether (360 mg, 88%). MS found: $(M+Na)^+ = 616$.

15 (116e) Following a procedure analogous to step (1f), the ester from (116d) (152 mg, 0.256 mmol) was reacted with hydroxylamine to give the hydroxamic acid (71.0 mg, 47%). MS found: $(M-H)^- = 593$.

20 Example 117

[1(R)]- α -(4-aminobutyl)-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

The hydroxamic acid example 116 (39 mg, 0.065 mmol) was
25 stirred with trifluoroacetic acid (0.5 mL) and CH₂Cl₂ (2 mL) for 1 h at rt and concentrated to give example 117 (40 mg, 100%). MS found: $(M+H)^+ = 495$.

Example 118

30 [1(R)]- α -[4-(acetylamino)butyl]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide

(118a) The picolyl ether from (116d) (351 mg, 0.590 mmol) was stirred with trifluoroacetic acid (2 mL) and CH₂Cl₂ (8 mL) for
35 2 h at rt and concentrated to give the free amine trifluoroacetate in quantitative yield. MS found: $(M+H)^+ = 494$.

(118b) Beginning with the amine from (118a) and acetyl chloride, example 118 was prepared in an analogous series of reactions to (49a) and (1f). MS found: $(M-H)^- = 535$.

5

Example 119

[1(R)]-N-[5-[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]-3-pyridineacetamide

10

Beginning with the amine from (118a) and nicotinoyl chloride, example 119 was prepared in an analogous series of reactions to (49a) and (1f). MS found: $(M+H)^+ = 600$.

Example 120

15

[1(R)]-N-[5-[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]-4-morpholinecarboxamide

20

Beginning with the amine from (118a) and 4-morpholinecarbonyl chloride, example 120 was prepared in an analogous series of reactions to (49a) and (1f). MS found: $(M+Na)^+ = 630$.

Example 121

25

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl- α -[4-[(methylsulfonyl)aminobutyl]-2-oxo-1-pyrrolidineacetamide

30

Beginning with the amine from (118a) and methanesulfonyl chloride, example 121 was prepared in an analogous series of reactions to (49a) and (1f). MS found: $(M+Na)^+ = 595$.

Example 122

35

[1(R)]- α -[4-(acetylamino)butyl]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide

(122a) Following a procedure analogous to step (6b), the more polar phenol from (116c) (1.00 g, 2.30 mmol) was reacted with

4-bromomethyl-2,6-dimethylpyridine to give the picolyl ether (1.00 g, 79%). MS found: $(M+H)^+ = 554$.

(122b) Following a procedure analogous to step (118a), the picolyl ether from (122a) (1.00 g, 1.81 mmol) was deprotected with trifluoroacetic acid to give the amine trifluoroacetate (1.28, 100%). MS found: $(M+H)^+ = 454$.

(122c) Beginning with the amine from (122b) and acetyl chloride, example 122 was prepared in an analogous series of reactions to (49a) and (1f). MS found: $(M+H)^+ = 497$.

Example 123

[1(R)]-1,1-dimethylethyl [5-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]carbamate

Beginning with the picolyl ether from (122a), example 123 was prepared in an analogous series of reactions to (55d) and (55e). MS found: $(M+H)^+ = 555$.

Example 124

[1(R)]- α -(4-aminobutyl)-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

Starting with the hydroxamic acid from example 123, example 124 was prepared in a procedure analogous to example 117. MS found: $(M+H)^+ = 455$.

Example 125

[1(R)]- α -[4-[(aminoacetyl)amino]butyl]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the amine from (122b) and N-(t-butoxycarbonyl)glycine, example 125 was prepared in an analogous series of reactions to (50a), (1e) and example 51. MS found: $(M+H)^+ = 512$.

Example 126

[1(R)]- α -[4-(acetylamino)butyl]-3-[4-[3,5-bis(trifluoromethyl)phenyl]methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide

Beginning with the more polar phenol from (116c) and 3,5-bis(trifluoromethyl)benzyl bromide, example 126 was prepared in an analogous series of reactions to (6b), (118a), (49a) and (1f). MS found: $(M+Na)^+ = 626$.

Example 127

[1(R)]-1,1-dimethylethyl [5-[3-[4-(3,5-dibromophenoxy)phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]carbamate

Beginning with the more polar phenol from (116c) and 3,5-dibromobenzeneboronic acid, example 127 was prepared in an analogous series of reactions to (61a) and (1f). MS found: $(M-H)^- = 668$.

Example 128

[1(R)]- α -(4-aminobutyl)-3-[4-(3,5-dibromophenoxy)phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

Starting with the hydroxamic acid from example 127, example 128 was prepared in a procedure analogous to example 117. MS found: $(M+H)^+ = 570$.

Example 129

[1(R)]-1,1-dimethylethyl [3-[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-4-(hydroxyamino)-4-oxobutyl]carbamate

(129a) Iodobenzene diacetate (38.6 g, 1.2 eq) was added to a mixture of Z-D-Gln-OH (28.1 g, 100 mmol), ethyl acetate (134 mL), acetonitrile (134 mL) and water (67 mL) at 5-10 °C. After 30 min at 10 °C and 4 h at 16 °C, the organic solvent was removed in vacuo. The aqueous residue was washed with ethyl acetate (2x20 mL) and concentrated to small volume. The product was precipitated out by addition of ethyl acetate (100 mL). Filtration and washing with ethyl acetate (50 mL)

provided the diamino acid (16.3 g, 64.5%). MS found: $(M+H)^+$ = 253.

(129b) Following a procedure analogous to (82a), the diamino acid from (129a) (5.40 g, 21.4 mmol) was cyclized with BOP reagent to give the lactam (2.33 g, 47%). MS found: $(M+Na)^+$ = 257.

(129c) Following a procedure analogous to (3a), the lactam from (129b) (9.10 g, 38.8 mmol) was hydrogenolized to give the free aminolactam hydrochloride (5.33 g, 100%). MS found: $(M+NH_4)^+$ = 118.

(129d) Following a procedure analogous to (1d), the aldehyde from (1c) (2.39 g, 7.65 mmol) and the lactam from (129c) (1.3 eq) were converted to the lactam (2.29 g, 82%) as a 1:1 mixture of two isomers. MS found: $(M+Na)^+$ = 387.

(129e) Following a procedure analogous to (3a), the lactam from (129d) (2.23 g, 6.12 mmol) was hydrogenolized to give the phenol (1.60 g, 95%). MS found: $(M+H)^+$ = 275.

(129f) Following a procedure analogous to (6b), the phenol from (129e) (1.51 g, 5.50 mmol) was coupled with 4-bromomethyl-2,6-dichloropyridine to give the picolyl ether (1.03 g, 43%). MS found: $(M+Na)^+$ = 456.

(129g) Triethylamine (0.32 mL, 1 eq), (BOC)₂O (1.00 g, 2 eq) and DMAP (0.281 g, 1 eq) were added to the lactam from (129f) (1.00 g, 2.30 mmol) in dichloromethane (10 mL) and the mixture was stirred at rt overnight. The solvent was removed and the mixture purified by silica gel chromatography (ethyl acetate-hexane, 40:60 then 50:50 then 60:40) to provide the less polar isomer (380 mg) and the more polar isomer (310 mg). MS found: $(M+Na)^+$ = 556.

(129h) Following a procedure analogous to (1f), the more polar lactam from (129g) (102 mg, 0.191 mmol) was converted to the hydroxamic acid (50.0 mg, 50%). MS found: $(M-H)^-$ = 565.

Example 130

[1(R)]- α -(2-aminoethyl)-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

Starting with the hydroxamic acid from example 129, example 130 was prepared in a procedure analogous to example 117. MS found: $(M+H)^+ = 467$.

5

Example 131

[1(R)]- α -[2-(acetylamino)ethyl]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide

10 (131a) Chlorotrimethylsilane (0.20 mL, 10 eq) was added to the more polar lactam from (129g) (90.0 mg, 0.168 mmol) in methanol at rt. After 12 h at reflux, additional chlorotrimethylsilane (10 eq) was added and the mixture kept at reflux to another 24 h. Concentration and purification by
15 silica gel chromatography (methanol-dichloromethane, 5:95 then 10:90) provided the aminoester (70 mg, 89%). MS found: $(M+H)^+ = 466$.

(131b) Following a procedure analogous to (49a), the aminoester from (131a) (64 mg, 0.137 mmol) was converted to
20 the acetamide (70 mg, 100%). MS found: $(M+Na)^+ = 630$.

(131c) Following a procedure analogous to (1f), the acetamide from (131b) (65 mg, 0.128 mmol) was converted to the hydroxamic acid (15 mg, 23%). MS found: $(M-H)^- = 508$.

25

Example 132

[1(R)]-1,1-dimethylethyl [3-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-4-(hydroxyamino)-4-oxobutyl]carbamate mono(trifluoroacetate)

30

(132a) Following a procedure analogous to (129g), the lactam mixture from (129d) (6.36 g, 17.4 mmol) was converted to the BOC protected lactam. Silica gel chromatography (ethyl acetate-hexane, 40:60 then 50:50 then 60:40) provided the less
35 polar isomer (3.70 g) and the more polar isomer (3.19 g). The total yield is 85%. MS found: $(M+Na)^+ = 487$.

(132b) Following a procedure analogous to example 117, the more polar isomer from (132a) (3.13 g, 8.59 mmol) was

deprotected to give the lactam (1.70 g, 69%). MS found:
(M+H)⁺ = 365.

(132c) Following a procedure analogous to (3a), the lactam
from (132b) (1.68 g, 4.61 mmol) was hydrogenolized to give the
phenol (1.23 g, 97%). MS found: (M+H)⁺ = 275.

(132d) Following a procedure analogous to (6b), the phenol
from (132c) (1.20 g, 4.37 mmol) was coupled with 4-
bromomethyl-2,6-dimethylpyridine to give the picolyl ether
(1.63 g, 95%). MS found: (M+H)⁺ = 394.

(132e) Following a procedure analogous to (131a), the lactam
from (132d) (1.58 g, 4.02 mmol) was converted to the methyl
ester bis(hydrochloride) (2.00 g, 100%). MS found: (M+H)⁺ =
426.

(132f) Following a procedure analogous to (49a), the
aminoester from (132e) (100 mg, 0.183 mmol) was reacted with
(BOC)₂O to give the t-butyl carbamate (70 mg, 60%). MS found:
(M+H)⁺ = 526.

(132g) Following a procedure analogous to (1f), the ester from
(132f) (65 mg, 0.124 mmol) was converted to the hydroxamic
acid (23.5 mg, 30%). MS found: (M+H)⁺ = 527.

Example 133

**[1(R)]-α-(2-aminoethyl)-3-[4-[(2,6-dimethyl-4-
pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-
pyrrolidineacetamide bis(trifluoroacetate)**

Starting with the hydroxamic acid from example 132,
example 133 was prepared in a procedure analogous to example
117. MS found: (M+H)⁺ = 427.

Example 134

**N-[3-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-
3-methyl-2-oxo-1-pyrrolidinyl]-4-(hydroxyamino)-4-
oxobutyl]-3-pyridinecarboxamide**

Beginning with the amine from (132e) and nicotinoyl
chloride, example 134 was prepared in an analogous series of
reactions to (49a) and (1f). MS found: (M+H)⁺ = 523.

Example 135

[1(R)]-N-[3-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-4-(hydroxyamino)-4-oxobutyl]-4-morpholinecarboxamide mono(trifluoroacetate)

Beginning with the amine from (132e) and 4-morpholinecarbonyl chloride, example 120 was prepared in an analogous series of reactions to (49a) and (1f). MS found: $(M+H)^+ = 540$.

Example 136

[1(R)]-1,1-dimethylethyl [2-[[3-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-4-(hydroxyamino)-4-oxobutyl]amino]-2-oxoethyl]carbamate mono(trifluoroacetate)

Beginning with the amine from (132e) and N-(t-butoxycarbonyl)glycine, example 136 was prepared in an analogous series of reactions to (50a) and (1e). MS found: $(M+H)^+ = 584$.

Example 137

[1(R)]-α-[2-[(aminoacetyl)amino]ethyl]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

Starting with the hydroxamic acid from example 136, example 137 was prepared in a procedure analogous to example 117. MS found: $(M+H)^+ = 484$.

Example 138

[1(R)]-1,1-dimethylethyl [2-[[2-[[3-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-4-(hydroxyamino)-4-oxobutyl]amino]-2-oxoethyl]amino]-2-oxoethyl]carbamate mono(trifluoroacetate)

Beginning with the amine from (132e) and BOC-Gly-Gly-OH, example 138 was prepared in an analogous series of reactions to (50a) and (1e). MS found: $(M+H)^+ = 641$.

Example 139

[1(R)]- α -[2-[[[(aminoacetyl)amino]acetyl]amino]ethyl]-
3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-
hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide
bis(trifluoroacetate)

Starting with the hydroxamic acid from example 138, example 139 was prepared in a procedure analogous to example 117. MS found: $(M+H)^+ = 541$.

Example 140

[1(R)]-N-hydroxy-3-methyl-2-oxo- α -
[(phenylmethoxy)methyl]-3-[4-(phenylmethoxy)phenyl]-1-
pyrrolidineacetamide

Beginning with the aldehyde from (1c) and (D)-Ser(OBn)-OMe, example 140 was prepared in an analogous series of reactions to (1d) and (1e). MS found: $(M-H)^- = 473$.

Example 141

[1(R)]-3-[4-[(2,6-dichloro-4-
pyridinyl)methoxy]phenyl]-N-hydroxy- α -(hydroxymethyl)-
3-methyl-2-oxo-1-pyrrolidineacetamide

Beginning with the aldehyde from (1c) and (D)-Ser(OBn)-OMe, example 141 was prepared in an analogous series of reactions to (1d), (3a), (6b) and (1e). MS found: $(M-H)^- = 437$.

Example 142

[1(R)]-1,1-dimethylethyl 4-[2-(hydroxyamino)-1-[3-
methyl-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-
pyrrolidinyl]-2-oxoethyl]-1-piperidinecarboxylate
mono(trifluoroacetate)

(142a) To 2-(R)-azido-2-(N-t-BOC-4-piperidinyl)acetic acid (50.0 g, 213 mmol, Ciba-Geigy, EP606046 1994) in methanol (125 mL) and benzene (500 mL) was added a 2 M hexane solution of trimethylsilyl diazomethane (110 mL, 1.03 eq). After 10 min at rt, the mixture was concentrated. Silica gel chromatography (ethyl acetate-hexane, 10:90 then 20:80) gave the methyl ester (36.8 g, 58%). MS found: $(M+H)^+ = 299$.

(142b) A mixture of the azido ester from (142a) (36.8 g, 123 mmol), 10% Pd on carbon (8.0 g) in water (600 mL), THF (600 mL) and acetic acid (200 mL) was stirred under balloon pressure hydrogen at rt overnight. The catalyst was removed by filtration and the filtrate was concentrated to give the amino ester (29.5 g, 88%). MS found: $(M+H)^+ = 273$.

(142c) Following a procedure analogous to step (1d), the aldehyde from (1c) (2.00 g, 6.40 mmol) was reacted with the amino ester from (142b) (2.09 g, 1 eq) to give the crude lactam as mixture of two isomers. The BOC protecting group came off during the cyclization. MS found: $(M+H)^+ = 437$.

(142d) Following a procedure analogous to step (116b), the crude material from (142c) was reacted with (BOC)2O to provide the carbamate (2.13 g, 62%) as a 1:1 mixture. MS found: $(M+Na)^+ = 559$.

(142e) Following a procedure analogous to step (3a), the lactam from (142d) (2.13 g, 3.97 mmol) was hydrogenolized to give the phenol (1.72 g, 97%). MS found: $(M-H)^- = 445$.

(142f) Following a procedure analogous to step (6b), the phenol from (142e) (700 mg, 1.57 mmol) was reacted with 4-chloromethylquinoline hydrochloride to give the ether (744 mg, 81%). MS found: $(M+H)^+ = 588$.

(142g) Following a procedure analogous to step (92d), the ester from (142f) (160 mg, 0.272 mmol) was reacted with hydroxylamine. The product was purified by reverse phase HPLC on a Dynamax C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the fast moving isomer (61.5 mg) and the slow moving isomer (53.0 mg). MS found: $(M+H)^+ = 589$.

Example 143

[1(R)]-N-hydroxy- α -[3-methyl-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-4-piperidineacetamide mono(trifluoroacetate)

Starting with the slow moving isomer from example 142, example 143 was prepared in a procedure analogous to example 117. MS found: $(M+H)^+ = 489$.

Example 144

[1(R)]-N-hydroxy- α -[3-methyl-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-1-(methylsulfonyl)-4-piperidineacetamide mono(trifluoroacetate)]

(144a) Following a procedure analogous to example 117, the lactam from (142f) (553 mg, 0.941 mmol) was reacted with TFA to give the piperidine mono(trifluoroacetate) (1.04, 100%). MS found: $(M+H)^+ = 488$.

(144b) Following a procedure analogous to (49a), the piperidine from (144a) (200 mg, 0.278 mmol) was reacted with MsCl to give the sulfonamide (112 mg, 71%). MS found: $(M+H)^+ = 566$.

(144c) Following a procedure analogous to step (92d), the ester from (144b) (112 mg, 0.198 mmol) was reacted with hydroxylamine. The product was purified by reverse phase HPLC on a Dynamax C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the fast moving isomer (14.0 mg) and the slow moving isomer (13.5 mg). MS found: $(M+H)^+ = 567$.

Example 145

[1(R)]-1-(2-furanylcarbonyl)-N-hydroxy- α -[3-methyl-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-4-piperidineacetamide mono(trifluoroacetate)]

Beginning with the piperidine from (144a) and 2-furic acid, example 145 was prepared in an analogous series of reactions to (50a) and (92d). MS found: $(M+H)^+ = 583$.

Example 146

[1(R)]-1,1-dimethylethyl 4-[1-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-2-(hydroxyamino)-2-oxoethyl]-1-piperidinecarboxylate mono(trifluoroacetate)]

(146a) Following a procedure analogous to step (6b), the phenol from (142e) (1.07 g, 2.40 mmol) was reacted with 4-chloromethyl-2,6-dimethylpyridine hydrochloride to give the picolyl ether (1.15 g, 85%). MS found: $(M+Na)^+ = 588$.

(146b) Following a procedure analogous to step (92d), the ester from (146a) (124 mg, 0.219 mmol) was reacted with hydroxylamine. The product was purified by reverse phase HPLC on a Dynamax C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the fast moving isomer (40.0 mg) and the slow moving isomer (30.0 mg). MS found: $(M+H)^+ = 567$.

Example 147

[1(R)]- α -[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

Starting with the slow moving isomer from example 146, example 147 was prepared in a procedure analogous to example 117. MS found: $(M+H)^+ = 467$.

Example 148

[1(R)]-methyl 4-[1-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-2-(hydroxyamino)-2-oxoethyl]-1-piperidinecarboxylate mono(trifluoroacetate)

(148a) Following a procedure analogous to example 117, the 1:1 mixture of lactams from (146a) (1.01 g, 1.79 mmol) was reacted with TFA to give the piperidine mono(trifluoroacetate) (1.22 g, 100%). MS found: $(M+H)^+ = 466$.

(148b) Following a procedure analogous to (49a), the piperidine from (148a) (75.4 mg, 0.109 mmol) was reacted with methyl chloroformate to give the crude carbamate. MS found: $(M+H)^+ = 524$.

(148c) Following a procedure analogous to step (92d), the crude ester from (148b) was reacted with hydroxylamine. The diastereomeric mixture was purified by reverse phase HPLC on a Dynamax C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the slow moving isomer (14.1 mg). MS found: $(M+H)^+ = 525$.

Example 149

[1(R)]- α -[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-N-hydroxy-1-(methylsulfonyl)-4-piperidineacetamide mono(trifluoroacetate)

Beginning with the piperidine from (148a) and mathanesulfonyl chloride, example 149 was prepared in an analogous series of reactions to (49a) and (92d). MS found: (M+H)⁺ = 545.

Example 150

[1(R)]-1-acetyl- α -[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide mono(trifluoroacetate)

Beginning with the piperidine from (148a) and acetyl chloride, example 150 was prepared in an analogous series of reactions to (49a) and (92d). MS found: (M+H)⁺ = 509.

Example 151

[1(R)]-1-(2,2-dimethyl-1-oxopropyl)- α -[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide mono(trifluoroacetate)

Beginning with the piperidine from (148a) and trimethylacetyl chloride, example 151 was prepared in an analogous series of reactions to (49a) and (92d). MS found: (M+H)⁺ = 551.

Example 152

[1(R)]- α -[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-N-hydroxy-1-methyl-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the piperidine from (148a) and formaldehyde, example 152 was prepared in an analogous series of reactions to (86a) and (92d). MS found: (M+H)⁺ = 481.

Example 153

[1(R)]- α -[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-N-hydroxy-1-(1-methylethyl)-4-piperidineacetamide bis(trifluoroacetate)]

Beginning with the piperidine from (148a), sodium cyanoborohydride and acetone, example 153 was prepared in an analogous series of reactions to (86a) and (92d). MS found: (M+H)⁺ = 510.

Example 300

[1(R)]-3-amino-N-hydroxy- α -(2-methylpropyl)-2-oxo-3-[4-(2-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide mono(trifluoroacetate)

(300a) The p-hydroxy phenyl glycine (74.0 g, 442 mmol) was suspended in methanol (500 mL), cooled in an ice bath and HCl (gas) was bubbled through the reaction mixture for 20 minutes, to give a clear solution. The reaction was stirred at rt for 48 h, concentrated in vacuo to give an oil which was triturated with ethyl ether to give the p-hydroxy phenyl glycine methyl ester (95.8 g, 99%) as a white powder. MS found: (M+H)⁺ = 182.

(300b) The Di-t-butyl dicarbonate (105.0 g, 484 mmol) dissolved in DMF (100 mL) was added slowly to an ice cooled solution of p-hydroxy phenyl glycine methyl ester (95.8 g, 440 mmol), triethyl amine (101 mL) and DMF (800 mL). The reaction was allowed to warm to rt, stirred for 5 h, partitioned between ethyl acetate and 1N HCl. The organic layer was washed with brine, dried over magnesium sulfate and was concentrated in vacuo to give the N-Boc product (123.0 g, 100%) an amber oil. MS (M-H)⁻ = 280.

(300c) The N-Boc p-hydroxy phenyl glycine methyl ester from step (300a) (123.0 g, 440 mmol) was combined with benzyl bromide (90.3 g, 528 mmol), potassium carbonate (182 g, 1.3 mol) and acetone (800 mL) under a nitrogen atmosphere. The reaction was heated to reflux for 5 h, allowed to cool to rt,

diluted with ethyl acetate (800 mL) filtered to remove the solids and concentrated in vacuo to give a semisolid residue. The product was crystallized from ethyl ether to give the N-Boc p-benzyloxy phenyl glycine methyl ester (106.7 g, 65%) as a white powder. MS (M+H)⁺ = 372, (M+NH₄)⁺ = 389.

(300d) The LDA (148.1 mL, 296.2 mmol) was added slowly to a solution of the N-Boc p-benzyloxy phenyl glycine methyl ester from step (300c) (55.0 g, 148.1 mmol) in THF (500 mL) cooled to -78 °C under a nitrogen atmosphere. The reaction was allowed to stir for 1 h and the allyl bromide (17.9 g, 148.1 mmol) was added. The reaction was allowed to warm to 0 °C and stir for 1.5 h. The reaction was partitioned between ethyl acetate and 1 N HCl. The organic layer was washed with brine dried over magnesium sulfate and concentrated in vacuo to give an oil. The product was purified by flash chromatography on silica gel (hexane:ethyl acetate, 85:15, v:v) to give olefin (50.1 g, 82%). MS (M+Na)⁺ = 434.

(300e) Following a procedure analogous to that used in step (1c), the olefin from (300d) (5.0 g, 11.37 mmol) was oxidized to the aldehyde. The product was purified by flash chromatography on silica gel (hexane:ethyl acetate, 70:30, v:v) to give the desired aldehyde (4.6 g, 98%). MS (M+Na)⁺ = 436.

(300f) The aldehyde from (300e) (4.0 g, 9.67 mmol) was combined with leucine methyl ester hydrochloride (2.1 g, 11.6 mmol) and DIEA (1.49 g, 11.6 mmol) in 1,2 1,2-dichloroethane (50 mL) at rt and stirred for 1 h. To this solution the sodium triacetoxymethylborohydride (3.1 g, 14.5 mmol) was added. The reaction was stirred for 2 h, diluted with methylene chloride washed with brine, dried over magnesium sulfate and concentrated in vacuo, to give the amine (5.2 g, 100%) as a clear oil. MS (M+H)⁺ = 543.

(300g) The amine from (300f) (5.2 g, 9.67 mmol) was dissolved in toluene (100 mL) under a nitrogen atmosphere and was heated to 90 °C for 4 h. The reaction was allowed to cool to rt, concentrated in vacuo to give a crude oil which was purified by flash chromatography on silica gel (hexane: ethyl acetate,

85:15, v:v) to give the desired lactam as two separated diastereomers (4.8 g, 97%) as a glass. MS (M+H)⁺ = 511. (300h) The lactam from (300g) (2.6 g, 3.9 mmol) was dissolved in methanol (50 mL), degassed with nitrogen, 10% Pd/C was added and the reaction was charged to 50 PSI hydrogen. The reaction was shaken for 3 h, filtered through celite to remove the catalyst, concentrated in vacuo to give the phenol product (1.6 g, 100%) as a white foam. MS (M+H)⁺ = 421, MS (M+Na)⁺ = 443.

(300i) The phenol product from (300h) (0.15 g, 0.35 mmol) was combined with 2(chloromethyl)quinoline (0.15 g, 0.71 mmol), cesium carbonate (3 eq) and sodium iodide in acetone (15 mL), then heated to reflux. The reaction was heated for 3 h, cooled, diluted with ethyl acetate, filtered to remove the solids and concentrated in vacuo to give a crude oil. The product was purified by flash chromatography on silica gel (methylene chloride:ethyl acetate, 80:20, v:v) to give the desired lactam product (0.15 g, 76%) as a white foam. MS (M+H)⁺ = 562 (M-NH₂)⁺ = 445.

(300j) The N-Boc lactam from (300i) (0.14 g, 0.25 mmol) was dissolved in methylene chloride (2 mL) and TFA (2 mL) under a nitrogen atmosphere. The reaction was stirred for 2 h, concentrated in vacuo to give the expected amino lactam (0.14 g, 100%) as an oil. MS (M+H)⁺ = 462, (M-NH₂)⁺ = 445.

(300k) Following a procedure analogous to that used in step (1f), the methyl ester amino lactam product from (300j) (0.14 g, 0.30 mmol) was converted to the crude hydroxamic acid which was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.085 g, 49%) as a white amorphous solid. MS (M+H)⁺ = 463, (M-NH₂)⁺ = 446.

Example 301

[1(R)]-3-amino-3-[4-[(3,5-

dimethylphenyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(301a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester

in step (300f) and 3,5-dimethyl benzyl bromide in step (300i), the crude hydroxamic acid was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the
5 hydroxamic acid product (0.021 g, 42%) as a white amorphous solid. MS (M+H)⁺ = 398, (M-NH₂)⁺ = 381.

Example 302

10 [1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-3-
[[(ethylamino)carbonyl]amino]-N-hydroxy-alpha-methyl-
2-oxo-1-pyrrolidineacetamide

(302a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester
15 in step (300f) and 3,5-dimethyl benzyl bromide in step (300i), the amino lactam methyl ester from step (j) was prepared and purified by crystallization from ethyl ether (0.28 g, 40%).
MS (M+Na)⁺ = 419, (M-NH₂)⁺ = 380.

(302b) The ethyl isocyanate (0.0035 g, 0.05 mmol) was added to
20 a solution of amino lactam methyl ester (302a) (0.025 g, 0.05 mmol), methylene chloride (1 mL) and N-methyl morpholine (2 eq) at rt under a nitrogen atmosphere. After stirring for 1 h the reaction was concentrated in vacuo to give the ethyl urea (0.023 g, 98%) as a viscous oil. MS (M+H)⁺ = 468.

25 (302c) Following a procedure analogous to that used in step (1f), the ethyl urea lactam methyl ester product from (302b) (0.023 g, 0.049 mmol) was converted to the crude hydroxamic acid which was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to
30 give the title compound (0.015 g, 64%) as a white amorphous solid. MS (M+Na)⁺ = 491.

Example 303

35 [1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-
hydroxy-alpha-methyl-3-[(methylsulfonyl)amino]-2-oxo-
1-pyrrolidineacetamide

(303a) Following the procedures analogous to that used for the preparation of example (302), but using methane sulfonyl chloride in step (302b) the crude hydroxamic acid was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the title compound (0.010 g, 35%) as a white amorphous solid. MS (M+Na)⁺ = 498.

Example 304

10 [1(R)]-N-[3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-
1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-
pyrrolidinyl]-3-pyridineacetamide
mono(trifluoroacetate)

(304a) The amino lactam methyl ester (302a) (0.05 g, 0.098 mmol) was combined with 3-pyridinyl acetic acid (0.026 g, 0.15 mmol), HATU (0.057 g, 0.15 mmol), NMM (3 eq), and DMF (1 mL) at rt under nitrogen atmosphere. The reaction was stirred for 18 h, partitioned between ethyl acetate and 1 N HCl. The organic layer was washed with brine, dried over MgSO₄, and concentrated in vacuo to give the amide product as a crude oil. MS (M+H)⁺ = 515, MS (M+Na)⁺ = 538.

(304b) Following a procedure analogous to that used in step (1f), the pyridinyl acetamide lactam methyl ester from step (304a) (0.05 g, 0.098 mmol) was converted to the crude hydroxamic acid which was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the title compound (0.025 g, 49%) as a white amorphous solid. MS (M+H)⁺ = 517.

Example 305

30 [1(R)]-N-[3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-
1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-
pyrrolidinyl]-4-pyridinecarboxamide
mono(trifluoroacetate)

35 (305a) Following the procedures analogous to that used for the preparation of example (302), but using isonicotinoyl chloride in step (302b) the crude hydroxamic acid was prepared. The product was purified by reverse phase HPLC on a Vydac C-18

semiprep column eluting an acetonitrile:water:TFA gradient, to give the title compound (0.035 g, 71%) as a white amorphous solid. MS (M+H)⁺ = 503.

5

Example 306

[1(R)]-3-amino-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy] phenyl]-N-hydroxy-alpha-methyl-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

(306a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the crude hydroxamic acid was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the title compound (0.045 g, 33%) as a white amorphous solid. MS (M-H)⁻ = 437, 439.

Example 307

[1(R)]-N-[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]-4-pyridinecarboxamide bis(trifluoroacetate)

(307a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with isonicotinoyl chloride similar to example (305a), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.02 g, 20%) as a white amorphous solid. MS (M+H)⁺ = 544. 546.

Example 308

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-[[[(ethylamino)carbonyl]amino]-N-hydroxy-alpha-methyl-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)]

(308a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with ethyl isocyanate similar to example (302b), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.04 g, 25%) as a white amorphous solid. MS (M+Na)⁺ = 532, 534.

Example 309

[1(R)]-1,1-dimethylethyl [2-[[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]amino]-2-oxoethyl]carbamate mono(trifluoroacetate)

(309a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with N-Boc glycine acid similar to example (304a), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.02 g, 25%) as a white amorphous solid. MS (M+Na)⁺ = 618, 620.

Example 310

[1(R)]-3-[(aminoacetyl)amino]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

(310a) The N-Boc glycine compound example (309) was dissolved in methylene chloride (0.5 mL) and TFA (0.5 mL) at rt under a nitrogen atmosphere. The reaction was stirred for 1 h, concentrated in vacuo to give a residue which was triturated with ethyl ether to give the title compound (0.01 g 82%) as a white solid. MS (M+H)⁺ = 496, 498.

Example 311

[1(R)]-N-[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]-3-pyridineacetamide bis(trifluoroacetate)

(311a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with 3-pyridinyl acetic acid similar to example (304a), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.045 g, 23%) as a white amorphous solid. MS (M+H)⁺ = 558, 560.

Example 312

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-3-[[[(phenylmethyl)amino]carbonyl]amino]-1-pyrrolidineacetamide

(312a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with benzyl isocyanate similar to example (302b), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.05 g, 33%) as a white amorphous solid. MS (M+Na)⁺ = 594, 596.

Example 313

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-[[[(2,4-dimethoxyphenyl)amino]carbonyl]amino]-N-hydroxy-alpha-methyl-2-oxo-1-pyrrolidineacetamide

(313a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with 2,4-dimethoxy phenylisocyanate similar to example (302b), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.035 g, 27%) as a white amorphous solid. MS (M+Na)⁺ =640,642.

Example 314

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-3-[[(phenylamino)carbonyl]amino]-1-pyrrolidineacetamide

(314a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with phenylisocyanate similar to example (302b), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.016 g, 13%) as a white amorphous solid. MS (M+Na)⁺ =580,582.

Example 315

[1(R)]-1,1-dimethylethyl [3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]carbamate

(315a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the N-Boc lactam methyl ester from step (i) was reacted with hydroxylamine hydrochloride similar to example (1f), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18

semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.04 g, 42%) as a white amorphous solid. MS (M+Na)⁺ =561, 563.

5

Example 316

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-3-[[[2-(4-morpholinyl)ethyl]amino]carbonyl]amino]-2-oxo-1-pyrrolidineacetamide

- 10 (316a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) (0.10 g, 0.18 mmol) was dissolved in methylene chloride (3 mL) and saturated sodium bicarbonate solution (1 mL), cooled to 0°C, phosgene in toluene solution was added and the reaction was stirred vigorously for 15 minutes. The reaction was diluted with methylene chloride washed with brine, dried over magnesium sulfate and concentrated to give an oil. The oil was taken up in methylene chloride (2 mL) and the amino ethyl morpholine (0.047 g, 0.36 mmol) was added. The reaction was stirred for 0.5 h at rt and was concentrated to give the urea (0.09 g, 84%) as a crude product. MS (M+H)⁺ =594, 596.
- 25 (316b) The urea lactam methyl ester from step (316a) was reacted with hydroxylamine hydrochloride similar to example (1f), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.03 g, 28%) as a white amorphous solid. MS (M+H)⁺ =595, 597.
- 30

Example 317

[1(R)]-1,1-dimethylethyl N-[[[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]amino]carbonyl]glycine

35

(317a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with t-butyl glycine ester similar to steps (316a and 316b), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.04 g, 37%) as a white amorphous solid. MS (M+Na)⁺ =618, 620.

Example 318

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-3-[[(2-thiazolylamino)carbonyl]amino]-1-pyrrolidineacetamide

(318a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted 2-amino thiazole similar to steps (316a and 316b), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.045 g, 44%) as a white amorphous solid. MS (M+H)⁺ =565, 567.

Example 319

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-3-[[(4-pyridinylamino)carbonyl]amino]-1-pyrrolidineacetamide mono(trifluoroacetate)

(319a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with 4-amino pyridine similar to steps (316a and 316b), to prepare the title compound. The

product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.035 g, 32%) as a white amorphous solid. MS (M+Na)⁺ =581, 583.

5

Example 320

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-[[[(3-hydroxyphenyl)amino]carbonyl]amino]-alpha-methyl-2-oxo-1-pyrrolidineacetamide

10

(320a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted 3-hydroxy aniline similar to steps (316a and 316b), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.011 g, 14%) as a white amorphous solid. MS (M+Na)⁺ =596,598.

15

20

Example 321

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-[[[(2,3-dihydro-2-oxo-1H-benzimidazol-5-yl)amino]carbonyl]amino]-N-hydroxy-alpha-methyl-2-oxo-1-pyrrolidineacetamide

25

(321a) Following the procedures analogous to that used for the preparation of example (300), but using alanine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with 5-amino-1,3-dihydro-2H-benzimidazol-2-one similar to steps (316a and 316b), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.02 g, 22%) as a white amorphous solid. MS (M+Na)⁺ =636, 638.

30

35

Example 322

[1(R)]-3-amino-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide
mono(trifluoroacetate)

(322a) Following the procedures analogous to that used for the preparation of example (300), but using methionine methyl ester in step (300f), the sulfide from step (g) was oxidized (2.6 g, 5.10 mmol) by Oxone (12.55 g, 20.5 mmol) in methanol water solution, at rt. The methanol was removed in vacuo and the aqueous layer was extracted with methylene chloride (2X). The combined organic layer was washed with brine, dried over magnesium sulfate and concentrated in vacuo to give the sulfone (2.6 g, 91%) as a white foam. MS (M+Na)⁺ = 583.

(322b) Following the procedures analogous to that used for the preparation of example (300) steps (h through k), but using the sulfide compound from step (322a) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.03 g, 30%) as a white amorphous solid. MS (M+H)⁺ = 532, 533.

Example 323

[1(R)]-3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidine acetamide
mono(trifluoroacetate)

(323a) Following the procedures analogous to that used for the preparation of example (300), but using methionine methyl ester in step (300f), oxidation methods similar to example (322a) and 3,5-dimethyl-4-picolyl chloride hydrochloride in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.035 g, 35%) as a white amorphous solid. MS (M+H)⁺ = 491.

Example 324

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-3-[[2-thiazolylamino)carbonyl]amino]-1-pyrrolidineacetamide

(324a) Following the procedures analogous to that used for the preparation of example (300), but using methionine methyl ester in step (300f), oxidation methods similar to example (322a) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted 2-amino thiazole similar to example (316a), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.054 g, 20%) as a white amorphous solid. MS (M+H)⁺ =657, 659.

Example 325

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-3-[[2-thiazolylamino)carbonyl]amino]-1-pyrrolidineacetamide mono(trifluoroacetate)

(325a) Following the procedures analogous to that used for the preparation of example (300), but using methionine methyl ester in step (300f), oxidation methods similar to example (322a) and 3,5-dimethyl-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted 2-amino thiazole similar to example (316a), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.055 g, 40%) as a white amorphous solid. MS (M+H)⁺ =617.

Example 326

[5(R)]-2-propenyl [5-[3-amino-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]carbamate
mono(trifluoroacetate)

5 (326a) Following the procedures analogous to that used for the preparation of example (300), but using g-N-Alloc lysine methyl ester in step (300f) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a
10 Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.012 g, 18%) as a white amorphous solid. MS (M+H)⁺ =580, 582.

Example 327

15 [5(R)]-2-propenyl [5-[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]carbamate
bis(trifluoroacetate)

(327a) Following the procedures analogous to that used for the
20 preparation of example (300), but using g-N-Alloc lysine methyl ester in step (300f) and 3,5-dimethyl-4-picolyl chloride hydrochloride in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA
25 gradient, to give the hydroxamic acid product (0.025 g, 25%) as a white amorphous solid. MS (M+Na)⁺ =562.

Example 328

30 [1(R)]-3-amino-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide
mono(trifluoroacetate)

(328a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dichloro-4-picolyl
35 chloride hydrochloride in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA

gradient, to give the hydroxamic acid product (0.03 g, 35%) as a white amorphous solid. MS (M+H)⁺ =481,483.

Example 329

5 [1(R)]-3-[4-[(2,6-dichloro-4-
 pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-
 methylpropyl)-2-oxo-3-[[2-
 thiazolylamino)carbonyl]amino]-1-pyrrolidineacetamide
 (329a) Following the procedures analogous to that used for the
10 preparation of example (300), but using 3,5-dichloro-4-picolyl
chloride hydrochloride in step (300i), the amino lactam methyl
ester from step (j) was reacted 2-amino thiazole similar to
step (316a), the title compound was prepared. The product was
purified by reverse phase HPLC on a Vydac C-18 semiprep column
15 eluting an acetonitrile:water:TFA gradient, to give the
hydroxamic acid product (0.01 g, 25%) as a white amorphous
solid. MS (M+Na)⁺ =629,631.

Example 330

20 [1(R)]-3-[4-[(2,6-dimethyl-4-
 pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-
 methylpropyl)-2-oxo-3-[[2-
 thiazolylamino)carbonyl]amino]-1-pyrrolidineacetamide
 mono(trifluoroacetate)
25 (330a) Following the procedures analogous to that used for the
preparation of example (300), but using 3,5-dimethyl-4-
picolyl chloride hydrochloride in step (300i), the amino
lactam methyl ester from step (j) was reacted 2-amino thiazole
similar to step (316a), the title compound was prepared. The
30 product was purified by reverse phase HPLC on a Vydac C-18
semiprep column eluting an acetonitrile:water:TFA gradient to
give the hydroxamic acid product (0.01 g, 20%) as a white
amorphous solid. MS (M+H)⁺ =567.

35

Example 331

[1(R)]-3-[4-[(2,6-dichloro-4-
 pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-
 methylpropyl)-2-oxo-3-[[2-

**pyridinylamino)carbonyl]amino]-1-pyrrolidineacetamide
mono(trifluoroacetate)**

(331a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with 2-amino pyridine similar to step (316a), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.02 g, 20%) as a white amorphous solid. MS (M+Na)⁺ =623,625.

Example 332

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[(trifluoroacetyl)amino]-1-pyrrolidineacetamide mono(trifluoroacetate)

(332a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dimethyl-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with trifluoroacetic anhydride similar to step (302b), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.051 g, 25%) as a white amorphous solid. MS (M+H)⁺ =537.

Example 333

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[(2-pyridinylamino)carbonyl]amino]-1-pyrrolidineacetamide bis(trifluoroacetate)

(333a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dimethyl-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl

ester from step (j) was reacted with 2-amino pyridine similar to step (316a), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the
5 hydroxamic acid product (0.03 g, 25%) as a white amorphous solid. MS (M+H)⁺ =561.

Example 334

[1(R)]-3-[4-[(2,6-dichloro-4-
10 pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-
methylpropyl)-2-oxo-3-
[[[(phenylsulfonyl)amino]carbonyl]amino]-1-
pyrrolidineacetamide

(334a) Following the procedures analogous to that used for the
15 preparation of example (300), but using 3,5-dichloro-4-picoly chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with benzenesulfonyl isocyanate similar to step (302b), the title compound was prepared. The product was purified by reverse phase HPLC on a
20 Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.025 g, 20%) as a white amorphous solid. MS (M+Na)⁺ =686,688.

Example 335

25 [1(R)]-3-[4-[(2,6-dimethyl-4-
pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-
methylpropyl)-2-oxo-3-
[[[(phenylsulfonyl)amino]carbonyl]amino]-1-
pyrrolidineacetamide mono(trifluoroacetate)

30 (335a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dimethyl-4-picoly chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with benzenesulfonyl isocyanate similar to step (302b), the title compound was
35 prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.035 g, 30%) as a white amorphous solid. MS (M+H)⁺ =624.

Example 336

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-[[[(3-methyl-5-isothiazolyl)amino]carbonyl]amino]-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide

(336a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with 5-amino-3-methyl isothiazole similar to step (316a), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.01 g, 20%) as a white amorphous solid. MS (M+H)⁺ = 621, 623.

Example 337

[1(R)]-3-[[[(1H-benzimidazol-2-ylamino)carbonyl]amino]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide

(337a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with 2-amino benzimidazole similar to step (316a), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.005 g, 5%) as a white amorphous solid. MS (M+H)⁺ = 640, 642.

Example 338

[1(R)]-3-[[[(1H-benzimidazol-2-ylamino)carbonyl]amino]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(338a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dimethyl-4-picolyl

chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with 2-amino benzimidazole similar to step (316a), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.015 g, 25%) as a white amorphous solid. MS (M+H)⁺ =600.

Example 339

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[[phenylamino]carbonyl]amino]-1-pyrrolidineacetamide mono(trifluoroacetate)

(339a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dimethyl-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with benzene isocyanate similar to step (302b), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.02 g, 20%) as a white amorphous solid. MS (M+H)⁺ =560.

Example 340

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[[phenylamino]carbonyl]amino]-1-pyrrolidineacetamide

(340a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dichloro-4-picolyl chloride hydrochloride in step (300i), the amino lactam methyl ester from step (j) was reacted with benzene isocyanate similar to step (302b), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.015 g, 20%) as a white amorphous solid. MS (M+Na)⁺ =622,624.

Example 341

[1(R)]-1-[1-[(hydroxyamino)carbonyl]-3-methylbutyl]-N,N,N-trimethyl-2-oxo-3-[4-(phenylmethoxy)phenyl]-1-pyrrolidinemethanaminium trifluoroacetate

- 5 (341a) Following the procedures analogous to that used for the preparation of example (300), but using benzyl bromide in step (300i), the amino lactam methyl ester from step (j) was reacted with methyl iodide and triethylamine in DMSO at rt. The reaction was partitioned between ethyl acetate and
- 10 saturated sodium bicarbonate. The organic layer was washed with brine, dried over magnesium sulfate and concentrated in vacuo to give an oil. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the trimethyl amino
- 15 lactam product (0.025 g, 61%) as an oil. MS (M+H)⁺ =453.
- (341b) Following the procedures analogous to that used for the preparation of step (1f) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to
- 20 give the hydroxamic acid product (0.01 g, 50%) as a white amorphous solid. MS (M+H)⁺ =454.

Example 342

[1(R)]-3-amino-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide mono(trifluoroacetate)

- 25 (342a) Following the procedures analogous to that used for the preparation of example (300), but using 4-chloromethyl quinoline hydrochloride in step (300i), the title compound was
- 30 prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.075 g, 52%) as a white amorphous solid. MS (M+H)⁺ =463, MS (M-NH2)⁺ =
- 35 446.

Example 343

[1(R)]-3-amino-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[4-(2-oxo-2-phenylethoxy)phenyl]-1-pyrrolidineacetamide mono(trifluoroacetate)

(343a) Following the procedures analogous to that used for the preparation of example (300), but using 2-bromoacetophenone in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.075 g, 52%) as a white amorphous solid. MS (M+H)⁺ =455.

Example 344

[1(R)]-3-amino-3-[4-[(3,5-dimethyl-4-isoxazolyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(344a) Following the procedures analogous to that used for the preparation of example (300), but using 4-(chloromethyl)-3,5-dimethyl-isoxazole in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.075 g, 53%) as a white amorphous solid. MS (M+H)⁺ =431, MS (M-NH2)⁺ = 414.

Example 345

[1(R)]-3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

(345a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dimethyl-4-picoly chloride hydrochloride in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.160 g, 55%) as a white amorphous solid. MS (M+H)⁺ =441.

Example 346

[1(R)]-3-amino-3-[4-[2-(2-benzothiazolylamino)-2-oxoethoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

5 (346a) Following the procedures analogous to that used for the preparation of example (300), but using 2-chloro-N(2-benzthiazole)acetamide in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA
10 gradient, to give the hydroxamic acid product (0.08 g, 56%) as a white amorphous solid. MS (M+H)⁺ =512, MS (M-NH2)⁺ =495.

Example 347

[1(R)]-3-amino-N-hydroxy-3-[4-[(2-methoxy-4-quinolinyl)methoxy]phenyl]-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

15 (360a) Following the procedures analogous to that used for the preparation of example (300), but using 2-methoxy-4-bromomethyl quinoline in step (300i) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA
20 gradient, to give the hydroxamic acid product (0.12 g, 34%) as a white amorphous solid. MS (M+H)⁺ =493, MS (M-NH2)⁺ = 476.

Example 348

25 [1(R)]-3-amino-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[4-[(2-phenyl-4-quinolinyl)methoxy]phenyl]-1-pyrrolidineacetamide mono(trifluoroacetate)

(362a) Following the procedures analogous to that used for the preparation of example (300), but using 2-phenyl-4-chloromethyl quinoline hydrochloride in step (300i) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid
30 product (0.12 g, 34%) as a white amorphous solid. MS (M+H)⁺ =539.

Example 349

[1(R)]-3-amino-3-[4-[(2,6-dimethyl-4-quinolinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(363a) Following the procedures analogous to that used for the preparation of example (300), but using 2,6-dimethyl-4-chloromethyl quinoline hydrochloride in step (300i) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.12 g, 34%) as a white amorphous solid. MS (M+H)⁺ =491.

Example 350

[1(R)]-3-amino-3-[4-[(2-chloro-4-quinolinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(350a) Following the procedures analogous to that used for the preparation of example (300), but using 2-chloro-4(chloromethyl)quinoline hydrochloride in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.03 g, 20%) as a white amorphous solid. MS (M+H)⁺ =497,499.

Example 351

[1(R)]-3-amino-3-[4-[2-(2,5-dimethoxyphenyl)-2-(hydroxyimino)ethoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(351a) Following the procedures analogous to that used for the preparation of example (300), but using 2-bromo-2',5'-dimethoxy acetophenone in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA

gradient, to give the hydroxamic acid product (0.0 g, %) as a white amorphous solid. MS (M+H)⁺ =515.

Example 352

5 [1(R)]-3-amino-N-hydroxy-3-[4-[(2-methylimidazo[1,2-
alpyridin-3-yl)methoxy]phenyl]-alpha-(2-methylpropyl)-
2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(352a) Following the procedures analogous to that used for the preparation of example (300), the phenol from step (300h)

10 (0.15 g, 0.35 mmol) was combined with 3-hydroxymethyl-2-methyl-imidazo[1,2-
pyridine (0.086 g, 0.53 mmol), DEAD,

triphenylphosphine and benzene at rt. The reaction was stirred for 2 h, partitioned between ethyl acetate and water, the organic layer was washed with brine dried over magnesium

15 sulfate and concentrated in vacuo to give an oil. The product was purified by flash chromatography on silica gel eluting ethyl acetate to give the alkylated product (0.088 g, 44%) as an oil. MS (M+H)⁺ =565.

(352b) Following the procedures analogous to that used for the preparation of example (300) and step (1f) the compound from step (352a) was reacted to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.065 g, 72%) as a white

25 amorphous solid. MS (M+H)⁺ =466.

Example 353

30 [1(R)]-3-amino-3-[4-[[1,4-dimethyl-2-(methylthio)-1H-
imidazol-5-yl)methoxy]phenyl]-N-hydroxy-alpha-(2-
methylpropyl)-2-oxo-1-pyrrolidineacetamide
mono(trifluoroacetate)

(353a) Following the procedures analogous to that used for the preparation of example (300), the phenol from step (h) was treated with 2-thiomethyl-3N-5-dimethyl-4-hydroxymethyl

35 imidazole similar to step (352a), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA

gradient, to give the hydroxamic acid product (0.09 g, 44%) as a white amorphous solid. MS (M+H)⁺ =476.

Example 354

5 [1(R)]-3-amino-3-[4-[[1,5-dimethyl-2-(methylthio)-1H-imidazol-4-yl]methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(354a) Following the procedures analogous to that used for the preparation of example (300), the phenol from step (h) was treated with 2-thiomethyl-3N-methyl-4-methyl-5-hydroxymethyl imidazole similar to step (352a), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.04 g, 45%) as a white amorphous solid. MS (M+H)⁺ =476.

Example 355

20 [1(R)]-3-amino-3-[4-[(2,4-dimethyl-5-thiazolyl)methoxy]phenyl]-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(355a) Following the procedures analogous to that used for the preparation of example (300), the phenol from step (h) was treated with 2,4-dimethyl-5-hydroxymethyl thiazole similar to step (352a), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.150 g, 75%) as a white amorphous solid. MS (M+H)⁺ =447.

Example 356

[1(R)]-3-amino-N-hydroxy-alpha-(2-methylpropyl)-3-[4-[(2-methyl-4-quinolinyl)methoxy]phenyl]-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

35 (356a) Following the procedures analogous to that used for the preparation of example (300), but using 2-methyl-4-chloromethyl quinoline hydrochloride similar to step (300i), the title compound was prepared. The product was purified by

reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.055 g, 70%) as a white amorphous solid. MS (M+H)⁺ =477.

5

Example 357

[1(R)]-3-amino-3-[4-[(2-chloro-4-quinolinyl)methoxy]phenyl]-N-hydroxy-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

10

(357a) Following the procedures analogous to that used for the preparation of example (300), but using methionine methyl ester in step (300f), oxidation methods similar to example (322a), and 2-chloro-4-chloromethyl quinoline hydrochloride in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.12 g, 34%) as a white amorphous solid. MS (M+H)⁺ =547,549, MS (M-NH₂)⁺ 530,532.

15

20

Example 358

[1(R)]-3-amino-N-hydroxy-3-[4-[(2-methyl-4-quinolinyl)methoxy]phenyl]-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

25

(358a) Following the procedures analogous to that used for the preparation of example (300), but using methionine methyl ester in step (300f), oxidation methods similar to step (322a) and 2-methyl-4-chloromethyl quinoline hydrochloride in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.12 g, 34%) as a white amorphous solid. MS (M+H)⁺ =527.

30

35

Example 359

[1(R)]-3-amino-3-[4-[(3,5-dimethoxyphenyl)methoxy]phenyl]-N-hydroxy-alpha-[2-

(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide
mono(trifluoroacetate)

(359a) Following the procedures analogous to that used for the preparation of example (300), but using methionine methyl ester in step (300f), oxidation methods similar to step
5 (322a), 3,5-dimethoxy benzyl bromide in step (300i) and the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid
10 product (0.12 g, 34%) as a white amorphous solid. MS (M+H)⁺ = 522, MS (M-NH₂)⁺ = 505.

Example 360

[1(R)]-3-amino-N-hydroxy-3-[4-[(2-methoxy-4-
15 **quinolinyl)methoxy]phenyl]-alpha-[2-**
(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide
mono(trifluoroacetate)

(361a) Following the procedures analogous to that used for the preparation of example (300), but using methionine methyl ester in step (300f), 2-methoxy-4-bromomethyl quinoline in
20 step (300i) and oxidation similar to prep (322a) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid
25 product (0.12 g, 34%) as a white amorphous solid. MS (M+H)⁺ = 543, MS (M-NH₂)⁺ = 526.

Example 361

[1(R)]-3-amino-3-[4-[(3,5-
30 **dimethoxyphenyl)methoxy]phenyl]-N-hydroxy-alpha-(2-**
methylpropyl)-2-oxo-1-pyrrolidineacetamide
mono(trifluoroacetate)

(361a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dimethoxy
35 benzyl bromide in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA

gradient, to give the hydroxamic acid product (0.12 g, 34%) as a white amorphous solid. MS (M-NH₂)⁺ = 455.

Example 362

5

[1(R)]-3-amino-3-[4-[(2-methoxy-5-nitro-phenyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

10 (362a) Following the procedures analogous to that used for the preparation of example (300), but using 2-methoxy-5-nitro benzylbromide in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to
15 give the hydroxamic acid product (0.065 g, 25%) as a white amorphous solid. MS (M-NH₂)⁺ = 470.

Example 363

20

[1(R)]-3-amino-3-[4-[(5-quinolinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(363a) Following the procedures analogous to that used for the preparation of example (300), but using 5-chloromethyl
25 quinoline in step (300i), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.055 g, 50%) as a white amorphous solid. MS (M-NH₂)⁺ = 446.

30

Example 364

35

[1(R)]-3-amino-N-hydroxy-3-[4-[(2-methoxy-5-nitro-phenyl)methoxy]phenyl]-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(364a) Following the procedures analogous to that used for the preparation of example (300), but using methionine methyl

ester in step (300f), 2-methoxy-5-nitro-benzylbromide in step (300i) and oxidation similiar to step (322a) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.17 g, 60%) as a white amorphous solid. MS (M+H)⁺ = 543, MS (M-NH₂)⁺ = 520.

Example 365

10 [1(R)]-3-amino-N-hydroxy-3-[4-[(2-nitro-4,5-dimethoxy-phenyl)methoxy]phenyl]-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)]

15 (365a) Following the procedures analogous to that used for the preparation of example (300), but using methionine methyl ester in step (300f), 2-nitro-4,5-dimethoxy benzylbromide in step (300i) and oxidation similiar to step (322a) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.075 g, 42%) as a white amorphous solid. MS (M+H)⁺ = 567, MS (M-NH₂)⁺ = 550.

Example 366

25 [1(R)]-3-amino-N-hydroxy-3-[4-[(2-phenyl-4-quinolinyl)methoxy]phenyl]-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)]

30 (366a) Following the procedures analogous to that used for the preparation of example (300), but using methionine methyl ester in step (300f), 2-phenyl-4-bromomethyl quinoline in step (300i) and oxidation similiar to step (322a) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid

product (0.07 g, 25%) as a white amorphous solid. MS (M+H)⁺ = 589.

Example 367

5 [1(R)]-3-amino-N-hydroxy-3-[4-[(3,5-dimethyl-4-
 isoxazolyl)methoxy]phenyl]-alpha-[2-
 (methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide
 mono(trifluoroacetate)

10 (367a) Following the procedures analogous to that used for
the preparation of example (300), but using methionine methyl
ester in step (300f), 4-(chloromethyl)3,5-dimethyl-isoxazole
in step (300i) and oxidation similar to step (322a) the title
compound was prepared. The product was purified by reverse
15 phase HPLC on a Vydac C-18 semiprep column eluting an
acetonitrile:water:TFA gradient, to give the hydroxamic acid
product (0.12 g, 55%) as a white amorphous solid. MS (M+H)⁺
= 481, MS (M-NH₂)⁺ = 464.

20 **Example 368**

[1(R)]-3-amino-3-[4-[(phenyl)methoxy]phenyl]-N-
 hydroxy-alpha-[(4-hydroxyphenyl)methyl]-2-oxo-1-
 pyrrolidineacetamide mono(trifluoroacetate)

25 (368a) Following the procedures analogous to that used for
the preparation of example (300), but using tyrosine methyl
ester in step (300f), the title compound was prepared. The
product was purified by reverse phase HPLC on a Vydac C-18
semiprep column eluting an acetonitrile:water:TFA gradient, to
30 give the hydroxamic acid product (0.10 g, 50%) as a white
amorphous solid. MS (M+H)⁺ = 462, MS (M-NH₂)⁺ = 445.

Example 369

35 [1(R)]-3-amino-3-[4-[(2-methyl-4-
 quinolinyl)methoxy]phenyl]-N-hydroxy-alpha-[(4-
 methoxyphenyl)methyl]-2-oxo-1-pyrrolidineacetamide
 mono(trifluoroacetate)

(369a) Following the procedures analogous to that used for the preparation of example (300), but using O-methyl tyrosine methyl ester in step (300f) and 2-methyl-4-bromomethyl quinoline in step (300i) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.075 g, 53%) as a white amorphous solid. MS (M+H)⁺ = 541, MS (M-NH₂)⁺ = 524.

Example 370

[1(R)]-3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-[(4-methoxyphenyl)methyl]-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(370a) Following the procedures analogous to that used for the preparation of example (300), but using O-methyl tyrosine methyl ester in step (300f) and 2,6-dimethyl-4-bromomethyl pyridine in step (300i) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.095 g, 77%) as a white amorphous solid. MS (M+H)⁺ = 505, MS (M-NH₂)⁺ = 488.

Example 371

[1(R)]-3-amino-3-[4-[(phenyl)methoxy]phenyl]-N-hydroxy-alpha-[(4-methoxyphenyl)methyl]-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(371a) Following the procedures analogous to that used for the preparation of example (300), but using O-methyl tyrosine methyl ester in step (300f) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.051 g, 25%) as a white amorphous solid. MS (M+H)⁺ = 476, MS (M-NH₂)⁺ = 459.

Example 450

[1(R)]-3-(aminomethyl)-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide
bis(trifluoroacetate)

- 5 (450a) 4-Hydroxybenzyl cyanide (2.5 g, 18.77 mmol), benzyl bromide (3.8 g, 22.5 mmol) and potassium carbonate (45 mmol) were combined in acetone (50 mL) and heated to reflux for 8 h. The reaction was allowed to cool to rt, diluted with ethyl acetate and filtered to remove the solids. The organic layer
- 10 was concentrated in vacuo to give an oil. The crude benzyl ether was purified by chromatography on silica gel eluting hexane: ethyl acetate (90:10, v:v) to give 4-benzyloxybenzyl cyanide (4.0 g, 95%) which solidified. MS (M+NH₄)⁺ =241.
- 15 (450b) The 4-benzyloxybenzyl cyanide from step (450a) (3.2 g, 14.33 mmol), sodium ethoxide (1.07 g, 15.7 mmol), and diethyl carbonate (2.23 g, 18.9 mmol) were combined in toluene (100 mL), heated to reflux for 3 h, cooled to rt, and partitioned between ethyl acetate and 1 N HCl. The organic layer was washed with brine, dried over magnesium sulfate and
- 20 concentrated in vacuo. The crude was purified by chromatography on silica gel eluting hexane: ethyl acetate (80:20, v:v) to give ethyl 2-(4-benzyloxyphenyl)cyanoacetate (4.2 g, 99%) as an oil. MS (M+NH₄)⁺ =313.
- 25 (450c) The ethyl 2-(4-benzyloxyphenyl)cyanoacetate from step (450b) (3.7 g, 12.5 mmol) in DMF (20 mL) was added to a suspension of hexane washed sodium hydride (0.36 g, 15.0 mmol) in DMF (35 mL) cooled in an ice bath under nitrogen. The reaction was allowed to stir for 1 h and the allyl bromide (2.9 g, 24.0 mmol) was added. The reaction was allowed to
- 30 warm to rt and was stirred for 1 h. The reaction was partitioned between ethyl acetate and 1 N HCl. The organic layer was washed with brine, dried over magnesium sulfate and concentrated to give an oil. The crude was purified by chromatography on silica gel eluting hexane: ethyl acetate
- 35 (90:10, v:v) to give ethyl 2-(4-benzyloxyphenyl)-2-allyl cyanoacetate (4.0 g, 95%) as an oil. MS (M+NH₄)⁺ =353.
- (450d) Lithium hydroxide hydrate (1.13 g, 26.8 mmol) in water (20 mL) was added to a solution of ethyl 2-(4-

benzyloxyphenyl)-2-allyl cyanoacetate from step (450c) (4.5 g, 13.42 mmol) in methanol (100 mL) at rt. The reaction was stirred for 5 h, partitioned between ethyl acetate and 1 N HCL. The organic layer was washed with brine, dried over magnesium sulfate and concentrated to give 2-(4-benzyloxyphenyl)-2-allyl cyanoacetic acid (4.1 g, 100%) as an oil. MS (M+NH₄)⁺ =325.

(450e) The 2-(4-benzyloxyphenyl)-2-allyl cyanoacetic acid from step (450d) (2.34 g, 12.88 mmol), TBTU (5.17 g, 16.11 mmol), NMM (4 eq) and DMF (50 mL) were combined and stirred for 15 minutes then the leucine methyl ester (2.34 g, 12.86 mmol) was added. The reaction was allowed to stir at rt for 18 h, partitioned between ethyl acetate and 1 N HCL. The organic layer was washed with brine, dried over magnesium sulfate and concentrated to give an oil. The crude was purified by chromatography on silica gel eluting hexane: ethyl acetate (80:20, v:v) to give the amide (1.9 g, 34%) as an oil. MS (M+NH₄)⁺ =452.

(450f) Ozone was bubbled through a solution of the amide from step (450e) (1.9 g, 4.37 mmol) and methylene chloride (50 mL) cooled to -78 °C. After 20 minutes the reaction turned blue, oxygen and then nitrogen were bubbled through the reaction solution. The triphenylphosphine (1.15 g, 4.37 mmol) was added and the reaction was allowed to warm to rt and stirred for 4 h. The reaction was concentrated in vacuo to give an oil. The crude product was purified by chromatography on silica gel eluting ethyl ether (100%) to give the aldehyde (1.9 g, 100%) as an oil. MS (M+Na)⁺ =459.

(450g) The aldehyde of step (450f) (1.9 g, 4.37 mmol) was dissolved in methylene chloride (15 mL), triethylsilane (5 mL), and TFA (2 mL) at rt under nitrogen. The reaction was stirred for 4 h and was concentrated in vacuo to give an oil. The crude product was purified by chromatography on silica gel eluting hexane: ethyl acetate (70:30, v:v) to give the cyano lactam (1.55 g, 68%) as an oil. MS (M+NH₄)⁺ =438.

(450h) The cyano lactam from step (450g) (1.55 g, 3.68 mmol) was dissolved in methanol (50 mL) degassed with nitrogen, then HCL (conc) (5 drops) and 10% Pd/C were added, the

reaction was charged to 50 PSI hydrogen and shaken for 18 h. The catalyst was removed over celite, the organic layer concentrated in vacuo to give the aminomethyl lactam (1.2 g, 97%) as a foam. MS (M+Na)⁺ =335.

5 (450i) The di-t-butyl dicarbonate (0.85 g, 3.88 mmol) was added to a solution of aminomethyl lactam from step (450h) (1.2 g, 3.24 mmol) and TEA (4 eq) in DMF (20 mL) at rt. The reaction was stirred for 4 h, partitioned between ethyl acetate and 1 N HCl. The organic layer was washed with brine, 10 dried over magnesium sulfate and concentrated to give an oil. The crude was purified by chromatography on silica gel eluting hexane: ethyl acetate (50:50, v:v) to give the N-Boc aminomethyl lactam (0.9 g, 64%) as a foam. MS (M+Na)⁺ =457.

15 (450j) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dimethyl-4-picolyl chloride hydrochloride in step (300i), the removal of the N-Boc protecting group similar to step (300j) the compound from step (450i) was converted to the aminomethyl lactam methyl ester (0.64 g, 100%) isolated as an oil. MS (M+H)⁺ =454

20 (450k) Following the procedures analogous to that used for the preparation of example (1f), the aminomethyl lactam methyl ester from step (450j) (0.10 g, 0.146 mmol) was converted to title compound. The product was purified by reverse phase 25 HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.03 g, 30%) as a white amorphous solid. MS (M+H)⁺ =455.

30 Example 451

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[[[(2-thiazolylamino)carbonyl]amino]methyl]-1-pyrrolidineacetamide mono(trifluoroacetate)]

35 (451a) Following the procedures analogous to that used for the preparation of example (450), the aminomethyl lactam methyl ester from step (450j) was reacted with 2-isocyno

thiazole similar to step (302b), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.075 g, 60%) as a white amorphous solid. MS (M+H)⁺ =581.

Example 452

[1(R)]-3-(aminomethyl)-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(452a) Following the procedures analogous to that used for the preparation of example (450), but using alanine methyl ester in step (450e) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (450j), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.035 g, 35%) as a white amorphous solid. MS (M+H)⁺ =453,455.

Example 453

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-3-[[[(2-thiazolylamino)carbonyl]aminomethyl]-1-pyrrolidineacetamide]

(453a) Following the procedures analogous to that used for the preparation of example (450), but using alanine methyl ester in step (450e) and 3,5-dichloro-4-picolyl chloride hydrochloride in step (450j), the aminomethyl lactam methyl ester similar step (450j) was reacted with 2-isocyano thiazole similar to step (302b), to prepare the title compound. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.03 g, 47%) as a white amorphous solid. MS (M+H)⁺ =579,581.

Example 454

[1(R)]-4-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-
hydroxy- α ,4-dimethyl-5-oxo-1-
imidazolidineacetamide mono(trifluoroacetate)

- (454a) Following the procedures analogous to that used for the preparation of example (300), but using 3,5-dimethyl benzyl bromide in step (300c) and methyl iodide in step (300d) the 4-(3,5-dimethylbenzyloxy) phenyl glycine methyl ester was prepared (1.65 g, 80%) as an oil. MS (M+H, -t-but)⁺ =357.
- (454b) Following the procedures analogous to that used for step (450d), the methyl ester from step (454a) was converted to the 4-(3,5-dimethylbenzyloxy) phenyl glycine acid (1.5 g, 97%) as an oil. MS (M+Na)⁺ =422.
- (454c) Following the procedures analogous to that used for step (450e), but using alanine methyl ester the 4-(3,5-dimethylbenzyloxy) phenyl glycine acid from step (454b) (1.5 g, 97%) was converted to the diamino acid. The crude was purified by chromatography on silica gel eluting hexane: ethyl acetate (75:25, v:v) to give the alanine-phenyl glycine compound (1.4 g, 75%) as a foam. MS (M+H)⁺ =485.
- (454d) Following the procedures similar to that used for step (300j), the N-Boc group of the alanine-phenyl glycine compound from step (454c) was removed to give the amino compound (1.2 gm, 97%) as an oil. MS (M+H)⁺ =385, MS (M-NH₂)⁺ =368.
- (454e) Paraformaldehyde (0.006 g, 0.2 mmol) was added to a solution of the amino compound from step (454d) in toluene (5 mL) and NMM (2 eq), the reaction was heated to 80°C for 4.5 h. The reaction was concentrated in vacuo to give the cyclic compound (0.1 g, 100%) as an oil. MS (M+H)⁺ =397.
- (454f) Following the procedures similar to that used for step (1f), but using the cyclic compound from step (454e) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.015 g, 20%) as a white amorphous solid. MS (M+H)⁺ =398.

Example 455

[1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-
hydroxy-3-(hydroxymethyl)-alpha-methyl-2-oxo-1-
pyrrolidineacetamide

(455a) Methyl 4-hydroxyphenylacetate (8.0 g, 48.0 mmol), 3,5-dimethyl benzyl bromide (12.0 g, 60.0 mmol) and potassium carbonate (8.0 g, 58.0 mmol) were combined in acetone (120 mL) and heated to reflux for 8 h. The reaction was allowed to cool, diluted with ethyl acetate and filtered to remove the solids. The organic solvent was removed in vacuo to give an oil. The crude was purified by chromatography on silica gel eluting hexane: ethyl acetate (95:5, v:v) to give the methyl 4-(2,5-dimethylbenzyloxy)phenyl acetate compound (13.58 g, 99%) as an oil. MS (M+NH₄)⁺ = 302.

(455b) LDA (2.0 M in hexane, 3.5 mL, 7.0 mmol) was added to a solution of methyl 4-(2,5-dimethylbenzyloxy phenylacetate compound from step (455a), (2.0 g, 7.0 mmol) in THF (75 mL) cooled to -78 oC under a nitrogen atmosphere. The reaction was stirred for 40 minutes and the allyl bromide (0.73 mL, 8.4 mmol) was added. The reaction was stirred at -78 oC for 5 h, allowed to warm to rt overnight and was partitioned between ethyl acetate and 1N HCl. The organic layer was washed with brine, dried over magnesium sulfate and concentrated in vacuo. The crude was purified by chromatography on silica gel eluting hexane: ethyl acetate (93:2, v:v) to give the methyl 2-allyl-[4-(2,5-dimethylbenzyloxy)phenyl]acetate compound (1.2 g, 53%) as an oil. MS (M+NH₄)⁺ = 342.

(455c) Sodium methoxide (25% in methanol, 0.08 mL, 0.35 mmol) was added dropwise to a solution of the 2-allyl phenylacetate from step (455b) (1.2 g, 3.7 mmol) and parformaldehyde (0.135 g, 4.5 mmol) in DMSO (20 mL) at rt. The reaction was stirred for 1.2 h, diluted with water, acidified with 1N HCl, and extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and concentrated in vacuo, to give the 2-hydroxymethylene-2-allyl phenylacetate (0.91 g, 68%) as an oil. MS (M+NH₄-OCH₃)⁺ = 342.

(455d) Following the procedures analogous to that used for step (450d), the methyl ester from step (455c) was converted

to the 2-hydroxymethylene-2-allyl phenylacetic acid (0.45 g, 53%) as an oil. MS (M+Na)⁺ =.

(455e) Following the procedures analogous to that used for step (450e), but using alanine methyl ester, the 2-

hydroxymethylene-2-allyl phenylacetic acid from step (455d) (0.4 g, 1.2 mmol) was converted to the diamino acid. The

crude was purified by chromatography on silica gel eluting hexane: ethyl acetate (75:25, v:v) to give the

hydroxymethylene phenylacetamide compound (0.36 g, 71%) as an oil. MS (M-H)⁻ =339.

(455f) The hydroxymethylene compound from step (455e) (0.35 g, 0.82 mmol) was combined with TEA (1.3 eq), DMAP (0.025 g, 0.2 mmol), and t-butyldimethylchlorosilane (0.136 g, 0.90 mmol) in DMF (10 mL) at rt. The reaction was stirred for 48

h, diluted with ethyl acetate, washed with saturated ammonium chloride, dried over magnesium sulfate and concentrated to give an oil. The crude was purified by chromatography on silica gel eluting hexane: ethyl acetate (75:25, v:v) to give the O-t-butyldimethylsilyl hydroxymethylene compound (0.16 g, 36%) as an oil. MS (M+Na)⁺ =539.

(455g) Following the procedures analogous to that used for step (450f), but using allyl phenylacetamide from step (455f) (0.4 g, 0.74 mmol) the aldehyde was prepared. The crude was purified by chromatography on silica gel eluting hexane: ethyl ether (95:5, v:v) to give the aldehyde phenylacetamide compound (0.35 g, 87%) as an oil. MS (M+Na)⁺ =564.

(455h) Following the procedures analogous to that used for step (450g), but using aldehyde phenylacetamide compound from step (455g) (0.35 g, 0.65 mmol) the hydroxymethylene lactam was prepared. The crude was purified by chromatography on silica gel eluting methylene chloride: methanol (99:1, v:v) to give the hydroxymethylene lactam compound (0.185 g, 69%) as an oil. MS (M+H)⁺ =412.

(455i) Following the procedures analogous to that used for step (450d), but using hydroxymethylene lactam methyl ester compound from step (455h) (0.35 g, 0.65 mmol) the hydroxymethylene lactam acid (0.18 g, 100%) was prepared as an oil. MS (M+Na)⁺ =420.

(455j) Following the procedures analogous to that used for the preparation of step (450e), but using hydroxylamine hydrochloride and the hydroxymethylene lactam acid compound from step (455i) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.055 g, 30%) as a white amorphous solid. MS (M+Na)⁺ =435.

Example 456

[1(R)]-[3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]methyl ethylcarbamate

(456a) Following the procedures analogous to that used for the preparation of step (302b), but using ethyl isocyanate the hydroxymethylene lactam from step (455h), the lactam carbamate methyl ester compound (0.058 g, 100%) was prepared as an oil. MS (M+Na)⁺ =505.

(456b) Following the procedures similar to that used for step (1f), but using the carbamate lactam compound from step (456a), the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.019 g, 36%) as a white amorphous solid. MS (M+Na)⁺ =506.

Example 457

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-(hydroxymethyl)-alpha-methyl-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(457a) Following the procedures analogous to that used for the preparation of step (300h), but using the hydroxymethylene lactam from step (455h) and 3,5-dichloro-4-picolyl bromide hydrochloride similar to step (300i) and procedures similar to steps (455i and 455j) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to

give the hydroxamic acid product (0.03 g, 18%) as a white amorphous solid. MS (M+Na)⁺ = 476, 478.

Example 458

5 **[1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-azetidineacetamide**

~~(458a)~~ Following the procedures analogous to that used for the preparation of example (455), but using methyl iodide in step (455b) the hydroxymethylene acetamide methyl ester (0.10 g, 0.25 mmol) from step (e) was reacted with methanesulfonyl chloride (0.025 mL, 0.32 mmol) in pyridine at rt, to give the methanesulfonylmethyl acetamide (0.1 g, 84%) as an oil. MS (M+Na)⁺ = 500.

10 (458b) The methanesulfonylmethyl acetamide (0.1 g, 0.21 mmol) from step (458a) was combined with potassium carbonate (0.125 g, 0.9 mmol) in acetone (3 mL), heated to reflux for 6 h, allowed to cool to rt, diluted with ethyl acetate, filtered to remove the solids and concentrated to give an oil. The crude was purified by chromatography on silica gel eluting hexane: ethyl acetate (80:20, v:v) to give the beta-lactam compound (0.05 g, 63%) as an oil. MS (M+H)⁺ = 382.

20 (458c) Following the procedures similar to that used for steps (455i and 455j), but using the beta-lactam compound from step (458b) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.03 g, 80%) as a white amorphous solid. MS (M+H)⁺ = 381.

Example 459

30 **[1(R)]-3-[5-[(3,5-dimethylphenoxy)methyl]-2-thiazolyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide**

35 (459a) Following the procedures similar to that used for step (300a), but using thiopheneacetic acid (7.5 g, 52.7 mmol), the methyl ester was prepared. The crude ester was purified by chromatography on silica gel eluting hexane: ethyl acetate

(90:10, v:v) to give the methyl thiopheneacetate (7.5 g, 92%) as a foam. MS (M+H)⁺ =157.

(459b) Following the procedures similar to that used for step (455b), but using the methyl thiopheneacetate from step

5 (459a), the methyl 2-allyl thiopheneacetate was prepared. The crude ester was purified by chromatography on silica gel eluting hexane: ethyl acetate (95:5, v:v) to give the methyl allyl thiopheneacetate (5.9 g, 73%) as a foam. MS (M+H)⁺ =197.

(459c) Following the procedures similar to that used for step

10 (455b), but using methyl iodide and the methyl allyl thiopheneacetate from step (459b), the methyl 2-allyl-2-methyl thiopheneacetate was prepared. The crude ester was purified by chromatography on silica gel eluting hexane: ethyl acetate (95:5, v:v) to give the methyl 2-ally-2-methyl

15 thiopheneacetate (5.6 g, 89%) as an oil. MS (M+NH₄)⁺ =228.

(459d) Following the procedures similar to that used for step (450d), but using methyl 2-ally-2-methyl thiopheneacetate from step (459c), the 2-allyl-2-methyl thiopheneacetic acid was prepared. The crude ester was purified by chromatography on

20 silica gel eluting toluene: ethyl acetate:acetic acid (60:40:2, v:v:v) to give the thiopheneacetic acid (2.5 g, 99%) as an oil. MS (M+NH₄)⁺ =214.

(459e) Following the procedures similar to that used for step (450e), but using 2-ally-2-methyl thiopheneacetic acid from

25 step (459d) and alanine methyl ester, the thiopheneacetamide compound was prepared. The crude ester was purified by chromatography on silica gel eluting hexane: ethyl acetate (80:20, v:v) to give the thiopheneacetamide (1.5 g, 83%) as an oil. MS (M+NH₄)⁺ =299.

30 (459f) Osmium tetroxide (catalytic) was added to a solution of thiopheneacetamide compound from step (459e) (1.5 g, 5.3 mmol), N-methyl morpholine N-oxide (1.25 g, 10.6 mmol), THF (25 mL) and water (2 mL) at rt under a nitrogen atmosphere. The reaction was stirred overnight, poured into 10% NaHSO₃ and
35 1N HCl (50 mL) and was extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and concentrated to give an oil. The crude oil was dissolved in methylene chloride (25 mL) and water (5 mL). The

NaIO₄ (2.28 g, 10.6 mmol) was added and the reaction was stirred vigorously for 4 h. This was diluted with ethyl acetate, washed with brine, dried over magnesium sulfate and concentrated to give the aldehyde (1.5 g, 99%) as an oil. MS

(M+H-H₂O)⁺ =266.

(459g) Following the procedures similar to that used for step (450g), but using aldehyde thiopheneacetacetamide from step (459f) the lactam compound was prepared. The crude ester was purified by chromatography on silica gel eluting hexane:

ethyl acetate (70:30, v:v) to give the lactam thiophene (1.1 g, 77%) as an oil. MS (M+H)⁺ =268.

(459h) Phosphorous oxychloride (0.95 g, 6.17 mmol) was added slowly to a solution of lactam thiophene from step (451g), (1.1 g, 4.11 mmol) in DMF (0.45 g, 6.17 mmol) and heated to 85° C for 4 h. The reaction was allowed to cool, partitioned between ethyl acetate and ice water. The organic layer was washed with brine, dried over magnesium sulfate and concentrated in vacuo to give the thiophene aldehyde (0.75 g, 62%) as an oil.

(459i) Sodium borohydride (0.059 g, 1.69 mmol) was added to a solution of thiophene aldehyde from step (459h) (0.5 g, 1.69 mmol) dissolved in THF (5 mL) and methanol (1 mL), at rt. The reaction was stirred for 20 minutes, partitioned between ethyl acetate and 1N HCl. The organic layer was washed with brine, dried over magnesium sulfate and concentrated in vacuo to give the 5-hydroxymethylene-thiophene (0.5 g, 100%) as an oil.

(459j) The 5-hydroxymethylene-thiophene from step (459i) (5.0 g, 1.69 mmol) was combined with carbon tetrabromide (0.67 g, 2.03 mmol), triphenylphosphine (0.53, 2.03 mmol) in methylene chloride (5 mL) at rt. The reaction was stirred for 4 h and became a dark solution. This was partitioned between methylene chloride and 1N HCl. The organic layer was washed with brine, dried over magnesium sulfate and concentrated in vacuo to give a dark oil. The product was purified by

chromatography on silica gel eluting hexane: ethyl acetate (50:50, v:v), to give the 5-bromomethylene thiophene (0.15 g 25 %) as an oil. MS (M+H-Br+OCH₃)⁺ =312.

(459k) Following the procedures similar to that used for step (300i), but using 5-bromomethylene thiophene from step (459j) and 3,5-dimethyl phenol, the lactam thiophene compound was prepared. The crude ester was purified by chromatography on silica gel eluting methylene chloride: ethyl acetate (95:5, v:v) to give the lactam thiophene (0.08 g, 47%) as an oil. MS (M+NH₄)⁺ =419.

(459l) Following the procedures similar to that used for steps (1f), but using the lactam thiophene compound from step (459k) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.015 g, 20%) as a white amorphous solid. MS (M+Na)⁺ =425.

Example 460

[1(R)]-4-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2,5-dioxo-4-(2-propenyl)-1-imidazolidineacetamide

(460a) Following the procedures similar to that used for step (300j), but using N-Boc phenyl glycine from step (300c) (0.5 g, 1.13 mmol), the deprotected phenyl glycine compound (0.51 g, 99%) was prepared as an oil.

(460b) A solution of alanine methyl ester (0.046 g, 0.33 mmol) in methylene chloride (1 mL) and DIEA (0.130 mL) was added slowly to a solution of triphosgene (0.098 g, 0.33 mmol) in methylene chloride (2 mL) at rt. The reaction was stirred for 0.5 h and a solution of deprotected phenyl glycine from step (460a) in methylene chloride (1 mL) and DIEA (0.13 mL) was added. The reaction was stirred for 2 h, partitioned between ethyl acetate and 1N HCl. The organic layer was washed with brine, dried over magnesium sulfate and concentrated in vacuo to give an oil. The product was purified by chromatography on silica gel eluting methylene chloride: ethyl acetate (90:10, v:v), to give the mixed urea (0.035 g 23 %) as an oil. MS (M+NH₄-OCH₃)⁺ =454.

(460c) A suspension of the mixed urea from step (460b) (0.035 g, 0.075 mmol) and potassium carbonate (3 eq) in acetone (5

mL) was heated to reflux for 2 h. The reaction was allowed to cool, diluted with ethyl acetate and filtered to remove the solids, washed with brine and concentrated to give the hydantoin compound (0.025 g, 76%) as an oil. MS (M+NH₄)⁺ =454.

(460d) Following the procedures similar to that used for steps (1f), but using the hydantoin compound from step (460c) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.015 g, 60%) as a white amorphous solid. MS (M+Na)⁺ =460.

Example 461

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[[4-(phenylmethoxy)phenyl]methyl]-1-pyrrolidineacetamide

(461a) Triphenylphosphine (3.67 g, 14.0 mmol) and carbon tetrabromide (4.46 g, 14.0 mmol) were added to a solution of 4-benzyloxybenzyl alcohol (2.0 g, 9.3 mmol) in dichloromethane (25 mL) at 0 °C. The mixture was warmed to rt for 2.5 h and then concentrated. The residue was triturated with ether, and the solids filtered off. Filtrate was concentrated. Residue purified by silica gel chromatography (ethyl acetate:hexanes, 5:95, v:v). Residue from chromatography was purified further with treatment with ether and filtration of solids. Filtrate was concentrated in vacuo to yield the desired bromide (2.34 g, 90%) as a white solid. MS found: (M-Br)⁺ = 197.

(461b) A 2.0 M THF solution of lithium diisopropylamide (2.6 mL, 1.15 eq) was added over 10 minutes to a solution of ethyl 2-methyl-4-pentenoate (0.75 mL, 4.6 mmol) in THF (18 mL) at -78 °C. The mixture was warmed to -55 °C for 40 minutes then cooled to -78 °C. A solution of bromide compound from step (461a) (1.92 g, 6.9 mmol) in THF was added over 5 minutes to the cooled mixture. After 1 h at -78 °C the mixture was warmed to rt and 1 M HCl (30 mL) was added. The mixture was extracted with ethyl acetate (2 X 30 mL). The combined organic extracts were washed successively with 1N HCl (20 mL), saturated aqueous sodium bicarbonate (20 mL), water (20 mL),

brine (20 mL), dried (MgSO₄) and concentrated. The residue was purified by silica gel chromatography (hexane, then ethyl acetate:hexanes 2:98, v:v) to give the desired product (950 mg, 60%) as a clear oil. MS found: (M+NH₄)⁺ = 356.

- 5 (461c) Ozone was bubbled through a solution of compound (461b) (0.90 g, 2.6 mmol) in dichloromethane (30 mL) at -78 °C until a blue color persisted in the solution. The mixture was purged with oxygen and treated with triphenylphosphine (0.84 g, 3.2 mmol). The reaction mixture was allowed to warm to rt and stirred for 1 h, then was concentrated in vacuo. The residue was purified by silica gel chromatography (hexane, then ethyl acetate:hexanes 6:94, v:v) to give the desired aldehyde (0.70 g, 75%) as a clear oil. MS found: (M+H)⁺ = 341.
- 15 (461d) Following the procedures similar to that used for steps (1d, 1e and 1f), but using the aldehyde compound from step (461c) (650 mg, 1.9 mmol) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.065 g, 20%) as a white amorphous solid. MS (M+Na)⁺ = 405.

Example 462

- 25 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-(methylamino)-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

- (462a) Following the procedures analogous to that used for the preparation of example (300), the N-Boc phenyl glycine compound from step (300d) (3.59 g, 8.72 mmol) was treated with sodium hydride (0.42 g, 17.45 mmol) in DMF (25 mL) at 0 °C for 1 h. The methyl iodide (2.47 g, 17.45 mmol) was added, the reaction was allowed to stir for 2 h at rt, partitioned between ethyl acetate and 1 N HCl. The organic layer was washed with brine, dried over magnesium sulfate and concentrated to give the N-methyl-N-Boc phenyl glycine (3.6 g, 97%) as an oil. MS (M+Na)⁺ = 448.

(462b) Following the procedures analogous to that used for the preparation of example (300), but using the N-methyl-N-Boc phenyl glycine compound from step (462a) and using 2,6-dimethyl picolyl chloride hydrochloride in step (300i) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.12 g, 34%) as a white amorphous solid. MS (M+H)⁺ =455.

Example 463

[1(R)]-N-hydroxy-3-(methylamino)-alpha-(2-methylpropyl)-3-[4-[(2-methyl-4-quinolinyl)methoxy]phenyl]-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

(463a) Following the procedures analogous to that used for the preparation of example (462), but using 2-methyl-4-chloromethyl quinoline hydrochloride in step (300i) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.12 g, 34%) as a white amorphous solid. MS (M+H)⁺ =491.

Example 464

[1(R)]-alpha,3-dimethyl-N-hydroxy-2-oxo-3-[4-(phenylmethoxy)phenyl]-1-piperidineacetamide

(464a) Following the procedures analogous to that used for the preparation of example (1), the ester from step (1b) was treated with lithium hydroxide similar to step (450d) to give the carboxylic acid, which was coupled to alanine methyl ester similar to step (450e), to give the alanyl-phenyl glycine diamino acid as an oil. MS (M+H)⁺ =382.

(464b) 9-BBN (5.0 eq) was added to a solution of the olefin from Step (464a) (0.45 g, 1.18 mmol) in THF (10 mL) cooled to 0°C under nitrogen. The reaction was allowed to warm to rt and stir overnight at rt. The reaction was cooled to 0°C and water (2 mL) was added. The reaction was stirred for 20

minutes, then sodium acetate (1 g, in 2 mL water) and H₂O₂ (30%) (2.5 mL) were added simultaneously. This was stirred for 40 minutes, concentrated in vacuo, diluted with ethyl acetate and washed with water, brine, dried over magnesium sulfate and concentrated in vacuo to give the alcohol. The crude product was purified by chromatography on silica gel eluting methylene chloride: ethyl acetate (1:1, v:v) to give the alcohol (0.41 g, 87%) as an oil. MS (M+H)⁺ = 400.

(464c) Following the procedure similar to that used for the preparation of step (459j), but using the alcohol from step (464b), the bromide was prepared. The crude product was purified by chromatography on silica gel eluting hexane: ethyl acetate (2:1, v:v) to give the bromide (0.145 g, 71%) as an oil. MS (M+H)⁺ = 462, 464.

(464d) The bromide from step (464c) (0.145 g, 0.313 mmol) was treated with sodium hydride (0.019 g, 0.47 mmol) in THF (10 mL) cooled to 0°C under nitrogen. The reaction was stirred for 1.5 h, then partitioned between ethyl acetate and 1N HCl. The organic layer was washed with water, brine, dried over magnesium sulfate and concentrated in vacuo to give the lactam (0.105 g, 84%) as an oil. MS (M+H)⁺ = 382.

(464e) Following the procedures analogous to that used for step (1f), but using the lactam from step (464d) the title compound was prepared. The product was purified by reverse phase HPLC on a Vydac C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid product (0.062 g, 60%) as a white amorphous solid. MS (M+Na)⁺ = 405.

Example 501

[1(R)]-α-[3-amino-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide tris(trifluoroacetate)]

(501a) Following a procedure analogous to (300f), the aldehyde from (300e) (2.80 g, 6.77 mmol) and amino ester from (142b) (2.42 g, 1.1 eq) were coupled to give the secondary amine as a crude material. MS found: (M+H)⁺ = 670.

(501b) Following a procedure analogous to (300g), the crude amine from (501a) was converted to the lactam. Silica gel chromatography (ethyl acetate-hexane, 20:80 then 30:70) provided the less polar isomer (1.40 g) and the more polar isomer (1.30 g). The total yield is 63% for two steps. MS found: $(M+Na)^+ = 660$.

(501c) Following a procedure analogous to step (3a), the less polar lactam from (501b) (1.30 g, 2.04 mmol) was hydrogenolized to give the phenol (1.10 g, 98%). MS found: $(M+H)^+ = 548$.

(501d) Following a procedure analogous to step (6b), the phenol from (501c) (100 mg, 0.183 mmol) was reacted with 4-chloromethylquinoline hydrochloride to give the ether (75.5 mg, 60%). MS found: $(M+H)^+ = 689$.

(501e) Following a procedure analogous to step (92d), the ester from (501d) (69.0 mg, 0.100 mmol) was reacted with hydroxylamine to give the hydroxamic acid (36.0 mg, 52%). MS found: $(M+H)^+ = 690$.

(501f) Following a procedure analogous to example 117, the hydroxamic acid from (501e) (30.0 mg, 0.0362 mmol) was reacted with trifluoroacetic acid to give the hydroxamic acid tris(trifluoroacetate) (40.0 mg, 100%). MS found: $(M+H)^+ = 490$.

Example 502

[1(R)]- α -[3-amino-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (501c) and 4-bromomethyl-2,6-dichloropyridine, example 502 was prepared in an analogous series of reactions to (6b), (92d) and example 117. MS found: $(M+H)^+ = 508$.

Example 503

[1(R)]-1,1-dimethylethyl 4-[1-[3-[(1,1-dimethylethoxy)carbonyl]amino]-3-[4-[(1,1-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-2-

(hydroxyamino)-2-oxoethyl-1-piperidinecarboxylate
mono(trifluoroacetate)

(503a) Following a procedure analogous to step (6b), the phenol from (501c) (1.67 g, 3.05 mmol) was reacted with 4-chloromethyl-2,6-dimethylpyridine hydrochloride to give the picolyl ether (1.576 g, 77%). MS found: $(M+H)^+ = 667$.

(503b) Following a procedure analogous to step (92d), the ester from (501d) (76.0 mg, 0.114 mmol) was reacted with hydroxylamine to give the hydroxamic acid (32.6 mg, 37%). MS found: $(M+H)^+ = 668$.

Example 504

[1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide tris(trifluoroacetate)

Starting with the hydroxamic acid from example 503, example 504 was prepared in a procedure analogous to example 117. MS found: $(M+H)^+ = 468$.

Example 505

[1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-hydroxy-1-(methylsulfonyl)-4-piperidineacetamide bis(trifluoroacetate)

(505a) Following a procedure analogous to example 117, the lactam from (503a) (624 mg, 0.936 mmol) was reacted with TFA to give the piperidine tris(trifluoroacetate) (750 mg, 99%). MS found: $(M+H)^+ = 467$.

(505b) Following a procedure analogous to (49a), the piperidine from (148a) (125 mg, 0.155 mmol) was reacted with methylsulfonyl chloride to give the monosulfonamide (67.0 mg, 80%). MS found: $(M+Na)^+ = 567$.

(505c) Following a procedure analogous to step (92d), the crude ester from (505b) was reacted with hydroxylamine. The mixture was purified by reverse phase HPLC on a Dynamax C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid bis(trifluoroacetate) (45.0 mg, 52%). MS found: $(M+H)^+ = 546$.

Example 506

[1(R)]-1-acetyl- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the piperidine from (505a) and acetyl chloride, example 506 was prepared in an analogous series of reactions to (49a) and (92d). MS found: $(M+H)^+ = 510$.

Example 507

[1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-1-(2,2-dimethyl-1-oxopropyl)-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the piperidine from (505a) and trimethylacetyl chloride, example 507 was prepared in an analogous series of reactions to (49a) and (92d). MS found: $(M+H)^+ = 552$.

Example 508

[1(R)]-1,1-dimethylethyl 4-[1-[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-2-(hydroxyamino)-2-oxoethyl]-1-piperidinecarboxylate bis(trifluoroacetate)

Beginning with the piperidine from (505a) and di-t-butyl dicarbonate, example 508 was prepared in an analogous series of reactions to (49a) and (92d). MS found: $(M+H)^+ = 568$.

Example 509

[1(R)]-methyl 4-[1-[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-2-(hydroxyamino)-2-oxoethyl]-1-piperidinecarboxylate bis(trifluoroacetate)

Beginning with the piperidine from (505a) and methyl chloroformate, example 509 was prepared in an analogous series of reactions to (49a) and (92d). MS found: $(M+H)^+ = 526$.

Example 510

[1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-hydroxy-1-methyl-4-piperidineacetamide
tris(trifluoroacetate)

Beginning with the piperidine from (505a) and formaldehyde, example 506 was prepared in an analogous series of reactions to (86a) and (92d). MS found: $(M+H)^+ = 482$.

Example 511

[1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-1-dimethylcarbamyl-N-hydroxy-4-piperidineacetamide
bis(trifluoroacetate)

Beginning with the piperidine from (505a) and dimethylcarbamyl chloride, example 511 was prepared in an analogous series of reactions to (49a) and (92d). MS found: $(M+H)^+ = 539$.

Example 512

[1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-1-cyclopropanecarbonyl-N-hydroxy-4-piperidineacetamide
bis(trifluoroacetate)

Beginning with the piperidine from (505a) and cyclopropanecarbonyl chloride, example 512 was prepared in an analogous series of reactions to (49a) and (92d). MS found: $(M+H)^+ = 536$.

Example 513

[1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide
bis(trifluoroacetate)

(513a) Following a procedure analogous to (300f), the aldehyde from (300e) (8.00 g, 19.3 mmol) and D-Val-OMe were coupled to give the secondary amine as a crude material. MS found: $(M+H)^+ = 529$.

(513b) Following a procedure analogous to (300g), the crude amine from (513a) was converted to the lactam. Silica gel chromatography (ethyl acetate-hexane, 20:80 then 25:75) provided the less polar isomer (4.60 g) and the more polar isomer (3.60 g). The total yield is 85% for two steps.

(513c) Following a procedure analogous to step (3a), the less polar lactam from (513b) (4.10 g, 8.27 mmol) was hydrogenolized to give the phenol (3.30, 98%). MS found: $(M+Na)^+ = 429$.

(513d) Following a procedure analogous to step (6b), the phenol from (513c) (500 mg, 1.23 mmol) was reacted with 4-chloromethylquinoline hydrochloride to give the ether (575 mg, 85%). MS found: $(M+Na)^+ = 570$.

(513e) Following a procedure analogous to step (92d), the ester from (513d) (575 mg, 1.05 mmol) was reacted with hydroxylamine to give the hydroxamic acid (380 mg, 66%). MS found: $(M-H)^- = 547$.

(513f) Following a procedure analogous to example 117, the hydroxamic acid from (513e) (380 mg, 0.693 mmol) was reacted with trifluoroacetic acid. The material was purified by reverse phase HPLC on a Dynamax C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid bis(trifluoroacetate) (268 mg, 57%). MS found: $(M+H)^+ = 449$.

Example 514

[1(R)]-3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α -(1-methylethyl)-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (513c) and 4-chloromethyl-2,6-dimethylpyridine hydrochloride, example 514 was prepared in an analogous series of reactions to (6b), (92d) and example 117. MS found: $(M+H)^+ = 427$.

Example 515

[1(R)]-3-amino- α -cyclohexyl-N-hydroxy-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the aldehyde from (300e) and D-cyclohexylglycine methyl ester hydrochloride, example 515 was prepared in an analogous series of reactions to example 513. MS found: $(M+H)^+ = 589$.

5

Example 516

[1(R)]-3-amino- α -cyclohexyl-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

10

Beginning with the aldehyde from (300e) and D-cyclohexylglycine methyl ester hydrochloride, example 516 was prepared in an analogous series of reactions to example 513. MS found: $(M+H)^+ = 467$.

15

Example 517

3-amino- α -(1,1-dimethylethyl)-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

(517a) Following a procedure analogous to (300f), the aldehyde from (300e) (8.40 g, 20.3 mmol) and D-t-Leu-OMe were coupled to give the secondary amine as a crude material. MS found: $(M+H)^+ = 543$.

(517b) Following a procedure analogous to (300g), the crude amine from (517a) was converted to the lactam. Silica gel chromatography (ethyl acetate-hexane, 15:85 then 20:80) provided the less polar isomer (4.60 g, 45%). MS found: $(M+H)^+ = 511$.

(517c) Following a procedure analogous to step (3a), the less polar lactam from (517b) (4.50 g, 8.80 mmol) was hydrogenolized to give the phenol (3.62 g, 98%). MS found: $(M+Na)^+ = 443$.

(517d) Following a procedure analogous to step (6b), the phenol from (517c) (210 mg, 0.500 mmol) was reacted with 4-chloromethyl-2,6-dimethylpyridine hydrochloride to give the ether (240 mg, 89%). MS found: $(M+H)^+ = 540$.

(517e) The ester from (517d) (220 mg, 0.408 mmol) in concentrate HCl (5 mL) and HOAc (7.5 mL) was heated to 100 °C

overnight and concentrated to give the crude carboxylic acid.
MS found: $(M-H)^- = 424$.

(517f) The carboxylic acid from (517e), hydroxylamine hydrochloride (160 mg, 5.6 eq), NMM (0.5 mL), BOP (300 mg, 1.7 eq) in DMF (8 mL) were stirred at rt for 4 h. Following addition of sat NH_4Cl (25 mL), the mixture was extracted with ethyl acetate several times. The extracts were concentrated and purified by reverse phase HPLC on a Dynamax C-18 semiprep column eluting an acetonitrile:water:TFA gradient, to give the hydroxamic acid bis(trifluoroacetate) (140 mg, 51% for 2 steps). MS found: $(M+H)^+ = 441$.

Example 518

[1(R)]-3-amino- α -(1,1-dimethylethyl)-N-hydroxy-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (517c) and 4-chloromethylquinoline hydrochloride, example 518 was prepared in an analogous series of reactions to (6b), (517e) and (517f). MS found: $(M-H)^- = 461$.

Example 519

[1(R)]-3-amino- α -(1,1-dimethylethyl)-N-hydroxy-2-oxo-3-[4-[(2-methyl-4-quinolinyl)methoxy]phenyl]-1-pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (517c) and 4-chloromethyl-2-methylquinoline hydrochloride, example 519 was prepared in an analogous series of reactions to (6b), (517e) and (517f). MS found: $(M+H)^+ = 477$

Example 520

[1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-[(2-methyl-4-quinolinyl)methoxy]phenyl]-1-pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (513c) and 4-chloromethyl-2-methylquinoline hydrochloride, example 520 was prepared in

an analogous series of reactions to (6b), (92d) and example 117. MS found: $(M+H)^+ = 463$.

Example 521

5 [1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-
[(2,6-dimethyl-4-quinolinyl)methoxy]phenyl]-1-
pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (513c) and 4-chloromethyl-2,6-dimethylquinoline hydrochloride, example 521 was prepared in an analogous series of reactions to (6b), (92d) and example 117. MS found: $(M+H)^+ = 477$.

Example 522

15 [1(R)]-N-[4-[1-[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-2-(hydroxyamino)-2-oxoethyl]-1-piperidine]-4-morpholinecarboxamide bis(trifluoroacetate)

Beginning with the piperidine from (505a) and 4-morpholinecarbonyl chloride, example 522 was prepared in an analogous series of reactions to (49a) and (92d). MS found: $(M+H)^+ = 581$.

Example 523

25 [1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-1-(2-methyl-1-oxopropyl)-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the piperidine from (505a) and isobutyryl chloride, example 523 was prepared in an analogous series of reactions to (49a) and (92d). MS found: $(M+H)^+ = 538$.

Example 524

35 [1(R)]-3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α -(4-methoxycyclohexyl)-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

(524a) Sodium carbonate (6.13 g, 2 eq) and (BOC)₂O (6.30 g, 1 eq) were successively added to D-4-hydroxycyclohexylglycine

(5.00 g, 28.9 mmol, Ciba-Geigy, WO9722587, 1994) in water (120 mL) and dioxane (60 mL) at 0 °C. The mixture was stirred at rt overnight and then adjusted to pH 5-6 with 6 N HCl.

Following removal of dioxane, the mixture was diluted with
5 water (150 mL), acidified to pH 2-3, saturated with solid NaCl, and extracted with ethyl acetate (3x250 mL). The

combined extracts were dried (MgSO₄), and concentrated to give the BOC-protected amino acid (7.80 g, 99%). MS found: (M-H)⁻ = 272.

10 (524b) A 2.0 M hexane solution of trimethylsilyl diazomethane (18.3 mL, 1.3 eq) was added to the acid from (524a) (7.70 g, 28.8 mmol) in methanol (50 mL) and benzene (200 mL). The mixture was stirred at rt for 30 min, then concentrated.

Silica gel chromatography (ethyl acetate-hexane, 50:50) gave
15 the ester (7.40 g, 91%). MS found: (M+Na)⁺ = 310.

(524c) The ester from (524b) (7.20 g, 25.2 mmol) was stirred in 4 N dioxane solution of hydrogen chloride (200 mL) for 30 min and then concentrated to give the amino ester hydrochloride (5.70 g, 100%). MS found: (M+H)⁺ = 188.

20 (524d) Following a procedure analogous to (300f), the aldehyde from (300e) (2.00 g, 4.83 mmol) and the methyl ester hydrochloride from (525c) were coupled to give the secondary amine as a crude material. MS found: (M+H)⁺ = 585.

(524e) Following a procedure analogous to (300g), the crude
25 amine from (525d) were cyclized to give the lactam as a crude material (2.71 g). MS found: (M+Na)⁺ = 575.

(524f) Proton sponge (1.16 g, 3 eq) and trimethyloxonium tetrafluoroborate (803 mg, 3 eq) were added to the crude material from (524d) (1.00 g) in dichloromethane (20 mL).

30 After 4 h at rt, ethyl acetate (200 mL) was added. The mixture was washed with water (2x25 mL), brine (25 mL), dried (MgSO₄) and concentrated. Silica gel chromatography (35:65 then 40:60 then 45:55) gave the desired methyl ether (628 mg, 62% for 3 steps). MS found: (M+Na)⁺ = 589.

35 (524g) Following a procedure analogous to step (3a), the lactam from (524f) (838 mg, 1.48 mmol) was hydrogenolized to give the phenol (643.2 mg, 91%). MS found: (M+Na)⁺ = 499.

(524h) Following a procedure analogous to step (6b), the phenol from (524g) (200 mg, 0.420 mmol) was reacted with 4-chloromethyl-2,6-dimethylpyridine hydrochloride to give the ether (197.4 mg, 79%). MS found: $(M+Na)^+ = 619$.

5 (524i) Following a procedure analogous to step (92d), the ester from (524h) (185.4 mg, 0.311 mmol) was reacted with hydroxylamine to give the hydroxamic acid (top isomer: 67.3 mg; bottom isomer: 60.1 mg). The total yield is 127.4 mg (69%). MS found: $(M+H)^+ = 597$.

10 (524j) Following a procedure analogous to step (117), the bottom isomer of the hydroxamic acid from (524i) (56.1 mg, 0.094 mmol) was reacted with TFA to give the deprotected hydroxamic acid (68.1 mg, 100%). MS found: $(M+H)^+ = 497$.

15

Example 525

[1' (R)]-N-hydroxy-1,2-dihydro- α -(1-methylethyl)-2,2'-dioxo-6-(phenylmethoxy)spiro[3H-indole-3,3'-pyrrolidine]-1'-acetamide

(525a) Cesium carbonate (8.86 g, 2 eq) was added to a solution of dimethyl [4-(benzyloxy)-2-nitrophenyl]malonate (4.87 g, 13.6 mmol; Warpehoski, et al. *J. Med. Chem.* **1988**, 31, 590) and allyl bromide (3.53 mL, 3 eq) in DMSO at rt. After 1 h at this temperature, ether (800 mL) and sat ammonium chloride (100 mL) were added. The organic phase was separated, washed with water (3x50 mL), brine (50 mL), dried (MgSO₄) and concentrated. Silica gel chromatography (ethyl acetate-hexane, 15:85 then 20:80) provided the allylated product (5.28 g, 97%). MS found: $(M+H)^+ = 400$.

30 (525b) Following a procedure analogous to step (1c), the olefin from (219a) (5.18 g, 13.0 mmol) was degraded by ozonolysis. Silica gel chromatography (ethyl acetate-hexane, 20:80 then 30:70 then 35:65 then 40:60) provided the aldehyde (4.96 g, 95%). MS found: $(M+NH_4)^+ = 419$.

35 (525c) Following a procedure analogous to (300f), the aldehyde from (525b) (510 mg, 1.27 mmol) and D-valine methyl ester hydrochloride were coupled to give the secondary amine as a crude material.

(525d) Following a procedure analogous to (1d), the crude material from (525c) was treated with zinc in acetic acid at reflux. The crude spirolactam was purified by silica gel chromatography (ethyl acetate-hexane, 40:60 then 50:50) to give less polar isomer (180 mg) and more polar isomer (130 mg). The total yield for two steps is 310 mg (58%). MS found: $(M-H)^- = 421$.

(525e) Following a procedure analogous to step (92d), the ester from (525d) (25.5 mg, 0.060 mmol) was reacted with hydroxylamine to give the hydroxamic acid (15.2 mg, 60%). MS found: $(M-H)^- = 422$.

Example 526

[1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-hydroxy-1-(phenylcarbonyl)-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the piperidine from (505a) and benzoyl chloride, example 526 was prepared in an analogous series of reactions to (49a) and (92d). MS found: $(M+H)^+ = 572$.

Example 527

[1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-hydroxy-1-(1-oxopropyl)-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the piperidine from (505a) and propionyl chloride, example 527 was prepared in an analogous series of reactions to (49a) and (92d). MS found: $(M+H)^+ = 524$.

Example 528

[1(R)]- α -[3-amino-2-oxo-3-[4-(2-methyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-1-acetyl-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 4-chloromethyl-2-methylquinoline in step (6b) and acetyl chloride in step (49a). MS found: $(M+H)^+ = 546$.

Example 529

[1(R)]- α -[3-amino-2-oxo-3-[4-(2-methyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-N-hydroxy-1-(methylsulfonyl)-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 4-chloromethyl-2-methylquinoline in step (6b) and methanesulfonyl chloride in step (49a). MS found: $(M+H)^+ = 582$.

Example 530

[1(R)]- α -[3-amino-2-oxo-3-[4-(2-methyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-1-(2,2-dimethyl-1-oxopropyl)-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 4-chloromethyl-2-methylquinoline in step (6b) and pivoyl chloride in step (49a). MS found: $(M+H)^+ = 588$.

Example 531

[1(R)]- α -[3-amino-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-1-acetyl-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 4-chloromethylquinoline in

step (6b) and acetyl chloride in step (49a). MS found: $(M+H)^+$ = 532.

Example 532

[1(R)]- α -[3-amino-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-N-hydroxy-1-(methylsulfonyl)-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 4-chloromethylquinoline in step (6b) and methanesulfonyl chloride in step (49a). MS found: $(M+H)^+$ = 568.

Example 533

[1(R)]- α -[3-amino-2-oxo-3-[4-[(3,5-dimethoxyphenyl)methoxy]phenyl]-1-pyrrolidinyl]-1-acetyl-N-hydroxy-4-piperidineacetamide trifluoroacetate

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 3,5-dimethoxybenzyl bromide in step (6b) and acetyl chloride in step (49a). MS found: $(M+H)^+$ = 541.

Example 534

[1(R)]- α -[3-amino-2-oxo-3-[4-[(5-methyl-3-nitrophenyl)methoxy]phenyl]-1-pyrrolidinyl]-1-acetyl-N-hydroxy-4-piperidineacetamide trifluoroacetate

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 5-methyl-3-nitrobenzyl bromide in step (6b) and acetyl chloride in step (49a). MS found: $(M+H)^+$ = 540.

Example 535

[1(R)]- α -[3-amino-2-oxo-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-1-pyrrolidinyl]-1-acetyl-N-hydroxy-4-piperidineacetamide trifluoroacetate

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (61a), (117), (49a) and (92d), but using 3,5-bis(trifluoromethyl)benzene boronic acid in step (61a) and acetyl chloride in step (49a). MS found: (M+H)⁺ = 603.

Example 536

[1(R)]- α -[3-amino-2-oxo-3-[4-[(3,5-dichlorophenyl)methoxy]phenyl]-1-pyrrolidinyl]-1-acetyl-N-hydroxy-4-piperidineacetamide trifluoroacetate

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 3,5-dichlorobenzyl bromide in step (6b) and acetyl chloride in step (49a). MS found: (M+H)⁺ = 549.

Example 537

[1(R)]- α -[3-amino-2-oxo-3-[4-(6-fluoro-2-methyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-1-acetyl-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 4-chloromethyl-6-fluoro-2-methylquinoline in step (6b) and acetyl chloride in step (49a). MS found: (M+H)⁺ = 564.

Example 538

[1(R)]- α -[3-amino-2-oxo-3-[4-(7-chloro-2-methyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-1-acetyl-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

5

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 4-chloromethyl-7-chloro-2-methylquinoline in step (6b) and acetyl chloride in step (49a). MS found: $(M+H)^+ = 580$.

10

Example 539

[1(R)]- α -[3-amino-2-oxo-3-[4-(6-chloro-2-methyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-1-acetyl-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

15

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 4-chloromethyl-6-chloro-2-methylquinoline in step (6b) and acetyl chloride in step (49a). MS found: $(M+H)^+ = 580$.

20

Example 540

[1(R)]- α -[3-amino-2-oxo-3-[4-(6-methoxy-2-methyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-1-acetyl-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

25

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 4-chloromethyl-6-methoxy-2-methylquinoline in step (6b) and acetyl chloride in step (49a). MS found: $(M+H)^+ = 576$.

30

Example 541

[1(R)]- α -[3-amino-2-oxo-3-[4-(2,7-dimethyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide tris(trifluoroacetate)

35

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117) and (92d), but using 4-chloromethyl-2,7-dimethylquinoline in step (6b). MS found: $(M+H)^+ = 518$.

5

Example 542

[1(R)]- α -[3-amino-2-oxo-3-[4-(2,7-dimethyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-1-acetyl-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

10

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 4-chloromethyl-2,7-dimethylquinoline in step (6b) and acetyl chloride in step (49a). MS found: $(M+H)^+ = 560$.

15

Example 543

[1(R)]- α -[3-amino-2-oxo-3-[4-(2-methoxy-4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide tris(trifluoroacetate)

20

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117) and (92d), but using 4-bromomethyl-2-methoxyquinoline in step (6b). MS found: $(M+H)^+ = 520$.

25

Example 544

[1(R)]- α -[3-amino-2-oxo-3-[4-[(3,5-dimethoxyphenyl)methoxy]phenyl]-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide bis(trifluoroacetate)

30

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117) and (92d), but using 3,5-dimethoxybenzyl bromide in step (6b). MS found: $(M+H)^+ = 499$.

35

Example 545

[1(R)]- α -[3-amino-3-[4-[(2,6-diethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide tris(trifluoroacetate)

5

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117) and (92d), but using 4-chloromethyl-2,6-diethylpyridine (prepared from 2,6-dichloro-4-hydroxymethylpyridine following the procedure of Tamao, et al Bull. Chem. Soc. Jpn. 1976, 49, 1958 and subsequent treatment with thionyl chloride) in step (6b). MS found: (M+H)⁺ = 496.

10

Example 546

[1(R)]- α -[3-amino-3-[4-[(2,6-diethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-1-acetyl-N-hydroxy-4-piperidineacetamide tris(trifluoroacetate)

15

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117), (49a) and (92d), but using 4-chloromethyl-2,6-diethylpyridine in step (6b) and acetyl chloride in step (49a). MS found: (M+H)⁺ = 538.

20

25

Example 547

[1(R)]- α -[3-amino-2-oxo-3-[4-(7-methyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide tris(trifluoroacetate)

30

Beginning with the phenol from (501c), the title compound was prepared in an analogous series of reactions to (6b), (117) and (92d), but using 4-chloromethyl-7-methylquinoline in step (6b). MS found: (M+H)⁺ = 504.

35

Example 548

[1(R)]-3-amino-N-hydroxy- α -(4-methoxycyclohexyl)-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide bis(trifluoroacetate)

5

Beginning with the phenol from (524g), the title compound was prepared in an analogous series of reactions to (6b), (92d) and (117), but using 4-chloromethylquinoline in step (6b). MS found: $(M+H)^+ = 519$.

10

Example 549

[1(R)]-3-amino- α -(1,1-dimethylethyl)-3-[4-[(2,6-dimethyl-4-quinolinyl)methoxy]phenyl]-N-hydroxy-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

15

Beginning with the phenol from (517c), the title compound was prepared in an analogous series of reactions to (517d-f), but using 4-chloromethyl-2,6-dimethylquinoline in step (517d). MS found: $(M+H)^+ = 491$.

20

Example 550

[1(R)]-3-[4-[(2,6-dimethyl-1-oxido-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)

25

(550a) Beginning with the phenol from (6a), the picolyl ether was prepared in an analogous reaction to (6b), but using 4-chloromethyl-2,6-dimethylpyridine. MS found: $(M+H)^+ = 397$.

(550b) A mixture of the picolyl ether from (550a) (100 mg, 0.252 mmol), mCPBA (100 mg, 2 eq), and 40% aqueous HF (0.015 mL), DMF (2 mL) and methanol (0.56 mL) was stirred at rt for 2 h. The mixture was quenched with sat NaHSO₃ (1 mL) and sat Na₂CO₃, and extracted with ethyl acetate. The organic extracts were washed with Na₂CO₃ (2x), brine (2x), dried (MgSO₄) and concentrated to give the pyridine N-oxide (90 mg, 86%). MS found: $(M+H)^+ = 413$.

30
35

(550c) Following procedure analogous to (92d), the material from (550b) was converted to the hydroxamic acid. MS found: $(M+H)^+ = 414$.

5

Example 551

[1(R)]-3-amino- α -(1,1-dimethylethyl)-3-[4-[(7-chloro-2-methyl-4-quinolinyl)methoxy]phenyl]-N-hydroxy-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

10

Beginning with the phenol from (517c), the title compound was prepared in an analogous series of reactions to (517d-f), but using 7-chloro-4-chloromethyl-2-methylquinoline in step (517d). MS found: $(M+H)^+ = 511$.

15

Example 552

[1(R)]-3-amino- α -(1,1-dimethylethyl)-3-[4-[(6-fluoro-2-methyl-4-quinolinyl)methoxy]phenyl]-N-hydroxy-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

20

Beginning with the phenol from (517c), the title compound was prepared in an analogous series of reactions to (517d-f), but using 4-chloromethyl-6-fluoro-2-methylquinoline in step (517d). MS found: $(M+H)^+ = 495$.

25

Example 553

[1(R)]-3-amino- α -(1,1-dimethylethyl)-3-[4-[(6-chloro-2-methyl-4-quinolinyl)methoxy]phenyl]-N-hydroxy-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

30

Beginning with the phenol from (517c), the title compound was prepared in an analogous series of reactions to (517d-f), but using 6-chloro-4-chloromethyl-2-methylquinoline in step (517d). MS found: $(M+H)^+ = 511$.

35

Example 554

[1(R)]-3-amino- α -(1,1-dimethylethyl)-3-[4-[(6-methoxy-2-methyl-4-quinolinyl)methoxy]phenyl]-N-hydroxy-2-oxo-1-pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (517c), the title compound was prepared in an analogous series of reactions to (517d-f), but using 4-chloromethyl-6-methoxy-2-methylquinoline in step
5 (517d). MS found: $(M+H)^+ = 507$.

Example 555

[1(R)]-3-amino- α -(1,1-dimethylethyl)-3-[4-[(2,7-
dimethyl-4-quinolinyl)methoxy]phenyl]-N-hydroxy-2-oxo-
10 1-pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (517c), the title compound was prepared in an analogous series of reactions to (517d-f), but using 4-chloromethyl-2,7-dimethylquinoline in step (517d).
15 MS found: $(M+H)^+ = 491$.

Example 556

[1(R)]-3-amino- α -(1,1-dimethylethyl)-3-[4-[(7-methyl-
4-quinolinyl)methoxy]phenyl]-N-hydroxy-2-oxo-1-
20 pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (517c), the title compound was prepared in an analogous series of reactions to (517d-f), but using 4-chloromethyl-7-methylquinoline in step (517d). MS
25 found: $(M+H)^+ = 477$.

Example 557

[1(R)]-3-amino- α -cyclohexyl-N-hydroxy-2-oxo-3-[4-(2-
methyl-4-quinolinylmethoxy)phenyl]-1-
30 pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the aldehyde from (300e), the title compound was prepared in an analogous series of reactions to (513a-f), but using D-cyclohexylglycine methyl ester
35 hydrochloride in step (513a) and 4-chloromethyl-2-methylquinoline in step (513d). MS found: $(M+H)^+ = 503$.

Example 558

[1(R)]-3-amino- α -cyclohexyl-N-hydroxy-2-oxo-3-[4-(2,6-
dimethyl-4-quinolinylmethoxy)phenyl]-1-
pyrrolidineacetamide bis(trifluoroacetate)

5

Beginning with the aldehyde from (300e), the title compound was prepared in an analogous series of reactions to (513a-f), but using D-cyclohexylglycine methyl ester hydrochloride in step (513a) and 4-chloromethyl-2,6-dimethylquinoline in step (513d). MS found: $(M+H)^+ = 517$.

10

Example 559

[1(R)]-3-amino-3-[4-[(5-methyl-3-
nitrophenyl)methoxy]phenyl]-N-hydroxy- α -(1-
methylethyl)-2-oxo-1-pyrrolidineacetamide
trifluoroacetate

15

Beginning with the phenol from (513c), the title compound was prepared in an analogous series of reactions to (513d-f), but using 5-methyl-3-nitrobenzyl bromide in step (513d). MS found: $(M+H)^+ = 457$.

20

Example 560

[1(R)]-3-amino-3-[4-[3,5-
bis(trifluoromethyl)phenoxy]phenyl]-N-hydroxy- α -(1-
methylethyl)-2-oxo-1-pyrrolidineacetamide
trifluoroacetate

25

Beginning with the phenol from (513c), the title compound was prepared in an analogous series of reactions to (61a) and (513e-f), but using 3,5-bis(trifluoromethyl)benzene boronic acid in step (61a). MS found: $(M+H)^+ = 518$.

30

Example 561

[1(R)]-3-amino-3-[4-[[3,5-
bis(trifluoromethyl)phenyl]methoxy]phenyl]-N-hydroxy-
 α -(1-methylethyl)-2-oxo-1-pyrrolidineacetamide
trifluoroacetate

35

Beginning with the phenol from (513c), the title compound was prepared in an analogous series of reactions to (513d-f), but using 3,5-bis(trifluoromethyl)benzyl bromide in step (513d). MS found: $(M+H)^+ = 534$.

Example 562

[1(R)]-3-amino-3-[4-(3,5-dibromophenoxy)phenyl]-N-hydroxy- α -(1-methylethyl)-2-oxo-1-pyrrolidineacetamide
trifluoroacetate

Beginning with the phenol from (513c), the title compound was prepared in an analogous series of reactions to (61a) and (513e-f), but using 3,5-dibromobenzenboronic acid in step (61a). MS found: $(M+H)^+ = 523$.

Example 563

[1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-(6-fluoro-2-methyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (513c), the title compound was prepared in an analogous series of reactions to (513d-f), but using 4-chloromethyl-6-fluoro-2-methylquinoline in step (513d). MS found: $(M+H)^+ = 481$.

Example 564

[1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-(6-methoxy-2-methyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (513c), the title compound was prepared in an analogous series of reactions to (513d-f), but using 4-chloromethyl-6-methoxy-2-methylquinoline in step (513d). MS found: $(M+H)^+ = 493$.

Example 565

[1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-(7-chloro-2-methyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide bis(trifluoroacetate)

5

Beginning with the phenol from (513c), the title compound was prepared in an analogous series of reactions to (513d-f), but using 7-chloro-4-chloromethyl-2-methylquinoline in step (513d). MS found: $(M+H)^+ = 497$.

10

Example 566

[1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-(6-chloro-2-methyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide bis(trifluoroacetate)

15

Beginning with the phenol from (513c), the title compound was prepared in an analogous series of reactions to (513d-f), but using 6-chloro-4-chloromethyl-2-methylquinoline in step (513d). MS found: $(M+H)^+ = 497$.

20

Example 567

[1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-(2-methoxy-4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide bis(trifluoroacetate)

25

Beginning with the phenol from (513c), the title compound was prepared in an analogous series of reactions to (513d-f), but using 4-bromomethyl-2-methoxyquinoline in step (513d). MS found: $(M+H)^+ = 479$.

30

Example 568

[1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-(2,7-dimethyl-4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide bis(trifluoroacetate)

35

Beginning with the phenol from (513c), the title compound was prepared in an analogous series of reactions to (513d-f),

but using 4-chloromethyl-2,7-dimethylquinoline in step (513d).
MS found: $(M+H)^+ = 477$.

Example 569

5 [1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-
 [(2,6-diethyl-4-pyridinyl)methoxy]phenyl]-1-
 pyrrolidineacetamide bis(trifluoroacetate)

Beginning with the phenol from (513c), the title compound
10 was prepared in an analogous series of reactions to (513d-f),
but using 4-chloromethyl-2,6-diethylpyridine in step (513d).
MS found: $(M+H)^+ = 455$.

Example 700

15 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[3-
 (phenylmethoxy)phenyl]-1-pyrrolidineacetamide
 (700a) To 0.061 grams of methyl ester, obtained in a manner
analogous to examples 1a-d, in 4 mL of anhydrous methanol was
added 0.116 grams of hydroxylamine hydrochloride and 0.135
20 grams of sodium methoxide. The reaction was stirred at
ambient temperature overnight at which time it was quenched
with acetic acid and the volatiles removed under reduced
pressure. The resulting material was purified by C18 reverse
phase HPLC affording the hydroxamic acid 700. LRMS found (M-
25 H) $^- = 367$.

Example 701

[1(R)]-3-[3-[(3,5-dimethylphenyl)methoxy]phenyl]-N-
 hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide
30 (701a) Following the procedures analogous to examples 1a-d,
3a, 6b and 700a the hydroxamic acid 701 was obtained. LRMS
found $(M+H)^+ = 397$, $(M-H)^- = 395$.

Example 702

35 [1(R)]-N-hydroxy- α ,3-dimethyl-3-[3-[(3-
 methoxyphenyl)methoxy]phenyl]-2-oxo-1-
 pyrrolidineacetamide

(702a) Following the procedures analogous to examples 1a-d, 3a, 6b and 700a the hydroxamic acid 702 was obtained. LRMS found $(M-H)^- = 381$.

5

Example 703

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[3-(1-methylethoxy)phenyl]-2-oxo-1-pyrrolidineacetamide

(703a) Following the procedures analogous to examples 1a-d, 3a, 6b and 700a the hydroxamic acid 703 was obtained. LRMS found $(2M+Na)^+ = 663$.

Example 704

[1(R)]-3-[3-(heptyloxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

(704a) Following the procedures analogous to examples 1a-d, 3a, 6b and 700a the hydroxamic acid 704 was obtained. LRMS found $(M-H)^- = 375$.

Example 705

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-1,3,4-thiadiazol-2-yl-1,3-pyrrolidinediacetamide

(705a) To a stirred, cooled (-78°C) solution of 5 grams methyl ester 705 was added 1.2 eq. of lithium diisopropylamide over 10 minutes. After stirring for 1 hour at -78°C 1.7 mL of allyl bromide was added over 5 minutes. The reaction was allowed to slowly warm to ambient temperature while stirring overnight. Volatiles were removed under reduced pressure and the resulting material was diluted with ethyl acetate and washed with 1N hydrochloric acid. The aqueous phase was extracted 2 additional times with ethyl acetate. The combined organic phases were washed with brine, saturated aqueous sodium bicarbonate, brine, dried over magnesium sulfate and the volatiles were removed under reduced pressure. The resulting material was chromatographed on

silica gel eluting with 5% ethyl acetate/hexane affording 4.9 grams of 705a as a white solid. LRMS found $(M+H)^+ = 297$.

(705b) To a stirred, cooled (-78°C) solution of 5 grams (705a) was added 1.02 eq. of lithium diisopropylamide over 10 minutes. After stirring for 1 hour at -78°C 2.55 mL of t-butyl bromoacetate was added over 5 minutes. The reaction was allowed to slowly warm to ambient temperature while stirring overnight. Volatiles were removed under reduced pressure and the resulting material was diluted with ethyl acetate and washed with 1N hydrochloric acid. The aqueous phase was extracted 3 additional times with ethyl acetate. The combined organic phases were washed with brine, saturated aqueous sodium bicarbonate, brine, dried over magnesium sulfate and the volatiles were removed under reduced pressure. The resulting material was chromatographed on silica gel eluting with 5% ethyl acetate/hexane affording 5 grams of 705b as a white solid. LRMS found $(M+Na)^+ = 433$.

(705c) To 55 grams of methyl ester 705b in 600 mL of dimethyl sulfoxide, 400 mL of water and 1000 mL of methanol was added 55 grams of lithium hydroxide monohydrate. The reaction was stirred at 79°C for 3 hours. The mixture was concentrated to about half original volume and poured into ice. The mixture was acidified with 1N hydrochloric acid and extracted 4 times with diethyl ether. The combined ether extracts were washed three times with water, twice with brine and dried over magnesium sulfate. The volatiles were removed under reduced pressure and the resulting material was recrystallized from acetone/hexane affording 45 grams of the acid 705c as a white solid. LRMS found $(M+Na)^+ = 419$.

(705d) To 1.3 grams of acid 705c in 20 mL of N,N-dimethylformamide was added 1.44 mL of 4-methylmorpholine and 1.44 grams of O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate. After stirring 30 minutes 0.46 grams of D-alanine methyl ester hydrochloride was added. The reaction was stirred 18 hours at ambient temperature and for 45 minutes at 60°C . The volatiles were removed under reduced pressure and the resulting material was

partitioned in ethyl acetate and washed with 1N hydrochloric acid saturated with sodium chloride. The aqueous phase was extracted another two times with ethyl acetate. The combined organic phases were washed with brine, saturated aqueous sodium bicarbonate, brine, dried over magnesium sulfate and the volatiles were removed under reduced pressure. The resulting material was chromatographed on silica gel eluting with 25% ethyl acetate/hexane affording 1.6 grams of 705d. LRMS found $(M+Na)^+ = 504$.

(705e) To a stirred, cooled (-78°C) solution of 0.90 grams of 705d in 20 mL of dichloromethane was bubbled ozone until the mixture attained a blue color. Ozone was added for an additional 10 minutes followed by a 15 minute oxygen flush. To this material was added 0.54 grams of triphenylphosphine and the reaction was allowed to slowly warm to ambient temperature while stirring 48 hours. The volatiles were removed under reduced pressure and the resulting material was chromatographed on silica gel eluting with a gradient of 25% ethyl acetate/hexane to 50% ethyl acetate/hexane affording 0.620 grams of 705e as a viscous oil. LRMS found $(M+Na)^+ = 506$.

(705f) To a stirred cooled (-20°C : carbon tetrachloride/dry ice) solution of 14.1 grams of 705e in 500 mL of dichloromethane was added 23.3 mL of triethylsilane and 11.2 mL of trifluoroacetic acid. The reaction was stirred 1 hour at 0°C and 2 hours at room temperature. The reaction was made basic by the addition of saturated aqueous sodium bicarbonate and partitioned with chloroform. The aqueous was extracted 3 more times with chloroform. The combined organic phases were washed with brine, dried over magnesium sulfate and the volatiles were removed under reduced pressure affording 11.3 grams of 705f. LRMS found $(M+Na)^+ = 490$.

(705g) To 3 grams of 705f in 20 mL of methanol was added 0.30 grams of 10% palladium on carbon. The reaction was stirred 3 hours under hydrogen (balloon). The catalyst was filtered through a 0.45 μm PTFE filter and the volatiles were removed

under reduced pressure affording 2.4 grams of phenol 705g.
LRMS found $(M+Na)^+ = 400$.

(705h) To 1.2 grams of 705g in 20 mL of DMSO was added 1.54 grams of 3-bromomethyl 2,5-dichloropyridine and 2.32 grams of cesium carbonate. After stirring for two hours at ambient temperature the reaction was diluted with diethyl ether and washed with brine. The aqueous was extracted an additional three times with ether. All organics were combined and washed with saturated aqueous sodium bicarbonate, water, brine, dried over magnesium sulfate and the volatiles were removed under reduced pressure. The resulting material was chromatographed on silica gel eluting with 2% methanol/chloroform affording 1.1 grams of 705h. LRMS found $(M+H)^+ = 481$.

(705i) To 1.1 grams of 705h in 50 mL of dichloromethane was added 10 mL of trifluoroacetic acid. After stirring 3 hours the volatiles were removed under reduced pressure affording 1 gram of 705i. LRMS found $(M+Na)^+ = 503$.

(705j) To 0.50 grams of 705i in 20 mL of N,N-dimethylformamide was added 0.46 mL of 4-methylmorpholine, 0.315 grams of 2-amino-1,3,4-thiadiazole and 0.474 grams of O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate. After stirring 10 hours at room temperature the reaction was heated at 60°C for 45 minutes. The volatiles were removed under reduced pressure and the resulting material was diluted with ethyl acetate and washed with 1N hydrochloric acid saturated with sodium chloride. The aqueous was extracted 3 times with ethyl acetate and all the organics were combined and extracted with brine, saturated aqueous sodium bicarbonate, brine, dried over magnesium sulfate, and the volatiles were removed under reduced pressure affording 0.60 grams of 705j. LRMS found $(M-H)^- = 562$.

(705k) To 0.55 grams of 705j in 20 mL of 1:1 tetrahydrofuran/water was added 0.12 grams of lithium hydroxide monohydrate. After stirring 3 hours at ambient temperature the reaction volume was reduced by half under reduced pressure, diluted with water and washed twice with diethyl ether. The ether phases were combined and extracted

twice with water. All aqueous phases were combined, acidified with 1N hydrochloric acid and extracted 3 times with ethyl acetate. The combined ethyl acetate extracts were washed with water, brine, dried over magnesium sulfate and the volatiles were removed under reduced pressure affording 0.52 grams of 705k. LRMS found $(M-H)^- = 548$.

(7051) To 0.40 grams of 705k in 20 mL of N,N-dimethylformamide was added 0.8 mL of 4-methyl morpholine, 0.202 grams of hydroxylamine hydrochloride and 0.354 grams of benzotriazol-1-yl-oxy-tris(dimethylamino)phosphonium-hexafluorophosphate. After stirring overnight at ambient temperature the volatiles were removed under reduced pressure and the resulting material was separated on C18 reverse phase HPLC isolating 0.18 grams of faster isomer 7051. LRMS found $(M-H)^- = 563$.

Example 706

[1(R)]-1,1-dimethylethyl 1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-[4-(phenylmethoxy)phenyl]-3-pyrrolidineacetate

(706a) Following the procedures analogous to examples 705a-j and 700a the hydroxamic acid 706 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found $(M-H)^- = 467$, $(M+H)^+ = 469$

Example 707

[1(R)]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-[4-(phenylmethoxy)phenyl]-3-pyrrolidineacetic acid

(707a) To 0.015 grams of hydroxamic acid 706 in 3 mL of dichloromethane was added 0.5 mL of trifluoroacetic acid. After stirring one hour the volatiles were removed under reduced pressure affording 0.009 grams of 707. LRMS found $(M+Na)^+ = 435$, $(M-H)^- = 411$

Example 708

[1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-N3-[2-(methylamino)-2-oxoethyl]-2-oxo-1,3-pyrrolidinediacetamide

(708a) Following the procedures analogous to examples 705a-j and 700a the hydroxamic acid 708 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found (M+Na)⁺ = 533.

Example 709

10 [1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide

(709a) Following the procedures analogous to examples 705a-j and 700a the hydroxamic acid 709 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found (M-H)⁻ = 521.

Example 710

20 [1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy- α -methyl-3-[2-(4-morpholinyl)-2-oxoethyl]-2-oxo-1-pyrrolidineacetamide

(710a) Following the procedures analogous to examples 705a-j and 700a the hydroxamic acid 710 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found (M+Na)⁺ = 532.

Example 711

30 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide mono(trifluoroacetate)

(711a) Following the procedures analogous to examples 705a-j and 700a the hydroxamic acid 711 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found (M+H)⁺ = 564.

Example 712

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-
5 N3-[2-(4-morpholinyl)ethyl]-1,3-pyrrolidinediacetamide)
bis(trifluoroacetate)

(712a) Following the procedures analogous to examples 705a-j and 700a the hydroxamic acid 712 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found (M+H)⁺ =

10 594

Example 713

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-
15 N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide
bis(trifluoroacetate)

(713a) Following the procedures analogous to examples 705a-j and 700a the hydroxamic acid 713 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found (M+Na)⁺ =

20 594

Example 714

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-
25 N3-2-thiazolyl-1,3-pyrrolidinediacetamide
mono(trifluoroacetate)

(714a) Following the procedures analogous to examples 705a-j and 700a the hydroxamic acid 714 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found (M+H)⁺ =

30

524.

Example 715

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-
35 N3-(3-pyridinylmethyl)-1,3-pyrrolidinediacetamide
mono(trifluoroacetate)

(715a) Following the procedures analogous to examples 705a-j and 700a the hydroxamic acid 715 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found $(M+Na)^+ = 594$.

5

Example 716

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-(2-pyridinylmethyl)-1,3-pyrrolidinediacetamide
mono(trifluoroacetate)

10

(716a) Following the 716 analogous to examples 705a-j and 700a the hydroxamic acid 706 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found $(M+H)^+ = 572$.

15

Example 717

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-4-pyridinyl-1,3-pyrrolidinediacetamide
mono(trifluoroacetate)

20

(717a) Following the procedures analogous to examples 705a-j and 700a the hydroxamic acid 717 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found $(M+H)^+ = 558$.

25

Example 718

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-N3-(3-methyl-5-isothiazolyl)-2-oxo-1,3-pyrrolidinediacetamide

30

(718a) Following the procedures analogous to examples 705a-1 the hydroxamic acid 718 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found $(M-H)^- = 576$.

Example 719

35

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N3-[5-(1,1-dimethylethyl)-

1,3,4-thiadizol-2-yl]-N1-hydroxy- α 1-methyl-2-oxo-1,3-pyrrolidinediacetamide

(719a) Following the procedures analogous to examples 705a-1
5 the hydroxamic acid 719 was obtained and isolated as the
faster isomer by reverse phase HPLC. LRMS (M-H)⁻ = 619.

Example 720

**[1(R)]-1,1-dimethylethyl 2-[[[3-[4-[(2,6-dichloro-4-
10 pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-
methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]acetyl]amino]-
4-thiazoleacetate**

(720a) Following the procedures analogous to examples 705a-1
the hydroxamic acid 720 was obtained and isolated as the
15 faster isomer by reverse phase HPLC. LRMS (M-H)⁻ = 676.

Example 721

**[1(R)]-2-[[[3-[4-[(2,6-dichloro-4-
pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-
20 methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]acetyl]amino]-
4-thiazoleacetic acid**

(721a) Following the procedures analogous to examples 705a-1
and 707a the hydroxamic acid 721 was obtained and isolated as
the faster isomer by reverse phase HPLC. LRMS found (M-H)⁻ =
25 620.

Example 722

**[1(R)]-3-[4-[(2,6-dichloro-4-
pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-N3-[4-
30 [2-(methylamino)-2-oxoethyl]-2-thiazolyl]-2-oxo-1,3-
pyrrolidinediacetamide**

(722a) Following the procedures analogous to examples 705a-j,
707a,705j-1 the hydroxamic acid 722 was obtained and isolated
as the faster isomer by reverse phase HPLC. LRMS found
35 (M+Na)⁺ = 657.

Example 723

[1(R)]-3-(1H-benzimidazol-2-ylmethyl)-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α -methyl-2-oxo-1-pyrrolidineacetamide

(723a) To 0.20 grams of acid obtained by procedures analogous to 705a-i in 5 mL of N,N-dimethylformamide was added 0.18 mL of 4-methyl morpholine, 0.135 grams of phenyldiamine, and 0.173 grams of O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate. After stirring for 12 hours at ambient temperature the volatiles were removed under reduced pressure and the resulting material was washed with brine and 1 mL of 10% aqueous citric acid. The aqueous was extracted twice with ethyl acetate and the combined organic phases were washed with brine, saturated aqueous sodium bicarbonate, brine, dried over magnesium sulfate and the volatiles were removed under reduced pressure affording 723a. LRMS found (M+H)⁺ = 571.

(723b) To 0.20 grams of 723a in 40 mL of 1:1 tetrahydrofuran/acetic acid was heated to reflux for 1.5 hours. The volatiles were removed under reduced pressure and the resulting material was dissolved in ethyl acetate and washed with water. The aqueous phase was extracted twice with ethyl acetate and the combined organic phases were washed with water, saturated aqueous sodium bicarbonate, brine, dried over magnesium sulfate and the volatiles were removed under reduced pressure affording 0.17 grams of 723b. LRMS found (M+H)⁺ = 553.

(723c) To 0.15 grams of 723b in 6 mL of 1:1 tetrahydrofuran/water was added 0.065 grams of lithium hydroxide monohydrate. After stirring for two hours at ambient temperature the volatiles were removed under reduced pressure and the resulting material was dissolved in ethyl acetate and washed 1N hydrochloric acid. The aqueous was extracted twice with ethyl acetate and the combined organic phases were washed with brine, dried over magnesium sulfate and the volatiles were removed under reduced pressure affording 0.11 grams of 723c. LRMS found (M+H)⁺ = 539.

(723d) Following the procedure analogous to 7051 the hydroxamic acid 723d was obtained and isolated as the faster isomer by C18 reverse phase HPLC. LRMS found (M+H)⁺ = 554.

5

Example 724

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-(3H-imidazo(4,5-c)pyridin-2-ylmethyl)- α -methyl-2-oxo-1-pyrrolidineacetamide

10 (724a) Following the procedures analogous to examples 723a-d the hydroxamic acid 724 was obtained and isolated as the faster isomer by reverse phase HPLC. LRMS found (M+H)⁺ = 555.

Example 725

15 [1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide

(725a) Following the procedures analogous to examples 705a-g, 61a, and 705i-1, the hydroxamic acid 725 was obtained and
20 isolated as the faster isomer by reverse phase HPLC. LRMS found (M-H)⁻ = 615.

Example 726

[1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide mono(trifluoroacetate)

(726a) Following the procedures analogous to examples 705a-g, 61a, and 705i-1, the hydroxamic acid 726 was obtained and
30 isolated as the faster isomer by reverse phase HPLC. LRMS found (M+H)⁺ = 625.

Example 780

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-(1-methylethyl)-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide

35

(780a) A 1.0 M tetrahydrofuran solution of sodium bis(trimethylsilyl)amide (192.5 mL, 1.1 eq) was added over 30 min to methyl 4-benzyloxyphenylacetate (44.95 g, 175 mmol) in tetrahydrofuran (900 mL) at -78 °C. After 1 h at -78 °C, DMPU (52.9 mL, 2.5 eq) was added over 15 min. The cold bath was replaced with an ice-water bath, and 2-benzyloxyethyl iodide (50.45 g, 1.1 eq) in THF (40 mL) was added dropwise. After 2 h at 0 °C, sat ammonium chloride (500 mL) was added. Following removal of THF in vacuo, the residue was diluted with water (250 mL) and extracted with 1:2 mixture of ether-hexane (3x500 mL). The combined extracts were washed with water (2x100 mL), brine (100 mL), dried (MgSO₄) and concentrated. The residue was filtered through a silica gel pad and the filter cake rinsed with ethyl acetate-hexane (20:80) until free of product. The filtrate was concentrated and used in the next step without purification. MS found: (M+H)⁺ = 391.

(780b) Following a procedure analogous to (1a), the crude material from (780a) was reacted with allyl bromide. The crude material was used in the next step without purification. MS found: (M+H)⁺ = 431.

(780c) Following a procedure analogous to (1c), the crude material from (780b) was ozonolized. Silica gel chromatography (ethyl acetate-hexane, 15:85 then 20:80 then 25:75) gave the desired aldehyde (43.27 g, 57% for three steps). MS found: (M+H)⁺ = 433.

(780d) Following a procedure analogous to (1d), the aldehyde from (780c) (3.00 g, 6.94 mmol) and D-valine ethyl ester hydrochloride was condensed to give the lactam (2.50 g, 68%) as a 1:1 mixture of two isomers. MS found: (M+H)⁺ = 530.

(780e) Following a procedure analogous to step (3a), the lactam from (780d) (4.50 g, 8.51 mmol) was hydrogenolized to give the phenol (2.30 g, 77%). MS found: (M+H)⁺ = 350.

(780f) Following a procedure analogous to step (6b), the phenol from (780e) (975 mg, 2.79 mmol) was reacted with 4-chloromethyl-2,6-dimethylpyridine hydrochloride to give the picolyl ether (818 mg, 62%). MS found: (M+H)⁺ = 455.

(780g) Ruthenium chloride monohydrate (18 mg, 0.05 eq) was added to a mixture of the picolyl ether from (780f) (790 mg, 1.69 mmol), sodium periodate (1.44 g, 4 eq), acetonitrile (2 mL), carbon tetrachloride (2 mL) and water (3.5 mL). After 5 h at rt, the mixture was extracted with chloroform (3x). The extracts were washed with brine, dried (MgSO₄) and concentrated to give the crude carboxylic acid (710 mg). MS found: (M+H)⁺ = 469.

(780h) Following a procedure analogous to step (705j), the carboxylic acid from (780g) (218 mg, 0.452 mmol) was coupled with 4-picolyamine to give the amide (179 mg, 69%). MS found: (M+H)⁺ = 573.

(780i) Following a procedure analogous to step (92d), the ester from (780h) was reacted with hydroxylamine to give the desired hydroxamic acid (40 mg, 23%). MS found: (M+H)⁺ = 560.

Example 781

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-α1-(1-methylethyl)-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide

Beginning with the phenol from (780e) and 4-bromomethyl-2,6-dichloropyridine, example 781 was prepared in an analogous series of reactions to (780f-i). MS found: (M+H)⁺ = 600.

Example 782

[1(R)]-α1-(cyclohexylmethyl)-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide

Beginning with the aldehyde from (780c) and D-cyclohexylmethylglycine methyl ester, example 782 was prepared in an analogous series of reactions to (780d-i). MS found: (M+H)⁺ = 614.

Example 783

[1(R)]- α 1-(cyclohexylmethyl)-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide

Beginning with the aldehyde from (780c) and D-cyclohexylmethylglycine methyl ester and using 4-bromomethyl-2,6-dichloropyridine in place of 4-chloromethyl-2,6-dimethylpyridine, example 783 was prepared in an analogous series of reactions to (780d-i). MS found: $(M+H)^+ = 654$.

Example 784

[1(R)]-1,1-dimethylethyl [5-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-3-[2-oxo-2-[(4-pyridinylmethyl)amino]ethyl]-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]carbamate

Following a sequence analogous to example 705, example 784 was prepared. MS found: $(M+H)^+ = 689$.

Example 785

[1(R)]- α 1-(4-aminobutyl)-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide tris(trifluoroacetate)

Example 785 was prepared from example 784 following a procedure similar to example 117. MS found: $(M+2H)^{2+} = 590$.

Example 800

[1(R)]-3-[3-(1H-benzotriazol-1-ylmethoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide

(800a) To 0.090 grams of methyl ester, obtained in a manner analogous to examples 1a-d, in 1 mL of anhydrous methanol was added 0.153 grams of hydroxylamine hydrochloride and 0.18 grams of sodium methoxide. The reaction was stirred at room temperature overnight at which time it was quenched with hydrochloric acid and the volatiles were removed under reduced pressure. The resulting material was purified by reverse phase HPLC affording the hydroxamic acid 800. LRMS found $(M-H)^- = 408$.

Example 801

[1(R)]-N-hydroxy-3,4,4-trimethyl- α -[3-methyl-2-oxo-3-[4-(phenylmethoxy)phenyl]-1-pyrrolidinyl]-2,5-dioxo-1-imidazolidinepropanamide

(801a) Following the procedures analogous to examples 1a-d, 6b and 800a the hydroxamic acid 801 was obtained. LRMS found $(M+H)^+ = 509$, $(M-H)^- = 507$ $(M+Na)^+ = 531$

Example 802

[1(R)]-1,1-dimethylethyl 1-[(hydroxyamino)carbonyl]-3-methylbutyl]-2-oxo-3-[4-(phenyl)-3-pyrrolidineacetate

(802a) Following the procedures analogous to examples 705a-f and 1e the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found $(M-H)^- = 509$, $(M+H)^+ = 511$, $(M+Na)^+ = 533$.

Example 803

[1(R)-N1-hydroxy-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N3-[2-(methylamino)-2-oxoethyl]- α -(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide

(803a) Following the procedures analogous to examples 705a-j and 1e the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found $(M+Na)^+ = 533$, $(M-H)^- = 551$, $(M+H)^+ = 553$.

Example 804

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-N3-[2-(methylamino)-2-oxoethyl]- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide

(804a) Following the procedures analogous to examples 705a-j and 1e the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found $(M+H)^+ = 595$.

Example 805

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide

(805a) Following the procedures analogous to examples 705a-j and 1e the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 607, (M-H)- = 605.

Example 806

[1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-N1-hydroxy-N3-[2-(methylamino)-2-oxoethyl]- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide

(806a) Following the procedures analogous to examples 705a-g, 61a, 705i, 705j, and 1e the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 647, (M-H)- = 645, (M+Na)+ = 669.

Example 807

[1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide mono(trifluoroacetate)

(807a) Following the procedures analogous to examples 705a-g, 61a, 705i, 705j, and 1e the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 667.

Example 808

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-N3-phenyl-1,3-pyrrolidinediacetamide

(808a) Following the procedures analogous to examples 705a-1 the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 600.

5

Example 809

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-N3-methyl- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide mono(trifluoroacetate)

10

(809a) Following the procedures analogous to examples 705a-1 the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 497.

15

Example 810

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-N3-[2-(1H-imidazol-4-yl)ethyl]- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide bis(trifluoroacetate)

20

(810a) Following the procedures analogous to examples 705a-1 the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 577.

25

Example 811

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-N3-[1-(phenylmethyl)-4-piperidinyl]-1,3-pyrrolidinediacetamide bis(trifluoroacetate)

30

(811a) Following the procedures analogous to examples 705a-1 the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 656.

35

Example 812

[1(R)]-N3-[2-(dimethylamino)ethyl]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide bis(trifluoroacetate)]

5

(812a) Following the procedures analogous to examples 705a-1 the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 554.

10

Example 813

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-N3-(4-hydroxyphenyl)- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide mono(trifluoroacetate)]

15

(813a) Following the procedures analogous to examples 705a-1 the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 575.

20

Example 814

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N3-hydroxy- α 1-(2-methylpropyl)-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide mono(trifluoroacetate)]

25

(814a) Following the procedures analogous to examples 705a-1 the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 566

30

Example 815

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N3-hydroxy-3-(2-hydroxyethyl)- α 1-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide mono(trifluoroacetate)]

35

(815a) Following the procedures analogous to examples 780 the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 470.

5

Example 816

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N3-(4,5-dimethyl-2-thiazolyl)-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide mono(trifluoroacetate)

10

(816a) Following the procedures analogous to examples 705a-1 the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 594.

15

Example 817

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-N3-1H-indazol-5-yl- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide mono(trifluoroacetate)

20

(817a) Following the procedures analogous to examples 705a-1 the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 599.

25


Example 818

[1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide

30

(818a) Following the procedures analogous to examples 705a-g, 61a, and 705i-1 the hydroxamic acid was obtained and isolated by reverse phase HPLC. LRMS found (M+H)+ = 659.

TABLE 1



Ex #	R ¹	R ²	R ³	MS
1	Me	Me	4-(phenylmethoxy)phenyl	367
2	Me	Me	4-methoxyphenyl	291
3	Me	Me	4-(1-isopropoxy)phenyl	319
4	Me	Me	4-(t-butoxy)phenyl	333
5	Me	Me	4-cyclohexyloxyphenyl	359
6	Me	Me	4-[[4-(t-butyl)phenyl]methoxy]phenyl	423
7	Me	Me	4-[(3-phenyl-2-propen-1-yl)oxy]phenyl	393
8	Me	Me	4-[(3-methylphenyl)methoxy]phenyl	381
9	Me	Me	4-[(3,5-dimethylphenyl)methoxy]phenyl	395
10	Me	Me	4-allyloxyphenyl	317
11	Me	Me	4-[(3-cyanophenyl)methoxy]phenyl	392
12	Me	Me	4-[(2-nitrophenyl)methoxy]phenyl	412
13	Me	Me	4-[(4-nitrophenyl)methoxy]phenyl	412
14	Me	Me	4-[(3-nitrophenyl)methoxy]phenyl	412
15	Me	Me	4-[(2-naphthalenyl)methoxy]phenyl	417
16	Me	Me	4-hydroxyphenyl	277
17	Me	Me	4-[(2-pyridinyl)methoxy]phenyl	368
18	Me	Me	4-[(3-pyridinyl)methoxy]phenyl	368
19	Me	Me	4-[(4-pyridinyl)methoxy]phenyl	368
20	Me	Me	4-(i-Bu)phenyl	317
21	Me	Me	phenyl	261
22	Me	Me	phenyl	233
23	H	H	phenyl	247
24	H	Me	phenyl	247
25	Me	H	4-methoxyphenyl	277
26	Me	H	cyclohexyl	267
27	Me	Me	2-phenylethyl	289
28	Me	Me	2-cyclohexylethyl	295
29	Me	Me	phenyl	337
30			see structure at bottom	287
31	Me	Me	4-[(3,5-dibromophenyl)methoxy]phenyl	523

32	Me	Me	4-[[3,5-bis(trifluoromethyl)phenyl]methoxy]phenyl	503
33	Me	Me	4-[(3,5-dichlorophenyl)methoxy]phenyl	435
34	Me	Me	4-[(2-methyl-1-naphthalenyl)methoxy]phenyl	455
35	Me	Me	4-[(3,5-dimethoxyphenyl)methoxy]phenyl	427
36	Me	Me	4-[[4-chloro-2-(trifluoromethyl)-6-quinoliny]methoxy]phenyl	520
37	Me	Me	4-[[4-(1,2,3-thiadiazol-4-yl)phenyl]methoxy]phenyl	451
38	Me	Me	4-([1,1'-biphenyl]-2-ylmethoxy)phenyl	443
39	Me	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	436
40	Me	Me	4-(1H-benzotriazol-1-ylmethoxy)phenyl	408
41	Me	Me	4-[(4,6-dimethyl-2-pyrimidinyl)methoxy]phenyl	397
42	Me	Me	4-(1,3-benzodioxol-5-ylmethoxy)phenyl	411
43	Me	Me	4-[(2-chloro-6-ethoxy-4-pyridinyl)methoxy]phenyl	446
44	Me	Me	4-(4-quinoliny)methoxy]phenyl	420
45	Me	Me	4-[(4,5-dimethyl-2-thiazolyl)methoxy]phenyl	402
46	Me	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	398
47	Me	Me	4-[(3-methyl-5-nitrophenyl)methoxy]phenyl	426
48	Me	Me	4-[(3-amino-5-methylphenyl)methoxy]phenyl	396
49	Me	Me	4-[[3-(acetyl amino)-5-methylphenyl]methoxy]phenyl	438
50	Me	Me	4-[[3-[[[(t-butoxy)carbonyl]amino]acetyl]amino]-5-methylphenyl]methoxy]phenyl	553
51	Me	Me	4-[[3-[(aminoacetyl)amino]-5-methylphenyl]methoxy]phenyl	455
52	Me	Me	4-[[3-[[[[[(t-butoxy)carbonyl]amino]acetyl]amino]acetyl]amino]-5-methylphenyl]methoxy]phenyl	634
53	Me	Me	4-[[3-[[[(aminoacetyl)amino]acetyl]amino]-5-methylphenyl]methoxy]phenyl	512
54	Me	Me	4-[[3-[[[(4-morpholinyl)carbonyl]amino]-5-methylphenyl]methoxy]phenyl	509
55			see structure at bottom	479
56	Me	Me	[1,1'-biphenyl]-4-yl	339
57	Me	Me	2'-methyl[1,1'-biphenyl]-4-yl	353

58	Me	Me	4'-methyl[1,1'-biphenyl]-4-yl	353
59	Me	Me	3',4'-dimethoxy[1,1'-biphenyl]-4-yl	397
60	Me	Me	2'-(trifluoromethyl)[1,1'-biphenyl]-4-yl	405
61	Me	Me	4-(4-methylphenoxy)phenyl	367
62	Me	Me	4-phenoxyphenyl	353
63	Me	Me	4-(2-methylphenoxy)phenyl	367
64	Me	Me	4-(3,5-dichlorophenoxy)phenyl	421
65	Me	Me	4-(3,4-dimethoxyphenoxy)phenyl	413
66	Me	Me	4-(1,3-benzodioxol-5-yloxy)phenyl	397
67	Me	Me	4-[3-(i-Pr)phenoxy]phenyl	395
68	Me	Me	4-(3-methoxyphenoxy)phenyl	383
69	Me	Me	4-(3-thienyloxy)phenyl	359
70	Me	Me	4-(3,4,5-trimethoxyphenoxy)phenyl	443
71	Me	Me	4-[3,5-bis(trifluoromethyl)phenoxy]phenyl	491
72	Me	Me	4-(1-naphthalenyloxy)phenyl	405
73	Me	Me	4-[3-[(hydroxyimino)methyl]phenoxy]phenyl	398
74	Me	Me	4-[4-[1-(hydroxyimino)ethyl]phenoxy]phenyl	410
75	Me	Me	4-([1,1'-biphenyl]-4-yloxy)phenyl	431
76	Me	Me	4-(3,5-dibromophenoxy)phenyl	510
77	Me	Me	4-[3-(acetylamino)phenoxy]phenyl	412
78	Me	Me	4-(4-nitrophenoxy)phenyl	398
79	Me	Me	4-methylphenyl	275
80	Me	Me	4-[(2,6-dimethyl-4-pyridinyl)oxy]methylphenyl	398
81	Me	Me	4-[(4-quinolinyl)oxy]methylphenyl	420
82	Me	Me	4-nitrophenyl	306
83	Me	Me	4-[(phenylcarbonyl)amino]phenyl	380
84	Me	Me	4-[(phenylsulfonyl)amino]phenyl	440
85	Me	Me	4-[(phenylamino)carbonyl]amino]phenyl	419
86	Me	Me	4-[(1-naphthalenylmethyl)amino]phenyl	440
87	Me	Me	4-[(4-quinolinylmethyl)amino]phenyl	419
88	Me	Me	4-[(3,5-dimethoxyphenyl)methyl]amino]phenyl	426
89	H	Me	4-[(3,5-dimethylphenyl)methoxy]phenyl	405
90	H	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	424
91	H	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	384

92	i-Pr	Me	4-(4-quinolinylmethoxy)phenyl	446
93	i-Pr	Me	4-(phenylmethoxy)phenyl	395
94	i-Pr	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	426
95	i-Bu	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	440
96	i-Bu	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	479
97	i-Bu	Me	4-[(3,5-bis(trifluoromethyl)phenyl)methoxy]phenyl	454
98	i-Bu	Me	4-[(3,5-dichlorophenyl)methoxy]phenyl	479
99	i-Bu	Me	3-(phenylmethoxy)propyl	375
101	i-Bu	Me	2-methyl-4-(phenylmethoxy)phenyl	423
102	i-Bu	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]-2-methylphenyl	492
103	i-Bu	Me	2-methyl-4-(2-naphthalenylmethoxy)phenyl	475
104	i-Bu	Me	2-methyl-4-(4-pyridinylmethoxy)phenyl	426
105	i-Bu	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]-2-methylphenyl	454
106	CH ₃ SCH ₂ CH ₂	Me	4-(phenylmethoxy)phenyl	427
107			see structure at bottom	492
108	CH ₃ SO ₂ -CH ₂ CH ₂	Me	4-[3,5-bis(trifluoromethyl)phenoxy]phenyl	581
109	CH ₃ SO ₂ -CH ₂ CH ₂	Me	4-(3,5-dibromophenoxy)phenyl	603
110	CH ₃ SO ₂ -CH ₂ CH ₂	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	528
111	CH ₃ SO ₂ -CH ₂ CH ₂	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	490
112	CH ₃ SO ₂ -CH ₂ CH ₂	Me	4-(4-quinolinylmethoxy)phenyl	512
113			see structure at bottom	379
114	(4-HO-phenyl)CH ₂	Me	4-(phenylmethoxy)phenyl	395
115	HOCH ₂ CH ₂	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	466
116	4-[(CH ₃) ₃ CO-C(O)NH ₂]butyl	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	593
117	4-aminobutyl	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	495
118	4-(acetyl-amino)butyl	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	535
119	4-[3-pyridinyl-C(O)NH]butyl	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	600

120	4-[4-morpholinyl C(O)NH] butyl	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	630
121	4-[CH ₃ SO ₂ - amino]butyl	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	595
122	4-(acetyl- amino)butyl	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	497
123	4- [(CH ₃) ₃ CO- C(O)NH] butyl	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	555
124	4- aminobutyl	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	455
125	4- [H ₂ NCH ₂ C(O) -NH]butyl	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	512
126	4-(acetyl- amino)butyl	Me	4-[[3,5-bis(trifluoromethyl) phenyl]methoxy]phenyl	626
127	4- [(CH ₃) ₃ CO- C(O)NH] butyl	Me	4-(3,5-dibromophenoxy)phenyl	=668
128	4- aminobutyl	Me	4-(3,5-dibromophenoxy)phenyl	570
129	2- [(CH ₃) ₃ CO- C(O)NH] ethyl	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	565
130	2- aminoethyl	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	467
131	2-(acetyl- amino)ethyl	Me	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	508
132	2- [(CH ₃) ₃ CO- C(O)NH] ethyl	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	527
133	2- aminoethyl	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	427
134	2-[3-pyridinyl- C(O)NH] ethyl	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	523
135	2-[4-morpholinyl C(O)NH] ethyl	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	540
136	2- [(CH ₃) ₃ CO- C(O)NHCH ₂ - C(O)NH] ethyl	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	584
137	2- [H ₂ NCH ₂ C(O) -NH]ethyl	Me	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	484

138	2- [(CH ₃) ₃ CO- C(O)NHCH ₂ - C(O)NH] ethyl	Me	4-[(2,6-dimethyl-4- pyridinyl)methoxy]phenyl	641
139	2- [H ₂ NCH ₂ C(O)- NHCH ₂ C(O)- NH]ethyl	Me	4-[(2,6-dimethyl-4- pyridinyl)methoxy]phenyl	541
140	phenyl- CH ₂ OCH ₂	Me	4-(phenylmethoxy)phenyl	473
141	HOCH ₂	Me	4-[(2,6-dichloro-4- pyridinyl)methoxy]phenyl	437
142	1- [(CH ₃) ₃ CO- C(O)]-4- piperidinyl	Me	4-(4- quinolinylmethoxy)phenyl	589
143	4- piperidinyl	Me	4-(4- quinolinylmethoxy)phenyl	489
144	1-(CH ₃ SO ₂)- 4- piperidinyl	Me	4-(4- quinolinylmethoxy)phenyl	567
145	1-[(2- furanyl) C(O)]-4- piperidinyl	Me	4-(4- quinolinylmethoxy)phenyl	583
146	1- [(CH ₃) ₃ CO- C(O)]-4- piperidinyl	Me	4-[(2,6-dimethyl-4- pyridinyl)methoxy]phenyl	567
147	4- piperidinyl	Me	4-[(2,6-dimethyl-4- pyridinyl)methoxy]phenyl	467
148	1- (CH ₃ C(O))- 4- piperidinyl	Me	4-[(2,6-dimethyl-4- pyridinyl)methoxy]phenyl	525
149	1-(CH ₃ SO ₂)- 4- piperidinyl	Me	4-[(2,6-dimethyl-4- pyridinyl)methoxy]phenyl	545
150	1-acetyl-4- piperidinyl	Me	4-[(2,6-dimethyl-4- pyridinyl)methoxy]phenyl	509
151	1-(2,2- dimethyl-1- oxopropyl)- 4- piperidinyl	Me	4-[(2,6-dimethyl-4- pyridinyl)methoxy]phenyl	551
152	1-methyl-4- piperidinyl	Me	4-[(2,6-dimethyl-4- pyridinyl)methoxy]phenyl	481
153	1-(i-Pr)-4- piperidinyl	Me	4-[(2,6-dimethyl-4- pyridinyl)methoxy]phenyl	510
300	i-Bu	amino	4-(2- quinolinylmethoxy)phenyl	463
301	Me	amino	4-[(3,5-dimethylphenyl) methoxy]phenyl	398
302	Me	EtNHC(O)NH	4-[(3,5-dimethylphenyl) methoxy]phenyl	491
303	Me	CH ₃ SO ₂ NH	4-[(3,5-dimethylphenyl) methoxy]phenyl	498

304	Me	[(3-pyridinyl)acetyl]NH	4-[(3,5-dimethylphenyl)methoxy]phenyl	517
305	Me	4-pyridinyl-C(O)NH	4-[(3,5-dimethylphenyl)methoxy]phenyl	503
306	Me	amino	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	437
307	Me	4-pyridinyl-C(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	544
308	Me	EtNHC(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	532
309	Me	(CH ₃) ₃ CO-C(O)NHCH ₂ -C(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	618
310	Me	H ₂ NCH ₂ -C(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	496
311	Me	(3-pyridinyl)CH ₂ -C(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	558
312	Me	phenylCH ₂ NH-C(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	594
313	Me	[(2,4-dimethoxyphenyl)NHC(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	640
314	Me	phenyl-NHC(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	580
315	Me	(CH ₃) ₃ CO-C(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	561
316	Me	[2-(4-morpholinyl)ethyl]NHC(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	595
317	Me	(CH ₃) ₃ CO-C(O)NHCH ₂ -C(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	618
318	Me	(2-thiazolyl)NH-C(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	565
319	Me	(4-pyridinyl)NH-C(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	581
320	Me	(3-HO-phenyl)NH-C(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	596
321	Me	(2,3-dihydro-2-oxo-1H-benzimidazol-5-yl)NHC(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	636
322	CH ₃ SO ₂ CH ₂ CH ₂	amino	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	532
323	CH ₃ SO ₂ CH ₂ CH ₂	amino	4-[(3,5-dimethylphenyl)methoxy]phenyl	491
324	CH ₃ SO ₂ CH ₂ CH ₂	[(2-thiazolyl)-NHC(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	657

325	CH ₃ SO ₂ CH ₂ CH ₂	[(2-thiazolyl)NH C(O)NH	4-[(3,5-dimethylphenyl)methoxy]phenyl	617
326	4-[(2-propenyl)OC (O)NH]butyl	amino	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	580
327	4-[(2-propenyl)OC (O)NH]butyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	562
328	i-Bu	amino	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	481
329	i-Bu	[(2-thiazolyl)NH C(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	629
330	i-Bu	[(2-thiazolyl)NH C(O)NH	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	567
331	i-Bu	[(2-pyridinyl)NH C(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	623
332	i-Bu	CF ₃ CH ₂ C(O)- NHC(O)NH	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	537
333	i-Bu	[(2-pyridinyl)NH C(O)NH	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	561
334	i-Bu	phenylSO ₂ - NHC(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	686
335	i-Bu	phenylSO ₂ - NHC(O)NH	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	624
336	i-Bu	[(3-Me-5-isothiazolyl)NHC(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	621
337	i-Bu	1H-benzimidazo 1-2-ylNHC(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	640
338	i-Bu	1H-benzimidazo 1-2-ylNHC(O)NH	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	600
339	i-Bu	phenylNH- C(O)NH	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	560
340	i-Bu	phenyl- NHC(O)NH	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	622
341	i-Bu	(CH ₃) ₃ N ⁺	(phenylmethoxy)phenyl	454
342	i-Bu	amino	4-(4-quinolinylmethoxy)phenyl	446
343	i-Bu	amino	4-(2-oxo-2-phenylethoxy)phenyl	455
344	i-Bu	amino	4-[(3,5-dimethyl-4-isoxazolyl)methoxy]phenyl	431
345	i-Bu	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	441
346	i-Bu	amino	4-[2-(2-benzothiazolylamino)-2-oxoethoxy]phenyl	512
347	i-Bu	amino	4-[(2-methoxy-4-quinolinyl)methoxy]phenyl	476

348	i-Bu	amino	4-[(2-phenyl-4-quinolinyl)methoxy]phenyl	539
349	i-Bu	amino	4-[(2,6-dimethyl-4-quinolinyl)methoxy]phenyl	491
350	i-Bu	amino	4-[(2-chloro-4-quinolinyl)methoxy]phenyl	497
351	i-Bu	amino	4-[2-(2,5-dimethoxyphenyl)-2-(hydroxyimino)ethoxy]phenyl	515
352	i-Bu	amino	4-[(2-methylimidazo[1,2-a]pyridin-3-yl)methoxy]phenyl	466
353	i-Bu	amino	4-[[1,4-dimethyl-2-(methylthio)-1H-imidazol-5-yl]methoxy]phenyl	476
354	i-Bu	amino	4-[[1,5-dimethyl-2-(methylthio)-1H-imidazol-4-yl]methoxy]phenyl	476
355	i-Bu	amino	4-[(2,4-dimethyl-5-thiazolyl)methoxy]phenyl	447
356	i-Bu	amino	4-[(2-methyl-4-quinolinyl)methoxy]phenyl	477
357	CH ₃ SO ₂ CH ₂ CH ₂	amino	4-[(2-chloro-4-quinolinyl)methoxy]phenyl	547
358	CH ₃ SO ₂ CH ₂ CH ₂	amino	4-[(2-methyl-4-quinolinyl)methoxy]phenyl	527
359	CH ₃ SO ₂ CH ₂ CH ₂	amino	4-[(3,5-dimethoxyphenyl)methoxy]phenyl	522
360	CH ₃ SO ₂ CH ₂ CH ₂	amino	4-[(2-methoxy-4-quinolinyl)methoxy]phenyl	526
361	i-Bu	amino	4-[(3,5-dimethoxyphenyl)methoxy]phenyl	455
362	i-Bu	amino	4-[(2-CH ₃ O-5-nitrophenyl)methoxy]phenyl	470
363	i-Bu	amino	4-[(5-quinolinyl)methoxy]phenyl	446
364	2-(CH ₃ SO ₂)-ethyl	amino	4-[(2-CH ₃ O-5-nitrophenyl)methoxy]phenyl	520
365	2-(CH ₃ SO ₂)-ethyl	amino	4-[(2-nitro-4,5-dimethoxyphenyl)methoxy]phenyl	567
366	2-(CH ₃ SO ₂)-ethyl	amino	4-[(2-phenyl-4-quinolinyl)methoxy]phenyl	589
367	2-(CH ₃ SO ₂)-ethyl	amino	4-[(3,5-dimethyl-4-isoxazolyl)methoxy]phenyl	481
368	(4-HO-phenyl)-methyl	amino	4-[(phenyl)methoxy]phenyl	462
369	(4-CH ₃ O-phenyl)-methyl	amino	4-[(2-methyl-4-quinolinyl)methoxy]phenyl	541
370	(4-CH ₃ O-phenyl)-methyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	505
371	(4-CH ₃ O-phenyl)-methyl	amino	4-[(phenyl)methoxy]phenyl	476
450	i-Bu	aminomethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	455

451	i-Bu	2-thiazolylNH C(O)NHCH ₂	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	581
452	Me	aminomethyl	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	453
453	Me	2-thiazolylNH C(O)NHCH ₂	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	579
454			see structure at bottom	398
455	Me	HOCH ₂	4-[(3,5-dimethylphenyl)methoxy]phenyl	435
456	Me	CH ₃ CH ₂ NH- C(O)OCH ₂	4-[(3,5-dimethylphenyl)methoxy]phenyl	506
457	Me	HOCH ₂	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	476
458			see structure at bottom	381
459	Me	Me	5-[(3,5-dimethylphenoxy)methyl]-2-thienyl	425
460			see structure at bottom	460
461	Me	Me	[4-(phenylmethoxy)phenyl]methyl	405
462	i-Bu	CH ₃ NH	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	455
463	i-Bu	CH ₃ NH	4-[(2-methyl-4-quinolinyl)methoxy]phenyl	491
464			see structure at bottom	405
501	4-piperidinyl	amino	4-(4-quinolinylmethoxy)phenyl	490
502	4-piperidinyl	amino	4-[(2,6-chloro-4-pyridinyl)methoxy]phenyl	508
503	1-[(CH ₃) ₃ CO- C(O)]-4-piperidinyl	(CH ₃) ₃ CO- C(O)NH	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	668
504	4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	468
505	1-(CH ₃ SO ₂)- 4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	546
506	1-acetyl-4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	510
507	1-(2,2-dimethyl-1-oxopropyl)- 4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	552
508	1-[(CH ₃) ₃ CO- C(O)]-4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	568
509	1-(CH ₃ OC(O))- 4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	526
510	1-methyl-4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	482

511	1-dimethyl-carbamyl-4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	539
512	1-cycPr-C(O)-4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	536
513	i-Pr	amino	4-(4-quinolinylmethoxy)phenyl	449
514	i-Pr	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	427
515	cyclohexyl	amino	4-(4-quinolinylmethoxy)phenyl	589
516	cyclohexyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	467
517	t-Bu	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	441
518	t-Bu	amino	4-(4-quinolinylmethoxy)phenyl	461
519	t-Bu	amino	4-(2-methyl-4-quinolinylmethoxy)phenyl	477
520	i-Pr	amino	4-(2-methyl-4-quinolinylmethoxy)phenyl	463
521	i-Pr	amino	4-(2,6-dimethyl-4-quinolinylmethoxy)phenyl	477
522	1-(4-morpholino-C(O))-4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	581
523	1-(2-methyl-1-oxopropyl)-4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	538
524	4-CH ₃ O-cycHex	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	497
525			see structure at bottom	422
526	1-(phenyl-C(O))-4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	572
527	1-(1-oxopropyl)-4-piperidinyl	amino	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	524
528	1-acetyl-4-piperidinyl	amino	4-(2-methyl-4-quinolinylmethoxy)phenyl	546
529	1-(CH ₃ SO ₂)-4-piperidinyl	amino	4-(2-methyl-4-quinolinylmethoxy)phenyl	582
530	1-(2,2-di-CH ₃ -1-oxopropyl)-4-piperidinyl	amino	4-(2-methyl-4-quinolinylmethoxy)phenyl	588
531	1-acetyl-4-piperidinyl	amino	4-(4-quinolinylmethoxy)phenyl	532
532	1-(CH ₃ SO ₂)-4-piperidinyl	amino	4-(4-quinolinylmethoxy)phenyl	568


533	1-acetyl-4-piperidinyl	amino	4-[(3,5-dimethoxyphenyl)methoxy]phenyl	541
534	1-acetyl-4-piperidinyl	amino	4-[(5-methyl-3-nitrophenyl)methoxy]phenyl	540
535	1-acetyl-4-piperidinyl	amino	4-[3,5-bis(trifluoromethyl)phenoxy]phenyl	603
536	1-acetyl-4-piperidinyl	amino	4-[(3,5-dichlorophenyl)methoxy]phenyl	549
537	1-acetyl-4-piperidinyl	amino	4-(6-fluoro-2-methyl-4-quinolinylmethoxy)phenyl	564
538	1-acetyl-4-piperidinyl	amino	4-(7-chloro-2-methyl-4-quinolinylmethoxy)phenyl	580
539	1-acetyl-4-piperidinyl	amino	4-(6-chloro-2-methyl-4-quinolinylmethoxy)phenyl	580
540	1-acetyl-4-piperidinyl	amino	4-(6-methoxy-2-methyl-4-quinolinylmethoxy)phenyl	576
541	4-piperidinyl	amino	4-(2,7-dimethyl-4-quinolinylmethoxy)phenyl	518
542	1-acetyl-4-piperidinyl	amino	4-(2,7-dimethyl-4-quinolinylmethoxy)phenyl	560
543	4-piperidinyl	amino	4-(2-CH ₃ O-4-quinolinylmethoxy)phenyl	520
544	4-piperidinyl	amino	4-[(3,5-dimethoxyphenyl)methoxy]phenyl	499
545	4-piperidinyl	amino	4-[(2,6-diethyl-4-pyridinyl)methoxy]phenyl	496
546	1-acetyl-4-piperidinyl	amino	4-[(2,6-diethyl-4-pyridinyl)methoxy]phenyl	538
547	4-piperidinyl	amino	4-(7-methyl-4-quinolinylmethoxy)phenyl	504
548	4-methoxy-cycHex	amino	4-(4-quinolinylmethoxy)phenyl	519
549	t-Bu	amino	4-(2,6-dimethyl-4-quinolinylmethoxy)phenyl	491
550	methyl	methyl	4-[(2,6-dimethyl-1-oxido-4-pyridinyl)methoxy]phenyl	414
551	t-Bu	amino	4-(7-chloro-2-methyl-4-quinolinylmethoxy)phenyl	511
552	t-Bu	amino	4-(6-fluoro-2-methyl-4-quinolinylmethoxy)phenyl	495
553	t-Bu	amino	4-(6-chloro-2-methyl-4-quinolinylmethoxy)phenyl	511
554	t-Bu	amino	4-(6-methoxy-2-methyl-4-quinolinylmethoxy)phenyl	507
555	t-Bu	amino	4-(2,7-dimethyl-4-quinolinylmethoxy)phenyl	491
556	t-Bu	amino	4-(7-methyl-4-quinolinylmethoxy)phenyl	477
557	cycHex	amino	4-(2-methyl-4-quinolinylmethoxy)phenyl	503
558	cycHex	amino	4-(2,6-dimethyl-4-quinolinylmethoxy)phenyl	517
559	i-Pr	amino	4-[(5-methyl-3-nitrophenyl)methoxy]phenyl	457
560	i-Pr	amino	4-[3,5-bis(trifluoromethyl)phenoxy]phenyl	518
561	i-Pr	amino	4-[[3,5-bis(trifluoromethyl)phenyl]methoxy]phenyl	534

562	i-Pr	amino	4-(3,5-dibromophenoxy)phenyl	523
563	i-Pr	amino	4-(6-fluoro-2-methyl-4-quinolinylmethoxy)phenyl	481
564	i-Pr	amino	4-(6-CH ₃ O-2-methyl-4-quinolinylmethoxy)phenyl	493
565	i-Pr	amino	4-(7-chloro-2-methyl-4-quinolinylmethoxy)phenyl	497
566	i-Pr	amino	4-(6-chloro-2-methyl-4-quinolinylmethoxy)phenyl	497
567	i-Pr	amino	4-(2-CH ₃ O-4-quinolinylmethoxy)phenyl	479
568	i-Pr	amino	4-(2,7-dimethyl-4-quinolinylmethoxy)phenyl	477
569	i-Pr	amino	4-[(2,6-diethyl-4-pyridinyl)methoxy]phenyl	455
700	Me	Me	3-(phenylmethoxy)phenyl	367
701	Me	Me	3-[(3,5-dimethylphenyl)methoxy]phenyl	395
702	Me	Me	3-[(3-methylphenyl)methoxy]phenyl	381
703	Me	Me	3-(1-methylethoxy)phenyl	663
704	Me	Me	3-heptyloxyphenyl	375
705	Me	2-oxo-2-[(1,3,4-thiadiazol-2-yl)NH]ethyl	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	563
706	Me	2-((CH ₃) ₃ CO)-2-oxoethyl	4-(phenylmethoxy)phenyl	467
707	Me	2-HO-2-oxoethyl	4-(phenylmethoxy)phenyl	411
708	Me	2-[2-(CH ₃ NH)-2-oxoethyl]NH]-2-oxoethyl	4-[(3,5-dimethylphenyl)methoxy]phenyl	533
709	Me	2-oxo-2-[(2-thiazolyl)NH]ethyl	4-[(3,5-dimethylphenyl)methoxy]phenyl	521
710	Me	2-(4-morpholinyl)-2-oxoethyl	4-[(3,5-dimethylphenyl)methoxy]phenyl	532
711	Me	2-oxo-2-[(2-thiazolyl)NH]ethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	564
712	Me	2-[2-[(4-morpholinyl)ethyl]NH]-2-oxoethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	594
713	Me	2-oxo-2-[(4-pyridinyl)CH ₂ NH]ethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	594


714	Me	2-oxo-2-[(2-thiazolyl)NH]ethyl	4-[(3,5-dimethylphenyl)methoxy]phenyl	524
715	Me	2-oxo-2-[(3-pyridinyl)CH ₂ NH]ethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	594
716	Me	2-oxo-2-[(2-pyridinyl)CH ₂ NH]ethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	572
717	Me	2-oxo-2-[(4-pyridinyl)NH]ethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	558
718	Me	2-[(3-Me-5-isothiazolyl)NH]-2-oxoethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	576
719	Me	2-[[5-(t-Bu)-1,3,4-thiadiazol-2-yl]NH]-2-oxoethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	619
720	Me	2-[[4-(2-(t-Butoxyethoxy)-2-oxoethyl)-2-thiazolyl]NH]-2-oxoethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	676
721	Me	2-[[4-(2-HO-2-oxoethyl)-2-thiazolyl]NH]-2-oxoethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	620
722	Me	2-[[4-(2-CH ₃ NH-2-oxoethyl)-2-thiazolyl]NH]-2-oxoethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	657
723	Me	1H-benzimidazo1-2-ylmethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	554
724	Me	3H-imidazo[4,5-c]pyridin-2-ylmethyl	4-[(3,5-dichlorophenyl)methoxy]phenyl	555
725	Me	2-oxo-2-(2-thiazolyl)NH-ethyl	4-[3,5-bis(trifluoromethyl)phenoxy]phenyl	615
726	Me	2-oxo-2-[(4-pyridinyl)CH ₂ NH]ethyl	4-[3,5-bis(trifluoromethyl)phenoxy]phenyl	625

780	i-Pr	2-oxo-2-(4-pyridin-ylCH ₂)NH-ethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	560
781	i-Pr	2-oxo-2-(4-pyridin-ylCH ₃)NH-ethyl	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	600
782	cyclohexylmethyl	2-oxo-2-(4-pyridinylCH ₂)NH-ethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	614
783	cyclohexylmethyl	2-oxo-2-(4-pyridinylCH ₂)NH-ethyl	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	654
784	4-[(CH ₃) ₃ CO-C(O)NH]butyl	2-oxo-2-(4-pyridinylCH ₂)NH-ethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	689
785	4-aminobutyl	2-oxo-2-[(4-pyridinylCH ₃)NH-ethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	590
800	methyl	methyl	3-(1H-benzotriazol-1-ylmethoxy)phenyl	408
801	(3,4,4-tri-Me-2,5-dioxo-1-imidazol-1-yl)CH ₂	methyl	4-(phenylmethoxy)phenyl	509
802	i-Bu	2-(t-butoxy)-2-oxoethyl	4-(phenylmethoxy)phenyl	509
803	i-Bu	2-[2-(CH ₃ NH)-2-oxoethyl]NH-2-oxoethyl	4-[(3,5-dimethylphenyl)methoxy]phenyl	533
804	i-Bu	2-[2-(CH ₃ NH)-2-oxoethyl]NH-2-oxoethyl	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	595
805	i-Bu	2-oxo-2-(2-thiazol-yl)NH-ethyl	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	607
806	i-Bu	2-[2-(CH ₃ NH)-2-oxoethyl]NH-2-oxoethyl	4-[3,5-bis(trifluoromethyl)phenoxy]phenyl	647
807	i-Bu	2-oxo-2-[(4-pyridinyl)CH ₂]NH-ethyl	4-[3,5-bis(trifluoromethyl)phenoxy]phenyl	667


808	i-Bu	2-oxo-2-(phenyl-NH)ethyl	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	600
809	i-Bu	2-oxo-2-(CH ₃ -NH)ethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	497
810	i-Bu	2-[2-(1H-imidazol-4-yl)ethyl]NH-2-oxoethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	577
811	i-Bu	2-2-[1-(phenylCH ₂)-4-piperidinyl NH]-2-oxoethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	656
812	i-Bu	2-[2-(dimethylamino)ethyl]NH-2-oxoethyl	4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl	554
813	i-Bu	2-[(4-HO-phenyl)NH]-2-oxoethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	575
814	i-Bu	2-oxo-2-(2-thiazol-yl)NH-ethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	566
815	i-Bu	2-HO-ethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	470
816	i-Bu	2-[(4,5-dimethyl-2-thiazol-yl)NH]-2-oxoethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	594
817	i-Bu	2-[(1H-indazol-5-yl)NH]-2-oxoethyl	4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl	599
818	i-Bu	2-oxo-2-[(2-thiazol-yl)NH]ethyl	4-[3,5-bis(trifluoromethyl)phenoxy]phenyl	659




(example 30)




(example 55)




(example 107)




(example 113)




(example 454)




(example 458)



(example 460)








(example 464)




(example 525)


The following tables contain representative examples of the present invention. Each entry in each table is intended to be paired with each formula at the start of the table. For example, in Table 2, example 1 is intended to be paired with each of formulae A1-FF3.

TABLE 2







U1 ($R^2=Me$)
U2 ($R^2=NH_2$)
U3 ($R^2=OH$)




V1 ($R^2=Me$)
V2 ($R^2=NH_2$)
V3 ($R^2=OH$)




W1 ($R^2=Me$)
W2 ($R^2=NH_2$)
W3 ($R^2=OH$)




X1 ($R^2=Me$)
X2 ($R^2=NH_2$)
X3 ($R^2=OH$)




Y1 ($R^2=Me$)
Y2 ($R^2=NH_2$)
Y3 ($R^2=OH$)




Z1 ($R^2=Me$)
Z2 ($R^2=NH_2$)
Z3 ($R^2=OH$)




AA1 ($R^2=Me$)
AA2 ($R^2=NH_2$)
AA3 ($R^2=OH$)




BB1 ($R^2=Me$)
BB2 ($R^2=NH_2$)
BB3 ($R^2=OH$)




CC1 ($R^2=Me$)
CC2 ($R^2=NH_2$)
CC3 ($R^2=OH$)



DD1 ($R^2=Me$)
DD2 ($R^2=NH_2$)
DD3 ($R^2=OH$)



EE1 ($R^2=Me$)
EE2 ($R^2=NH_2$)
EE3 ($R^2=OH$)



FF1 ($R^2=Me$)
FF2 ($R^2=NH_2$)
FF3 ($R^2=OH$)


5

Ex#	R^{10}
1	H
2	methyl
3	methoxy
4	1-methylethyl
5	1-methylethoxy
6	phenyl
7	[1,1'-biphenyl]-4-yl
8	phenoxy
9	2-phenylethyl
10	2-(3,5-dimethylphenyl)ethyl
11	1-(2,6-dimethylphenyl)ethyl
12	2-phenylethenyl
13	phenoxyethyl
14	(2-methylphenyl)methoxy
15	(3-methylphenyl)methoxy
16	3-methylphenoxy
17	2,6-dimethylphenoxy
18	(2,6-dimethylphenyl)methoxy
19	3,5-dimethylphenoxy


20	(3,5-dimethylphenyl)methoxy
21	2-(3,5-dimethylphenyl)ethyl
22	2-(3,5-dimethylphenyl)ethenyl
23	(3-amino-5-methylphenyl)methoxy
24	(2-amino-6-methylphenyl)methoxy
25	(3-cyano-5-methylphenyl)methoxy
26	(3-cyano-5-methylphenoxy)methyl
27	(3-cyano-5-nitrophenyl)methoxy
28	(3,5-diethoxyphenyl)methoxy
29	(3,5-dimethoxyphenyl)methoxy
30	3,5-dimethoxyphenoxy
31	2-(3,5-dimethoxyphenyl)ethyl
32	1-(3,5-dimethoxyphenyl)ethoxy
33	(3,5-dichlorophenyl)methoxy
34	(2,6-dichlorophenyl)methoxy
35	(3,5-dibromophenyl)methoxy
36	3,5-dibromophenoxy
37	(3-amino-5-cyanophenyl)methoxy
38	[2,6-bis(trifluoromethyl)phenyl]methoxy
39	2,6-bis(trifluoromethyl)phenoxy
40	(3-aminocarbonyl-5-methylphenyl)methoxy
41	([1,1'-biphenyl]-2-yl)methoxy
42	([1,1'-biphenyl]-3-yl)methoxy
43	[5-methyl-3-(methylsulfonyl)phenyl]methoxy
44	5-methyl-3-(methylsulfonyl)phenoxy
45	(2-pyridinyl)methoxy
46	(4-pyridinyl)methoxy
47	(2,6-dimethyl-4-pyridinyl)methoxy
48	2,6-dimethyl-4-pyridinyloxy
49	1-(2,6-dimethyl-4-pyridinyl)ethoxy
50	(3,5-dimethyl-4-pyridinyl)methoxy
51	(2,6-diethyl-4-pyridinyl)methoxy
52	(2,6-dichloro-4-pyridinyl)methoxy
53	(2,6-dimethoxy-4-pyridinyl)methoxy
54	(2-chloro-6-methyl-4-pyridinyl)methoxy
55	(2-chloro-6-methoxy-4-pyridinyl)methoxy
56	(2-methoxy-6-methyl-4-pyridinyl)methoxy
57	(1-naphthalenyl)methoxy
58	1-naphthalenyloxy
59	(2-naphthalenyl)methoxy
60	(2-methyl-1-naphthalenyl)methoxy
61	(4-methyl-2-naphthalenyl)methoxy
62	(4-quinolinyl)methoxy
63	1-(4-quinolinyl)ethoxy
64	4-quinolinylloxy
65	(4-quinolinylloxy)methyl
66	2-(4-quinolinyl)ethyl
67	(2-methyl-4-quinolinyl)methoxy
68	2-methyl-4-quinolinylloxy
69	(2-chloro-4-quinolinyl)methoxy
70	(2-methoxy-4-quinolinyl)methoxy
71	(2-hydroxy-4-quinolinyl)methoxy

72	(2-trifluoromethyl-4-quinolinyl)methoxy
73	(2-phenyl-4-quinolinyl)methoxy
74	(2,6-dimethyl-4-quinolinyl)methoxy
75	(2,7-dimethyl-4-quinolinyl)methoxy
76	(5-quinolinyl)methoxy
77	(7-methyl-5-quinolinyl)methoxy
78	(7-methoxy-5-quinolinyl)methoxy
79	(8-quinolinyl)methoxy
80	2-(1,2,3-benzotriazol-1-yl)ethyl
81	(2-benzimidazolyl)methoxy
82	(1,4-dimethyl-5-imidazolyl)methoxy
83	(3,5-dimethyl-4-isoxazolyl)methoxy
84	(4,5-dimethyl-2-oxazolyl)methoxy
85	(2,5-dimethyl-4-thiazolyl)methoxy
86	(3,5-dimethyl-1-pyrazolyl)ethyl
87	(1,3-benzodioxo-4-yl)methoxy
88	(1,3,5-trimethyl-4-pyrazolyl)methoxy
89	(2,6-dimethyl-4-pyrimidinyl)methoxy
90	(4,5-dimethyl-2-furanyl)methoxy
91	(4,5-dimethyl-2-thiazolyl)methoxy
92	2-(2-oxazolyl)ethyl


TABLE 3




A1 (X=linker Σ)
 A2 (X=linker Δ)
 A3 (X=linker Φ)
 A4 (X=linker Ω)
 A5 (X=linker Π)
 A6 (X=linker Ψ)
 A7 (X=linker Λ)




B1 (X=linker Σ)
 B2 (X=linker Δ)
 B3 (X=linker Φ)
 B4 (X=linker Ω)
 B5 (X=linker Π)
 B6 (X=linker Ψ)
 B7 (X=linker Λ)




C1 (X=linker Σ)
 C2 (X=linker Δ)
 C3 (X=linker Φ)
 C4 (X=linker Ω)
 C5 (X=linker Π)
 C6 (X=linker Ψ)
 C7 (X=linker Λ)




D1 (X=linker Σ)
 D2 (X=linker Δ)
 D3 (X=linker Φ)
 D4 (X=linker Ω)
 D5 (X=linker Π)
 D6 (X=linker Ψ)
 D7 (X=linker Λ)




E1 (X=linker Σ)
 E2 (X=linker Δ)
 E3 (X=linker Φ)
 E4 (X=linker Ω)
 E5 (X=linker Π)
 E6 (X=linker Ψ)
 E7 (X=linker Λ)




F1 (X=linker Σ)
 F2 (X=linker Δ)
 F3 (X=linker Φ)
 F4 (X=linker Ω)
 F5 (X=linker Π)
 F6 (X=linker Ψ)
 F7 (X=linker Λ)




G1 (X=linker Σ)
 G2 (X=linker Δ)
 G3 (X=linker Φ)
 G4 (X=linker Ω)
 G5 (X=linker Π)
 G6 (X=linker Ψ)
 G7 (X=linker Λ)




H1 (X=linker Σ)
 H2 (X=linker Δ)
 H3 (X=linker Φ)
 H4 (X=linker Ω)
 H5 (X=linker Π)
 H6 (X=linker Ψ)
 H7 (X=linker Λ)




I1 (X=linker Σ)
 I2 (X=linker Δ)
 I3 (X=linker Φ)
 I4 (X=linker Ω)
 I5 (X=linker Π)
 I6 (X=linker Ψ)
 I7 (X=linker Λ)




J1 (X=linker Σ)
 J2 (X=linker Δ)
 J3 (X=linker Φ)
 J4 (X=linker Ω)
 J5 (X=linker Π)
 J6 (X=linker Ψ)
 J7 (X=linker Λ)



K1 (X=linker Σ)
 K2 (X=linker Δ)
 K3 (X=linker Φ)
 K4 (X=linker Ω)
 K5 (X=linker Π)
 K6 (X=linker Ψ)
 K7 (X=linker Λ)



L1 (X=linker Σ)
 L2 (X=linker Δ)
 L3 (X=linker Φ)
 L4 (X=linker Ω)
 L5 (X=linker Π)
 L6 (X=linker Ψ)
 L7 (X=linker Λ)







Ex #	R ²	R ¹⁰
1	amino	methoxy
2	amino	1-methylethyl
3	amino	1-methylethoxy
4	amino	phenyl
5	amino	phenoxy
6	amino	2-phenylethyl
7	amino	2-(3,5-dimethylphenyl)ethyl
8	amino	2-phenylethenyl
9	amino	phenoxymethyl
10	amino	3,5-dimethylphenoxy
11	amino	(3,5-dimethylphenyl)methoxy
12	amino	2-(3,5-dimethylphenyl)ethyl
13	amino	2-(3,5-dimethylphenyl)ethenyl
14	amino	(3-amino-5-methylphenyl)methoxy
15	amino	(3,5-dimethoxyphenyl)methoxy
16	amino	3,5-dimethoxyphenoxy
17	amino	2-(3,5-dimethoxyphenyl)ethyl
18	amino	(3,5-dichlorophenyl)methoxy
19	amino	3,5-dibromophenoxy
20	amino	[2,6-bis(trifluoromethyl)phenyl]methoxy
21	amino	2,6-bis(trifluoromethyl)phenoxy
22	amino	[5-methyl-3-(methylsulfonyl)phenyl]methoxy
23	amino	5-methyl-3-(methylsulfonyl)phenoxy
24	amino	(2,6-dimethyl-4-pyridinyl)methoxy
25	amino	2,6-dimethyl-4-pyridinyloxy
26	amino	(2,6-dichloro-4-pyridinyl)methoxy
27	amino	(2-methoxy-6-methyl-4-pyridinyl)methoxy
28	amino	(1-naphthalenyl)methoxy
29	amino	1-naphthalenyloxy
30	amino	(2-naphthalenyl)methoxy
31	amino	(2-methyl-1-naphthalenyl)methoxy
32	amino	(4-methyl-2-naphthalenyl)methoxy
33	amino	(4-quinolinyl)methoxy
34	amino	1-(4-quinolinyl)ethoxy
35	amino	4-quinolinylloxy
36	amino	(4-quinolinylloxy)methyl
37	amino	(2-methyl-4-quinolinyl)methoxy

38	amino	2-methyl-4-quinolinylloxy
39	amino	(2-methoxy-4-quinolinyl)methoxy
40	amino	2-(1,2,3-benzotriazol-1-yl)ethyl
41	amino	(2-benzimidazolyl)methoxy
42	amino	(1,4-dimethyl-5-imidazolyl)methoxy
43	amino	(3,5-dimethyl-4-isoxazolyl)methoxy
44	amino	(4,5-dimethyl-2-oxazolyl)methoxy
45	amino	(2,5-dimethyl-4-thiazolyl)methoxy
46	amino	(3,5-dimethyl-1-pyrazolyl)ethyl
47	amino	(1,3-benzodioxo-4-yl)methoxy
48	amino	(1,3,5-trimethyl-4-pyrazolyl)methoxy
49	amino	(2,6-dimethyl-4-pyrimidinyl)methoxy
50	amino	(4,5-dimethyl-2-furanyl)methoxy
51	amino	(4,5-dimethyl-2-thiazolyl)methoxy
52	amino	2-(2-oxazolyl)ethyl
53	methyl	methoxy
54	methyl	1-methylethyl
55	methyl	1-methylethoxy
56	methyl	phenyl
57	methyl	phenoxy
58	methyl	2-phenylethyl
59	methyl	2-(3,5-dimethylphenyl)ethyl
60	methyl	2-phenylethenyl
61	methyl	phenoxymethyl
62	methyl	3,5-dimethylphenoxy
63	methyl	(3,5-dimethylphenyl)methoxy
64	methyl	2-(3,5-dimethylphenyl)ethyl
65	methyl	2-(3,5-dimethylphenyl)ethenyl
66	methyl	(3-amino-5-methylphenyl)methoxy
67	methyl	(3,5-dimethoxyphenyl)methoxy
68	methyl	3,5-dimethoxyphenoxy
69	methyl	2-(3,5-dimethoxyphenyl)ethyl
70	methyl	(3,5-dichlorophenyl)methoxy
71	methyl	3,5-dibromophenoxy
72	methyl	[2,6-bis(trifluoromethyl)phenyl)methoxy
73	methyl	2,6-bis(trifluoromethyl)phenoxy
74	methyl	[5-methyl-3-(methylsulfonyl)phenyl)methoxy
75	methyl	5-methyl-3-(methylsulfonyl)phenoxy
76	methyl	(2,6-dimethyl-4-pyridinyl)methoxy
77	methyl	2,6-dimethyl-4-pyridinyloxy
78	methyl	(2,6-dichloro-4-pyridinyl)methoxy
79	methyl	(2-methoxy-6-methyl-4-pyridinyl)methoxy
80	methyl	(1-naphthalenyl)methoxy
81	methyl	1-naphthalenyloxy
82	methyl	(2-naphthalenyl)methoxy
83	methyl	(2-methyl-1-naphthalenyl)methoxy
84	methyl	(4-methyl-2-naphthalenyl)methoxy
85	methyl	(4-quinolinyl)methoxy
86	methyl	1-(4-quinolinyl)ethoxy
87	methyl	4-quinolinylloxy
88	methyl	(4-quinolinylloxy)methyl
89	methyl	(2-methyl-4-quinolinyl)methoxy
90	methyl	2-methyl-4-quinolinylloxy
91	methyl	(2-methoxy-4-quinolinyl)methoxy
92	methyl	2-(1,2,3-benzotriazol-1-yl)ethyl
93	methyl	(2-benzimidazolyl)methoxy

94	methyl	(1,4-dimethyl-5-imidazolyl)methoxy
95	methyl	(3,5-dimethyl-4-isoxazolyl)methoxy
96	methyl	(4,5-dimethyl-2-oxazolyl)methoxy
97	methyl	(2,5-dimethyl-4-thiazolyl)methoxy
98	methyl	(3,5-dimethyl-1-pyrazolyl)ethyl
99	methyl	(1,3-benzodioxo-4-yl)methoxy
100	methyl	(1,3,5-trimethyl-4-pyrazolyl)methoxy
101	methyl	(2,6-dimethyl-4-pyrimidinyl)methoxy
102	methyl	(4,5-dimethyl-2-furanyl)methoxy
103	methyl	(4,5-dimethyl-2-thiazolyl)methoxy
104	methyl	2-(2-oxazolyl)ethyl

TABLE 4






















Ex #	R ²	R ¹⁰
1	amino	methoxy
2	amino	1-methylethyl
3	amino	1-methylethoxy
4	amino	phenyl
5	amino	phenoxy
6	amino	2-phenylethyl
7	amino	2-(3,5-dimethylphenyl)ethyl
8	amino	2-phenylethenyl
9	amino	phenoxymethyl
10	amino	3,5-dimethylphenoxy
11	amino	(3,5-dimethylphenyl)methoxy
12	amino	2-(3,5-dimethylphenyl)ethyl
13	amino	2-(3,5-dimethylphenyl)ethenyl
14	amino	(3-amino-5-methylphenyl)methoxy
15	amino	(3,5-dimethoxyphenyl)methoxy

16	amino	3,5-dimethoxyphenoxy
17	amino	2-(3,5-dimethoxyphenyl)ethyl
18	amino	(3,5-dichlorophenyl)methoxy
19	amino	3,5-dibromophenoxy
20	amino	[2,6-bis(trifluoromethyl)phenyl]methoxy
21	amino	2,6-bis(trifluoromethyl)phenoxy
22	amino	[5-methyl-3-(methylsulfonyl)phenyl]methoxy
23	amino	5-methyl-3-(methylsulfonyl)phenoxy
24	amino	(2,6-dimethyl-4-pyridinyl)methoxy
25	amino	2,6-dimethyl-4-pyridinyloxy
26	amino	(2,6-dichloro-4-pyridinyl)methoxy
27	amino	(2-methoxy-6-methyl-4-pyridinyl)methoxy
28	amino	(1-naphthalenyl)methoxy
29	amino	1-naphthalenyloxy
30	amino	(2-naphthalenyl)methoxy
31	amino	(2-methyl-1-naphthalenyl)methoxy
32	amino	(4-methyl-2-naphthalenyl)methoxy
33	amino	(4-quinolinyl)methoxy
34	amino	1-(4-quinolinyl)ethoxy
35	amino	4-quinolinylloxy
36	amino	(4-quinolinylloxy)methyl
37	amino	(2-methyl-4-quinolinyl)methoxy
38	amino	2-methyl-4-quinolinylloxy
39	amino	(2-methoxy-4-quinolinyl)methoxy
40	amino	2-(1,2,3-benzotriazol-1-yl)ethyl
41	amino	(2-benzimidazolyl)methoxy
42	amino	(1,4-dimethyl-5-imidazolyl)methoxy
43	amino	(3,5-dimethyl-4-isoxazolyl)methoxy
44	amino	(4,5-dimethyl-2-oxazolyl)methoxy
45	amino	(2,5-dimethyl-4-thiazolyl)methoxy
46	amino	(3,5-dimethyl-1-pyrazolyl)ethyl
47	amino	(1,3-benzodioxo-4-yl)methoxy
48	amino	(1,3,5-trimethyl-4-pyrazolyl)methoxy
49	amino	(2,6-dimethyl-4-pyrimidinyl)methoxy
50	amino	(4,5-dimethyl-2-furanyl)methoxy
51	amino	(4,5-dimethyl-2-thiazolyl)methoxy
52	amino	2-(2-oxazolyl)ethyl
53	methyl	methoxy
54	methyl	1-methylethyl
55	methyl	1-methylethoxy
56	methyl	phenyl
57	methyl	phenoxy
58	methyl	2-phenylethyl
59	methyl	2-(3,5-dimethylphenyl)ethyl
60	methyl	2-phenylethenyl
61	methyl	phenoxymethyl
62	methyl	3,5-dimethylphenoxy
63	methyl	(3,5-dimethylphenyl)methoxy
64	methyl	2-(3,5-dimethylphenyl)ethyl
65	methyl	2-(3,5-dimethylphenyl)ethenyl
66	methyl	(3-amino-5-methylphenyl)methoxy
67	methyl	(3,5-dimethoxyphenyl)methoxy
68	methyl	3,5-dimethoxyphenoxy
69	methyl	2-(3,5-dimethoxyphenyl)ethyl
70	methyl	(3,5-dichlorophenyl)methoxy
71	methyl	3,5-dibromophenoxy

72	methyl	[2,6-bis(trifluoromethyl)phenyl]methoxy
73	methyl	2,6-bis(trifluoromethyl)phenoxy
74	methyl	[5-methyl-3-(methylsulfonyl)phenyl]methoxy
75	methyl	5-methyl-3-(methylsulfonyl)phenoxy
76	methyl	(2,6-dimethyl-4-pyridinyl)methoxy
77	methyl	2,6-dimethyl-4-pyridinyloxy
78	methyl	(2,6-dichloro-4-pyridinyl)methoxy
79	methyl	(2-methoxy-6-methyl-4-pyridinyl)methoxy
80	methyl	(1-naphthalenyl)methoxy
81	methyl	1-naphthalenyloxy
82	methyl	(2-naphthalenyl)methoxy
83	methyl	(2-methyl-1-naphthalenyl)methoxy
84	methyl	(4-methyl-2-naphthalenyl)methoxy
85	methyl	(4-quinolinyl)methoxy
86	methyl	1-(4-quinolinyl)ethoxy
87	methyl	4-quinolinylloxy
88	methyl	(4-quinolinylloxy)methyl
89	methyl	(2-methyl-4-quinolinyl)methoxy
90	methyl	2-methyl-4-quinolinylloxy
91	methyl	(2-methoxy-4-quinolinyl)methoxy
92	methyl	2-(1,2,3-benzotriazol-1-yl)ethyl
93	methyl	(2-benzimidazolyl)methoxy
94	methyl	(1,4-dimethyl-5-imidazolyl)methoxy
95	methyl	(3,5-dimethyl-4-isoxazolyl)methoxy
96	methyl	(4,5-dimethyl-2-oxazolyl)methoxy
97	methyl	(2,5-dimethyl-4-thiazolyl)methoxy
98	methyl	(3,5-dimethyl-1-pyrazolyl)ethyl
99	methyl	(1,3-benzodioxo-4-yl)methoxy
100	methyl	(1,3,5-trimethyl-4-pyrazolyl)methoxy
101	methyl	(2,6-dimethyl-4-pyrimidinyl)methoxy
102	methyl	(4,5-dimethyl-2-furanyl)methoxy
103	methyl	(4,5-dimethyl-2-thiazolyl)methoxy
104	methyl	2-(2-oxazolyl)ethyl

TABLE 5





Ex #	R ¹	R ²
1	Me	H
2	OH	H
3	NH ₂	H

4	Me	methyl
5	OH	methyl
6	NH ₂	methyl
7	Me	ethyl
8	OH	ethyl
9	NH ₂	ethyl
10	Me	isopropyl
11	OH	isopropyl
12	NH ₂	isopropyl
13	Me	phenyl
14	OH	phenyl
15	NH ₂	phenyl
16	Me	benzyl
17	OH	benzyl
18	NH ₂	benzyl
19	Me	2-phenylethyl
20	OH	2-phenylethyl
21	NH ₂	2-phenylethyl
22	Me	2-(2-methylphenyl)ethyl
23	OH	2-(2-methylphenyl)ethyl
24	NH ₂	2-(2-methylphenyl)ethyl
25	Me	2-(3-methylphenyl)ethyl
26	OH	2-(3-methylphenyl)ethyl
27	NH ₂	2-(3-methylphenyl)ethyl
28	Me	2-(2,6-dimethylphenyl)ethyl
29	OH	2-(2,6-dimethylphenyl)ethyl
30	NH ₂	2-(2,6-dimethylphenyl)ethyl
31	Me	2-(3,5-dimethylphenyl)ethyl
32	OH	2-(3,5-dimethylphenyl)ethyl
33	NH ₂	2-(3,5-dimethylphenyl)ethyl
34	Me	2-(3-amino-5-methylphenyl)ethyl
35	OH	2-(3-amino-5-methylphenyl)ethyl
36	NH ₂	2-(3-amino-5-methylphenyl)ethyl
37	Me	2-(pyridin-4-yl)ethyl
38	OH	2-(pyridin-4-yl)ethyl
39	NH ₂	2-(pyridin-4-yl)ethyl
40	Me	2-(2,6-dimethylpyridin-4-yl)ethyl
41	OH	2-(2,6-dimethylpyridin-4-yl)ethyl
42	NH ₂	2-(2,6-dimethylpyridin-4-yl)ethyl
43	Me	2-(3,5-dimethylpyridin-4-yl)ethyl
44	OH	2-(3,5-dimethylpyridin-4-yl)ethyl
45	NH ₂	2-(3,5-dimethylpyridin-4-yl)ethyl
46	Me	styryl
47	OH	styryl
48	NH ₂	styryl
49	Me	hydroxy
50	OH	hydroxy
51	NH ₂	hydroxy
52	Me	methoxy
53	OH	methoxy
54	NH ₂	methoxy
55	Me	ethoxy
56	OH	ethoxy
57	NH ₂	ethoxy
58	Me	isopropoxy
59	OH	isopropoxy
60	NH ₂	isopropoxy
61	Me	tert-butoxy
62	OH	tert-butoxy
63	NH ₂	tert-butoxy


64	Me	cyclohexyloxy
65	OH	cyclohexyloxy
66	NH ₂	cyclohexyloxy
67	Me	phenoxy
68	OH	phenoxy
69	NH ₂	phenoxy
70	Me	o-methylphenoxy
71	OH	o-methylphenoxy
72	NH ₂	o-methylphenoxy
73	Me	m-methylphenoxy
74	OH	m-methylphenoxy
75	NH ₂	m-methylphenoxy
76	Me	cinnamyloxy
77	OH	cinnamyloxy
78	NH ₂	cinnamyloxy
79	Me	benzyloxy
80	OH	benzyloxy
81	NH ₂	benzyloxy
82	Me	phenoxymethyl
83	OH	phenoxymethyl
84	NH ₂	phenoxymethyl
85	Me	o-methylbenzyloxy
86	OH	o-methylbenzyloxy
87	NH ₂	o-methylbenzyloxy
88	Me	m-methylbenzyloxy
89	OH	m-methylbenzyloxy
90	NH ₂	m-methylbenzyloxy
91	Me	o,o-dimethylbenzyloxy
92	OH	o,o-dimethylbenzyloxy
93	NH ₂	o,o-dimethylbenzyloxy
94	Me	(2,6-dimethylphenoxy)methyl
95	OH	(2,6-dimethylphenoxy)methyl
96	NH ₂	(2,6-dimethylphenoxy)methyl
97	Me	m,m-dimethylbenzyloxy
98	OH	m,m-dimethylbenzyloxy
99	NH ₂	m,m-dimethylbenzyloxy
100	Me	(3,5-dimethylphenoxy)methyl
101	OH	(3,5-dimethylphenoxy)methyl
102	NH ₂	(3,5-dimethylphenoxy)methyl
103	Me	o,o-dicyanobenzyloxy
104	OH	o,o-dicyanobenzyloxy
105	NH ₂	o,o-dicyanobenzyloxy
106	Me	m,m-dicyanobenzyloxy
107	OH	m,m-dicyanobenzyloxy
108	NH ₂	m,m-dicyanobenzyloxy
109	Me	(2,6-dicyanophenoxy)methyl
110	OH	(2,6-dicyanophenoxy)methyl
111	NH ₂	(2,6-dicyanophenoxy)methyl
112	Me	(3,5-dicyanophenoxy)methyl
113	OH	(3,5-dicyanophenoxy)methyl
114	NH ₂	(3,5-dicyanophenoxy)methyl
115	Me	o-amino-o-cyanobenzyloxy
116	OH	o-amino-o-cyanobenzyloxy
117	NH ₂	o-amino-o-cyanobenzyloxy
118	Me	m-amino-m-cyanobenzyloxy
119	OH	m-amino-m-cyanobenzyloxy
120	NH ₂	m-amino-m-cyanobenzyloxy
121	Me	o-amino-o-nitrobenzyloxy
122	OH	o-amino-o-nitrobenzyloxy
123	NH ₂	o-amino-o-nitrobenzyloxy













124	Me	m-amino-m-nitrobenzyloxy
125	OH	m-amino-m-nitrobenzyloxy
126	NH ₂	m-amino-m-nitrobenzyloxy
127	Me	p-amino-m,m-dimethylbenzyloxy
128	OH	p-amino-m,m-dimethylbenzyloxy
129	NH ₂	p-amino-m,m-dimethylbenzyloxy
130	Me	o-amino-o-methylbenzyloxy
131	OH	o-amino-o-methylbenzyloxy
132	NH ₂	o-amino-o-methylbenzyloxy
133	Me	m-amino-m-methylbenzyloxy
134	OH	m-amino-m-methylbenzyloxy
135	NH ₂	m-amino-m-methylbenzyloxy
136	Me	o-cyano-o-methylbenzyloxy
137	OH	o-cyano-o-methylbenzyloxy
138	NH ₂	o-cyano-o-methylbenzyloxy
139	Me	m-cyano-m-methylbenzyloxy
140	OH	m-cyano-m-methylbenzyloxy
141	NH ₂	m-cyano-m-methylbenzyloxy
142	Me	o-cyano-o-nitrobenzyloxy
143	OH	o-cyano-o-nitrobenzyloxy
144	NH ₂	o-cyano-o-nitrobenzyloxy
145	Me	(2-cyano-6-nitrophenoxy)methyl
146	OH	(2-cyano-6-nitrophenoxy)methyl
147	NH ₂	(2-cyano-6-nitrophenoxy)methyl
148	Me	m-cyano-m-nitrobenzyloxy
149	OH	m-cyano-m-nitrobenzyloxy
150	NH ₂	m-cyano-m-nitrobenzyloxy
151	Me	(3-cyano-5-nitrophenoxy)methyl
152	OH	(3-cyano-5-nitrophenoxy)methyl
153	NH ₂	(3-cyano-5-nitrophenoxy)methyl
154	Me	m,m-dimethoxybenzyloxy
155	OH	m,m-dimethoxybenzyloxy
156	NH ₂	m,m-dimethoxybenzyloxy
157	Me	m,m-dichlorobenzyloxy
158	OH	m,m-dichlorobenzyloxy
159	NH ₂	m,m-dichlorobenzyloxy
160	Me	(3,5-dichlorophenoxy)methyl
161	OH	(3,5-dichlorophenoxy)methyl
162	NH ₂	(3,5-dichlorophenoxy)methyl
163	Me	m,m-dibromobenzyloxy
164	OH	m,m-dibromobenzyloxy
165	NH ₂	m,m-dibromobenzyloxy
166	Me	m,m-bis(trifluoromethyl)benzyloxy
167	OH	m,m-bis(trifluoromethyl)benzyloxy
168	NH ₂	m,m-bis(trifluoromethyl)benzyloxy
169	Me	[3,5-bis(trifluoromethyl)phenoxy)methyl
170	OH	[3,5-bis(trifluoromethyl)phenoxy)methyl
171	NH ₂	[3,5-bis(trifluoromethyl)phenoxy)methyl
172	Me	m-carboxamido-m-methylbenzyloxy
173	OH	m-carboxamido-m-methylbenzyloxy
174	NH ₂	m-carboxamido-m-methylbenzyloxy
175	Me	(3-carboxamido-5-methylphenoxy)methyl
176	OH	(3-carboxamido-5-methylphenoxy)methyl
177	NH ₂	(3-carboxamido-5-methylphenoxy)methyl
178	Me	m-hydroxycarbonyl-m-methylbenzyloxy
179	OH	m-hydroxycarbonyl-m-methylbenzyloxy
180	NH ₂	m-hydroxycarbonyl-m-methylbenzyloxy
181	Me	(3-hydroxycarbonyl-5-methylphenoxy)methyl
182	OH	(3-hydroxycarbonyl-5-methylphenoxy)methyl
183	NH ₂	(3-hydroxycarbonyl-5-methylphenoxy)methyl

184	Me	o-phenylbenzyloxy
185	OH	o-phenylbenzyloxy
186	NH ₂	o-phenylbenzyloxy
187	Me	m-phenylbenzyloxy
188	OH	m-phenylbenzyloxy
189	NH ₂	m-phenylbenzyloxy
190	Me	(naphth-1-yl)methoxy
191	OH	(naphth-1-yl)methoxy
192	NH ₂	(naphth-1-yl)methoxy
193	Me	(naphth-2-yl)methoxy
194	OH	(naphth-2-yl)methoxy
195	NH ₂	(naphth-2-yl)methoxy
196	Me	(2-methylnaphth-1-yl)methoxy
197	OH	(2-methylnaphth-1-yl)methoxy
198	NH ₂	(2-methylnaphth-1-yl)methoxy
199	Me	(4-methylnaphth-2-yl)methoxy
200	OH	(4-methylnaphth-2-yl)methoxy
201	NH ₂	(4-methylnaphth-2-yl)methoxy
202	Me	(pyridin-3-yl)methoxy
203	OH	(pyridin-3-yl)methoxy
204	NH ₂	(pyridin-3-yl)methoxy
205	Me	(pyridin-4-yl)methoxy
206	OH	(pyridin-4-yl)methoxy
207	NH ₂	(pyridin-4-yl)methoxy
208	Me	(3,5-dichloropyridin-4-yl)methoxy
209	OH	(3,5-dichloropyridin-4-yl)methoxy
210	NH ₂	(3,5-dichloropyridin-4-yl)methoxy
211	Me	(3,5-dimethylpyridin-4-yl)methoxy
212	OH	(3,5-dimethylpyridin-4-yl)methoxy
213	NH ₂	(3,5-dimethylpyridin-4-yl)methoxy
214	Me	(1,2,3-benzotriazol-1-yl)methoxy
215	OH	(1,2,3-benzotriazol-1-yl)methoxy
216	NH ₂	(1,2,3-benzotriazol-1-yl)methoxy
217	Me	benzhydroxy
218	OH	benzhydroxy
219	NH ₂	benzhydroxy
220	Me	p-(1,2,3-thiadiazol-5-yl)benzyloxy
221	OH	p-(1,2,3-thiadiazol-5-yl)benzyloxy
222	NH ₂	p-(1,2,3-thiadiazol-5-yl)benzyloxy
223	Me	o-(tetrazol-5-yl)benzyloxy
224	OH	o-(tetrazol-5-yl)benzyloxy
225	NH ₂	o-(tetrazol-5-yl)benzyloxy
226	Me	m-(tetrazol-5-yl)benzyloxy
227	OH	m-(tetrazol-5-yl)benzyloxy
228	NH ₂	m-(tetrazol-5-yl)benzyloxy
229	Me	[3-methyl-5-(tetrazol-5-yl)phenoxy]methyl
230	OH	[3-methyl-5-(tetrazol-5-yl)phenoxy]methyl
231	NH ₂	[3-methyl-5-(tetrazol-5-yl)phenoxy]methyl
232	Me	m-methyl-m-(tetrazol-5-yl)benzyloxy
233	OH	m-methyl-m-(tetrazol-5-yl)benzyloxy
234	NH ₂	m-methyl-m-(tetrazol-5-yl)benzyloxy
235	Me	2-oxo-2-phenylethoxy
236	OH	2-oxo-2-phenylethoxy
237	NH ₂	2-oxo-2-phenylethoxy
238	Me	carbo-t-butoxymethoxy
239	OH	carbo-t-butoxymethoxy
240	NH ₂	carbo-t-butoxymethoxy
241	Me	(benzimidazol-2-yl)methoxy
242	OH	(benzimidazol-2-yl)methoxy
243	NH ₂	(benzimidazol-2-yl)methoxy

244	Me	(imidazol-2-yl)methoxy
245	OH	(imidazol-2-yl)methoxy
246	NH ₂	(imidazol-2-yl)methoxy
247	Me	(1,4-dimethylimidazol-5-yl)methoxy
248	OH	(1,4-dimethylimidazol-5-yl)methoxy
249	NH ₂	(1,4-dimethylimidazol-5-yl)methoxy
250	Me	(thiazol-4-yl)methoxy
251	OH	(thiazol-4-yl)methoxy
252	NH ₂	(thiazol-4-yl)methoxy
253	Me	(quinolin-2-yl)methoxy
254	OH	(quinolin-2-yl)methoxy
255	NH ₂	(quinolin-2-yl)methoxy
256	Me	(1,3-benzodioxo-5-yl)methoxy
257	OH	(1,3-benzodioxo-5-yl)methoxy
258	NH ₂	(1,3-benzodioxo-5-yl)methoxy
259	Me	(3,5-dimethylisoxazol-4-yl)methoxy
260	OH	(3,5-dimethylisoxazol-4-yl)methoxy
261	NH ₂	(3,5-dimethylisoxazol-4-yl)methoxy
262	Me	(3,5-dimethylpyrazol-1-yl)methoxy
263	OH	(3,5-dimethylpyrazol-1-yl)methoxy
264	NH ₂	(3,5-dimethylpyrazol-1-yl)methoxy
265	Me	(1,3,5-trimethylpyrazol-4-yl)methoxy
266	OH	(1,3,5-trimethylpyrazol-4-yl)methoxy
267	NH ₂	(1,3,5-trimethylpyrazol-4-yl)methoxy

TABLE 6





Ex #	X	Y	R
1	CH ₂	CH ₂	H
2	CH ₂	O	H
3	O	CH ₂	H
4	CH ₂	CH ₂	methyl
5	CH ₂	O	methyl
6	O	CH ₂	methyl
7	CH ₂	CH ₂	ethyl
8	CH ₂	O	ethyl
9	O	CH ₂	ethyl
10	CH ₂	CH ₂	isopropyl
11	CH ₂	O	isopropyl
12	O	CH ₂	isopropyl
13	CH ₂	CH ₂	phenyl
14	CH ₂	O	phenyl
15	O	CH ₂	phenyl
16	CH ₂	CH ₂	benzyl
17	CH ₂	O	benzyl
18	O	CH ₂	benzyl
19	CH ₂	CH ₂	o-methylbenzyl
20	CH ₂	O	o-methylbenzyl
21	O	CH ₂	o-methylbenzyl
22	CH ₂	CH ₂	m-methylbenzyl
23	CH ₂	O	m-methylbenzyl
24	O	CH ₂	m-methylbenzyl

25	CH ₂	CH ₂	o,o-dimethylbenzyl
26	CH ₂	O	o,o-dimethylbenzyl
27	O	CH ₂	o,o-dimethylbenzyl
28	CH ₂	CH ₂	m,m-dimethylbenzyl
29	CH ₂	O	m,m-dimethylbenzyl
30	O	CH ₂	m,m-dimethylbenzyl
31	CH ₂	CH ₂	2-phenylethyl
32	CH ₂	O	2-phenylethyl
33	O	CH ₂	2-phenylethyl
34	CH ₂	CH ₂	2-(2-methylphenyl)ethyl
35	CH ₂	O	2-(2-methylphenyl)ethyl
36	O	CH ₂	2-(2-methylphenyl)ethyl
37	CH ₂	CH ₂	2-(3-methylphenyl)ethyl
38	CH ₂	O	2-(3-methylphenyl)ethyl
39	O	CH ₂	2-(3-methylphenyl)ethyl
40	CH ₂	CH ₂	2-(2,6-dimethylphenyl)ethyl
41	CH ₂	O	2-(2,6-dimethylphenyl)ethyl
42	O	CH ₂	2-(2,6-dimethylphenyl)ethyl
43	CH ₂	CH ₂	2-(3,5-dimethylphenyl)ethyl
44	CH ₂	O	2-(3,5-dimethylphenyl)ethyl
45	O	CH ₂	2-(3,5-dimethylphenyl)ethyl
46	CH ₂	CH ₂	2-(3-amino-5-methylphenyl)ethyl
47	CH ₂	O	2-(3-amino-5-methylphenyl)ethyl
48	O	CH ₂	2-(3-amino-5-methylphenyl)ethyl
49	CH ₂	CH ₂	2-(pyridin-4-yl)ethyl
50	CH ₂	O	2-(pyridin-4-yl)ethyl
51	O	CH ₂	2-(pyridin-4-yl)ethyl
52	CH ₂	CH ₂	2-(2,6-dimethylpyridin-4-yl)ethyl
53	CH ₂	O	2-(2,6-dimethylpyridin-4-yl)ethyl
54	O	CH ₂	2-(2,6-dimethylpyridin-4-yl)ethyl
55	CH ₂	CH ₂	2-(3,5-dimethylpyridin-4-yl)ethyl
56	CH ₂	O	2-(3,5-dimethylpyridin-4-yl)ethyl
57	O	CH ₂	2-(3,5-dimethylpyridin-4-yl)ethyl
58	CH ₂	CH ₂	styryl
59	CH ₂	O	styryl
60	O	CH ₂	styryl
61	CH ₂	CH ₂	hydroxy
62	CH ₂	O	hydroxy
63	O	CH ₂	hydroxy
64	CH ₂	CH ₂	methoxy
65	CH ₂	O	methoxy
66	O	CH ₂	methoxy
67	CH ₂	CH ₂	ethoxy
68	CH ₂	O	ethoxy
69	O	CH ₂	ethoxy
70	CH ₂	CH ₂	isopropoxy
71	CH ₂	O	isopropoxy
72	O	CH ₂	isopropoxy
73	CH ₂	CH ₂	tert-butoxy
74	CH ₂	O	tert-butoxy
75	O	CH ₂	tert-butoxy
76	CH ₂	CH ₂	cyclohexyloxy
77	CH ₂	O	cyclohexyloxy
78	O	CH ₂	cyclohexyloxy
79	CH ₂	CH ₂	phenoxy
80	CH ₂	O	phenoxy
81	O	CH ₂	phenoxy
82	CH ₂	CH ₂	o-methylphenoxy
83	CH ₂	O	o-methylphenoxy
84	O	CH ₂	o-methylphenoxy

85	CH ₂	CH ₂	m-methylphenoxy
86	CH ₂	O	m-methylphenoxy
87	O	CH ₂	m-methylphenoxy
88	CH ₂	CH ₂	o,o-dimethylphenoxy
89	CH ₂	O	o,o-dimethylphenoxy
90	O	CH ₂	o,o-dimethylphenoxy
91	CH ₂	CH ₂	m,m-dimethylphenoxy
92	CH ₂	O	m,m-dimethylphenoxy
93	O	CH ₂	m,m-dimethylphenoxy
94	CH ₂	CH ₂	cinnamyloxy
95	CH ₂	O	cinnamyloxy
96	O	CH ₂	cinnamyloxy
97	CH ₂	CH ₂	benzyloxy
98	CH ₂	O	benzyloxy
99	O	CH ₂	benzyloxy
100	CH ₂	CH ₂	phenoxymethyl
101	CH ₂	O	phenoxymethyl
102	O	CH ₂	phenoxymethyl
103	CH ₂	CH ₂	o-methylbenzyloxy
104	CH ₂	O	o-methylbenzyloxy
105	O	CH ₂	o-methylbenzyloxy
106	CH ₂	CH ₂	m-methylbenzyloxy
107	CH ₂	O	m-methylbenzyloxy
108	O	CH ₂	m-methylbenzyloxy
109	CH ₂	CH ₂	o,o-dimethylbenzyloxy
110	CH ₂	O	o,o-dimethylbenzyloxy
111	O	CH ₂	o,o-dimethylbenzyloxy
112	CH ₂	CH ₂	(2,6-dimethylphenoxy)methyl
113	CH ₂	O	(2,6-dimethylphenoxy)methyl
114	O	CH ₂	(2,6-dimethylphenoxy)methyl
115	CH ₂	CH ₂	m,m-dimethylbenzyloxy
116	CH ₂	O	m,m-dimethylbenzyloxy
117	O	CH ₂	m,m-dimethylbenzyloxy
118	CH ₂	CH ₂	(3,5-dimethylphenoxy)methyl
119	CH ₂	O	(3,5-dimethylphenoxy)methyl
120	O	CH ₂	(3,5-dimethylphenoxy)methyl
121	CH ₂	CH ₂	o,o-dicyanobenzyloxy
122	CH ₂	O	o,o-dicyanobenzyloxy
123	O	CH ₂	o,o-dicyanobenzyloxy
124	CH ₂	CH ₂	m,m-dicyanobenzyloxy
125	CH ₂	O	m,m-dicyanobenzyloxy
126	O	CH ₂	m,m-dicyanobenzyloxy
127	CH ₂	CH ₂	(2,6-dicyanophenoxy)methyl
128	CH ₂	O	(2,6-dicyanophenoxy)methyl
129	O	CH ₂	(2,6-dicyanophenoxy)methyl
130	CH ₂	CH ₂	(3,5-dicyanophenoxy)methyl
131	CH ₂	O	(3,5-dicyanophenoxy)methyl
132	O	CH ₂	(3,5-dicyanophenoxy)methyl
133	CH ₂	CH ₂	o-amino-o-cyanobenzyloxy
134	CH ₂	O	o-amino-o-cyanobenzyloxy
135	O	CH ₂	o-amino-o-cyanobenzyloxy
136	CH ₂	CH ₂	m-amino-m-cyanobenzyloxy
137	CH ₂	O	m-amino-m-cyanobenzyloxy
138	O	CH ₂	m-amino-m-cyanobenzyloxy
139	CH ₂	CH ₂	o-amino-o-nitrobenzyloxy
140	CH ₂	O	o-amino-o-nitrobenzyloxy
141	O	CH ₂	o-amino-o-nitrobenzyloxy
142	CH ₂	CH ₂	m-amino-m-nitrobenzyloxy
143	CH ₂	O	m-amino-m-nitrobenzyloxy
144	O	CH ₂	m-amino-m-nitrobenzyloxy

145	CH ₂	CH ₂	p-amino-m,m-dimethylbenzyloxy
146	CH ₂	O	p-amino-m,m-dimethylbenzyloxy
147	O	CH ₂	p-amino-m,m-dimethylbenzyloxy
148	CH ₂	CH ₂	o-amino-o-methylbenzyloxy
149	CH ₂	O	o-amino-o-methylbenzyloxy
150	O	CH ₂	o-amino-o-methylbenzyloxy
151	CH ₂	CH ₂	m-amino-m-methylbenzyloxy
152	CH ₂	O	m-amino-m-methylbenzyloxy
153	O	CH ₂	m-amino-m-methylbenzyloxy
154	CH ₂	CH ₂	o-cyano-o-methylbenzyloxy
155	CH ₂	O	o-cyano-o-methylbenzyloxy
156	O	CH ₂	o-cyano-o-methylbenzyloxy
157	CH ₂	CH ₂	m-cyano-m-methylbenzyloxy
158	CH ₂	O	m-cyano-m-methylbenzyloxy
159	O	CH ₂	m-cyano-m-methylbenzyloxy
160	CH ₂	CH ₂	o-cyano-o-nitrobenzyloxy
161	CH ₂	O	o-cyano-o-nitrobenzyloxy
162	O	CH ₂	o-cyano-o-nitrobenzyloxy
163	CH ₂	CH ₂	(2-cyano-6-nitrophenoxy)methyl
164	CH ₂	O	(2-cyano-6-nitrophenoxy)methyl
165	O	CH ₂	(2-cyano-6-nitrophenoxy)methyl
166	CH ₂	CH ₂	m-cyano-m-nitrobenzyloxy
167	CH ₂	O	m-cyano-m-nitrobenzyloxy
168	O	CH ₂	m-cyano-m-nitrobenzyloxy
169	CH ₂	CH ₂	(3-cyano-5-nitrophenoxy)methyl
170	CH ₂	O	(3-cyano-5-nitrophenoxy)methyl
171	O	CH ₂	(3-cyano-5-nitrophenoxy)methyl
172	CH ₂	CH ₂	m,m-dimethoxybenzyloxy
173	CH ₂	O	m,m-dimethoxybenzyloxy
174	O	CH ₂	m,m-dimethoxybenzyloxy
175	CH ₂	CH ₂	m,m-dichlorobenzyloxy
176	CH ₂	O	m,m-dichlorobenzyloxy
177	O	CH ₂	m,m-dichlorobenzyloxy
178	CH ₂	CH ₂	(3,5-dichlorophenoxy)methyl
179	CH ₂	O	(3,5-dichlorophenoxy)methyl
180	O	CH ₂	(3,5-dichlorophenoxy)methyl
181	CH ₂	CH ₂	m,m-dibromobenzyloxy
182	CH ₂	O	m,m-dibromobenzyloxy
183	O	CH ₂	m,m-dibromobenzyloxy
184	CH ₂	CH ₂	m,m-bis(trifluoromethyl)benzyloxy
185	CH ₂	O	m,m-bis(trifluoromethyl)benzyloxy
186	O	CH ₂	m,m-bis(trifluoromethyl)benzyloxy
187	CH ₂	CH ₂	[3,5-bis(trifluoromethyl)phenoxy)methyl
188	CH ₂	O	[3,5-bis(trifluoromethyl)phenoxy)methyl
189	O	CH ₂	[3,5-bis(trifluoromethyl)phenoxy)methyl
190	CH ₂	CH ₂	m-carboxamido-m-methylbenzyloxy
191	CH ₂	O	m-carboxamido-m-methylbenzyloxy
192	O	CH ₂	m-carboxamido-m-methylbenzyloxy
193	CH ₂	CH ₂	(3-carboxamido-5-methylphenoxy)methyl
194	CH ₂	O	(3-carboxamido-5-methylphenoxy)methyl
195	O	CH ₂	(3-carboxamido-5-methylphenoxy)methyl
196	CH ₂	CH ₂	m-hydroxycarbonyl-m-methylbenzyloxy
197	CH ₂	O	m-hydroxycarbonyl-m-methylbenzyloxy
198	O	CH ₂	m-hydroxycarbonyl-m-methylbenzyloxy
199	CH ₂	CH ₂	(3-hydroxycarbonyl-5-methylphenoxy)methyl
200	CH ₂	O	(3-hydroxycarbonyl-5-methylphenoxy)methyl
201	O	CH ₂	(3-hydroxycarbonyl-5-methylphenoxy)methyl
202	CH ₂	CH ₂	o-phenylbenzyloxy
203	CH ₂	O	o-phenylbenzyloxy
204	O	CH ₂	o-phenylbenzyloxy

205	CH ₂	CH ₂	m-phenylbenzyloxy
206	CH ₂	O	m-phenylbenzyloxy
207	O	CH ₂	m-phenylbenzyloxy
208	CH ₂	CH ₂	(naphth-1-yl)methoxy
209	CH ₂	O	(naphth-1-yl)methoxy
210	O	CH ₂	(naphth-1-yl)methoxy
211	CH ₂	CH ₂	(naphth-2-yl)methoxy
212	CH ₂	O	(naphth-2-yl)methoxy
213	O	CH ₂	(naphth-2-yl)methoxy
214	CH ₂	CH ₂	(2-methylnaphth-1-yl)methoxy
215	CH ₂	O	(2-methylnaphth-1-yl)methoxy
216	O	CH ₂	(2-methylnaphth-1-yl)methoxy
217	CH ₂	CH ₂	(4-methylnaphth-2-yl)methoxy
218	CH ₂	O	(4-methylnaphth-2-yl)methoxy
219	O	CH ₂	(4-methylnaphth-2-yl)methoxy
220	CH ₂	CH ₂	(pyridin-3-yl)methoxy
221	CH ₂	O	(pyridin-3-yl)methoxy
222	O	CH ₂	(pyridin-3-yl)methoxy
223	CH ₂	CH ₂	(pyridin-4-yl)methoxy
224	CH ₂	O	(pyridin-4-yl)methoxy
225	O	CH ₂	(pyridin-4-yl)methoxy
226	CH ₂	CH ₂	(3,5-dichloropyridin-4-yl)methoxy
227	CH ₂	O	(3,5-dichloropyridin-4-yl)methoxy
228	O	CH ₂	(3,5-dichloropyridin-4-yl)methoxy
229	CH ₂	CH ₂	(3,5-dimethylpyridin-4-yl)methoxy
230	CH ₂	O	(3,5-dimethylpyridin-4-yl)methoxy
231	O	CH ₂	(3,5-dimethylpyridin-4-yl)methoxy
232	CH ₂	CH ₂	(1,2,3-benzotriazol-1-yl)methoxy
233	CH ₂	O	(1,2,3-benzotriazol-1-yl)methoxy
234	O	CH ₂	(1,2,3-benzotriazol-1-yl)methoxy
235	CH ₂	CH ₂	benzhydroxy
236	CH ₂	O	benzhydroxy
237	O	CH ₂	benzhydroxy
238	CH ₂	CH ₂	p-(1,2,3-thiadiazol-5-yl)benzyloxy
239	CH ₂	O	p-(1,2,3-thiadiazol-5-yl)benzyloxy
240	O	CH ₂	p-(1,2,3-thiadiazol-5-yl)benzyloxy
241	CH ₂	CH ₂	o-(tetrazol-5-yl)benzyloxy
242	CH ₂	O	o-(tetrazol-5-yl)benzyloxy
243	O	CH ₂	o-(tetrazol-5-yl)benzyloxy
244	CH ₂	CH ₂	m-(tetrazol-5-yl)benzyloxy
245	CH ₂	O	m-(tetrazol-5-yl)benzyloxy
246	O	CH ₂	m-(tetrazol-5-yl)benzyloxy
247	CH ₂	CH ₂	[3-methyl-5-(tetrazol-5-yl)phenoxy]methyl
248	CH ₂	O	[3-methyl-5-(tetrazol-5-yl)phenoxy]methyl
249	O	CH ₂	[3-methyl-5-(tetrazol-5-yl)phenoxy]methyl
250	CH ₂	CH ₂	m-methyl-m-(tetrazol-5-yl)benzyloxy
251	CH ₂	O	m-methyl-m-(tetrazol-5-yl)benzyloxy
252	O	CH ₂	m-methyl-m-(tetrazol-5-yl)benzyloxy
253	CH ₂	CH ₂	2-oxo-2-phenylethoxy
254	CH ₂	O	2-oxo-2-phenylethoxy
255	O	CH ₂	2-oxo-2-phenylethoxy
256	CH ₂	CH ₂	carbo-t-butoxymethoxy
257	CH ₂	O	carbo-t-butoxymethoxy
258	O	CH ₂	carbo-t-butoxymethoxy
259	CH ₂	CH ₂	(benzimidazol-2-yl)methoxy
260	CH ₂	O	(benzimidazol-2-yl)methoxy
261	O	CH ₂	(benzimidazol-2-yl)methoxy
262	CH ₂	CH ₂	(imidazol-2-yl)methoxy
263	CH ₂	O	(imidazol-2-yl)methoxy
264	O	CH ₂	(imidazol-2-yl)methoxy

265	CH ₂	CH ₂	(1,4-dimethylimidazol-5-yl)methoxy
266	CH ₂	O	(1,4-dimethylimidazol-5-yl)methoxy
267	O	CH ₂	(1,4-dimethylimidazol-5-yl)methoxy
268	CH ₂	CH ₂	(thiazol-4-yl)methoxy
269	CH ₂	O	(thiazol-4-yl)methoxy
270	O	CH ₂	(thiazol-4-yl)methoxy
271	CH ₂	CH ₂	(quinolin-2-yl)methoxy
272	CH ₂	O	(quinolin-2-yl)methoxy
273	O	CH ₂	(quinolin-2-yl)methoxy
274	CH ₂	CH ₂	(1,3-benzodioxo-5-yl)methoxy
275	CH ₂	O	(1,3-benzodioxo-5-yl)methoxy
276	O	CH ₂	(1,3-benzodioxo-5-yl)methoxy
277	CH ₂	CH ₂	(3,5-dimethylisoxazol-4-yl)methoxy
278	CH ₂	O	(3,5-dimethylisoxazol-4-yl)methoxy
279	O	CH ₂	(3,5-dimethylisoxazol-4-yl)methoxy
280	CH ₂	CH ₂	(3,5-dimethylpyrazol-1-yl)methoxy
281	CH ₂	O	(3,5-dimethylpyrazol-1-yl)methoxy
282	O	CH ₂	(3,5-dimethylpyrazol-1-yl)methoxy
283	CH ₂	CH ₂	(1,3,5-trimethylpyrazol-4-yl)methoxy
284	CH ₂	O	(1,3,5-trimethylpyrazol-4-yl)methoxy
285	O	CH ₂	(1,3,5-trimethylpyrazol-4-yl)methoxy

UTILITY

The compounds of formula I are expected to be metalloproteinase inhibitors. The MMP-3 inhibitory activity of the compounds of the present invention is demonstrated using assays of MMP-3 activity, for example, using the assay described below for assaying inhibitors of MMP-3 activity. The compounds of the present invention are expected to be bioavailable in vivo as demonstrated, for example, using the ex vivo assay described below. The compounds of formula I are expected to have the ability to suppress/inhibit cartilage degradation in vivo, for example, as demonstrated using the animal model of acute cartilage degradation described below.

The compounds provided by this invention should also be useful as standards and reagents in determining the ability of a potential pharmaceutical to inhibit MPs. These would be provided in commercial kits comprising a compound of this invention.

Metalloproteinases have also been implicated in the degradation of basement membranes to allow infiltration of cancer cells into the circulation and subsequent penetration into other tissues leading to tumor metastasis. (Stetler-Stevenson, Cancer and Metastasis Reviews, 9, 289-303, 1990.) The compounds of the present invention should be useful for the prevention and treatment of invasive tumors by inhibition of this aspect of metastasis.

The compounds of the present invention should also have utility for the prevention and treatment of osteopenia associated with matrix metalloproteinase-mediated breakdown of cartilage and bone which occurs in osteoporosis patients.

Compounds which inhibit the production or action of TNF and/or Aggrecanase and/or MP's are potentially useful for the treatment or prophylaxis of various inflammatory, infectious, immunological or malignant diseases. These include, but are not limited to inflammation, fever, cardiovascular effects, hemorrhage, coagulation and acute phase response, an acute infection, septic shock, haemodynamic shock and sepsis syndrome, post ischaemic reperfusion injury, malaria, Crohn's disease, mycobacterial infection, meningitis, psoriasis,

periodontitis, gingivitis, congestive heart failure, fibrotic disease, cachexia, and anorexia, graft rejection, cancer, corneal ulceration or tumor invasion by secondary metastases, autoimmune disease, skin inflammatory diseases, multiple osteo
5 and rheumatoid arthritis, multiple sclerosis, radiation damage, HIV, and hyperoxic alveolar injury.

Some compounds of the present invention have been shown to inhibit TNF production in lipopolysaccharide stimulated mice, for example, using the assay for TNF Induction in Mice
10 and in human whole blood as described below.

Some compounds of the present invention have been shown to inhibit aggrecanase a key enzyme in cartilage breakdown as determined by the aggrecanase assay described below.

As used herein "µg" denotes microgram, "mg" denotes
15 milligram, "g" denotes gram, "µL" denotes microliter, "mL" denotes milliliter, "L" denotes liter, "nM" denotes nanomolar, "µM" denotes micromolar, "mM" denotes millimolar, "M" denotes molar and "nm" denotes nanometer. "Sigma" stands for the Sigma-Aldrich Corp. of St. Louis, MO.

20 A compound is considered to be active if it has an IC₅₀ or K_i value of less than about 1 mM for the inhibition of MMP-3.

Aggrecanase Enzymatic Assay

25 A novel enzymatic assay was developed to detect potential inhibitors of aggrecanase. The assay uses active aggrecanase accumulated in media from stimulated bovine nasal cartilage (BNC) or related cartilage sources and purified cartilage aggrecan monomer or a fragment thereof as a substrate.

30 The substrate concentration, amount of aggrecanase time of incubation and amount of product loaded for Western analysis were optimized for use of this assay in screening putative aggrecanase inhibitors. Aggrecanase is generated by stimulation of cartilage slices with interleukin-1 (IL-1),
35 tumor necrosis factor alpha (TNFα) or other stimuli. Matrix metalloproteinases (MMPs) are secreted from cartilage in an inactive, zymogen form following stimulation, although active enzymes are present within the matrix. We have shown that

following depletion of the extracellular aggrecan matrix, active MMPs are released into the culture media. (Tortorella, M.D. et. al. Trans. Ortho. Res. Soc. 20, 341, 1995).

Therefore, in order to accumulate BNC aggrecanase in culture media, cartilage is first depleted of endogenous aggrecan by stimulation with 500 ng/ml human recombinant IL- β for 6 days with media changes every 2 days. Cartilage is then stimulated for an additional 8 days without media change to allow accumulation of soluble, active aggrecanase in the culture media. In order to decrease the amounts of other matrix metalloproteinases released into the media during aggrecanase accumulation, agents which inhibit MMP-1, -2, -3, and -9 biosynthesis are included during stimulation. This BNC conditioned media, containing aggrecanase activity is then used as the source of aggrecanase for the assay. Aggrecanase enzymatic activity is detected by monitoring production of aggrecan fragments produced exclusively by cleavage at the Glu373-Ala374 bond within the aggrecan core protein by Western analysis using the monoclonal antibody, BC-3 (Hughes, CE, et al., Biochem J 306:799-804, 1995). This antibody recognizes aggrecan fragments with the N-terminus, 374ARGSVIL, generated upon cleavage by aggrecanase. The BC-3 antibody recognizes this neoepitope only when it is at the N-terminus and not when it is present internally within aggrecan fragments or within the aggrecan protein core. Other proteases produced by cartilage in response to IL-1 do not cleave aggrecan at the Glu373-Ala374 aggrecanase site; therefore, only products produced upon cleavage by aggrecanase are detected. Kinetic studies using this assay yield a K_m of 1.5 +/- 0.35 μM for aggrecanase.

To evaluate inhibition of aggrecanase, compounds are prepared as 10 mM stocks in DMSO, water or other solvents and diluted to appropriate concentrations in water. Drug (50 μl) is added to 50 μl of aggrecanase-containing media and 50 μl of 2 mg/ml aggrecan substrate and brought to a final volume of 200 μl in 0.2 M Tris, pH 7.6, containing 0.4 M NaCl and 40 mM $CaCl_2$. The assay is run for 4 hr at 37°C, quenched with 20 mM EDTA and analyzed for aggrecanase-generated products. A

sample containing enzyme and substrate without drug is included as a positive control and enzyme incubated in the absence of substrate serves as a measure of background.

Removal of the glycosaminoglycan side chains from
5 aggrecan is necessary for the BC-3 antibody to recognize the
ARGSVIL epitope on the core protein. Therefore, for analysis
of aggrecan fragments generated by cleavage at the Glu373-
Ala374 site, proteoglycans and proteoglycan fragments are
enzymatically deglycosylated with chondroitinase ABC (0.1
10 units/10 ug GAG) for 2 hr at 37°C and then with keratanase
(0.1 units/10 ug GAG) and keratanase II (0.002 units/10 ug
GAG) for 2 hr at 37°C in buffer containing 50 mM sodium
acetate, 0.1 M Tris/HCl, pH 6.5. After digestion, aggrecan in
the samples is precipitated with 5 volumes of acetone and
15 resuspended in 30 ul of Tris glycine SDS sample buffer (Novex)
containing 2.5% beta mercaptoethanol. Samples are loaded and
then separated by SDS-PAGE under reducing conditions with 4-
12% gradient gels, transferred to nitrocellulose and
immunolocated with 1:500 dilution of antibody BC3.
20 Subsequently, membranes are incubated with a 1:5000 dilution
of goat anti-mouse IgG alkaline phosphatase second antibody
and aggrecan catabolites visualized by incubation with
appropriate substrate for 10-30 minutes to achieve optimal
color development. Blots are quantitated by scanning
25 densitometry and inhibition of aggrecanase determined by
comparing the amount of product produced in the presence
versus absence of compound.

Bisacetylated Substance P / MMP-3 fluorescent Assay

30 A high capacity enzymatic assay was developed to detect
potential inhibitors of MMP-3. The assay uses a derivative of
a peptide substrate, substance P (Arg-Pro-Lys-Pro-Gln-Gln-Phe-
Phe-Gly-Leu-Met), which is cleaved by MMP-3 exclusively at the
glutamine-phenylalanine bond. In order to adapt this assay
35 for high throughput screening, we have developed a
fluorimetric method of product detection. The production of
the hydrolysis product, substance P 7-11, is measured by
reaction with fluorecamine, a fluorogenic compound which

reacts with the primary amine of this fragment. The substance P substrate is bisacetylated to block the primary amines of the intact substrate. Thus, the resulting fluorescence represents generation of product (7-11 peptide) formed upon cleavage by MMP-3, and is quantitated using a standard curve prepared with known concentrations of 7-11 peptide. Kinetic studies using the bisacetylated substrate yield the following parameters for MMP-3: $K_m = 769 \pm 52 \mu M$; $V_{max} = 0.090 \pm 0.003$ nmoles 7-11 peptide/min.

To evaluate inhibition of MMP-3, compounds were prepared at a concentration of 10 mM in 100% methanol, and then further diluted to a 20X molar stock. Five microliters of each drug stock was added to the assay in the presence of 20 nM truncated MMP-3 in 67.5 mM tricine (pH 7.5), 10 mM $CaCl_2$, 40 mM NaCl, and 0.005% Brij 35 in a final volume of 100 microliters. Bisacetylated substance P (1000 nM) was added, and the assay was run for 1 hour at 25°C. The reaction was quenched with EDTA (20 mM) and product was detected fluorometrically following addition of fluoescamine (0.075 mg/ml). Fluorescence of each sample was converted to an amount of product formed using a substance P 7-11 standard curve. Under these conditions, the assay is linear with respect to MMP-3 amount up to 10 pmoles. Inhibition of MMP-3 was determined by comparing the amount of product generated in the presence and absence of compound.

Selected compounds of the present invention were tested and shown to have activity in the above assay.

Ex vivo assay for bioavailability of MMP-3 inhibitors

Blood was collected by cardiac puncture from rats at different times after dosing I.V., I.P., or P.O. with compound in order to determine the levels of inhibitor present. Plasma was extracted with 10% TCA in 95% methanol, and placed on ice for 10 minutes. The plasma was then centrifuged for 15 minutes at 14,000 rpm in an Eppendorf microcentrifuge. The supernatant was removed, recentrifuged, and the resulting supernatant was diluted 1:10 in 50 mM tricine, pH 8.5. The pH of the sample was adjusted to 7.5, and then assayed in the

MMP-3 substance P fluorescent enzymatic assay. Plasma from naive rats was extracted by the same method and used as a negative control. This plasma was also used to prepare a spiked plasma curve of the compound of interest. Known concentrations of the compound were added to control plasma, the plasma was extracted by the same method, and then assayed in the MMP-3 enzymatic assay. A standard curve was prepared that related percent inhibition in the MMP-3 assay to the concentration of drug added in the spiked samples. Based on the percent inhibition in the presence of plasma from dosed rats, the concentration of compound was determined using the standard curve.

Acute Cartilage Degradation Rat Model

A novel in vivo model of acute cartilage degradation in rats has been characterized as a method to determine the proteoglycan content in the synovial fluid after the induction of cartilage degradation. Experimental groups exhibit increased levels of proteoglycan content in their synovial fluid versus control rats. The criteria to demonstrate a compound's activity in this model, is the ability to inhibit the demonstration of cartilage degradation, as measured by increased proteoglycan content in the synovial fluid of rats after compound administration. Indomethacin, a non-steroidal anti-inflammatory drug is inactive in this model. Indomethacin administration does not inhibit the demonstration of cartilage degradation in experimental animals. In contrast, administration of a compound of this invention significantly inhibited the demonstration of cartilage degradation in this model.

TNF Human Whole Blood Assay

Blood is drawn from normal donors into tubes containing 143 USP units of heparin/10ml. 225ul of blood is plated directly into sterile polypropylene tubes. Compounds are diluted in DMSO/serum free media and added to the blood samples so the final concentration of compounds are 50, 10, 5, 1, .5, .1, and .01 μ M. The final concentration of DMSO does

not exceed .5%. Compounds are preincubated for 15 minutes before the addition of 100ng/ml LPS. Plates are incubated for 5 hours in an atmosphere of 5% CO₂ in air. At the end of 5 hours, 750ul of serum free media is added to each tube and the samples are spun at 1200RPM for 10 minutes. The supernatant is collected off the top and assayed for TNF-alpha production by a standard sandwich ELISA. The ability of compounds to inhibit TNF-alpha production by 50% compared to DMSO treated cultures is given by the IC50 value.

TNF Induction In Mice

Test compounds are administered to mice either I.P. or P.O. at time zero. Immediately following compound administration, mice receive an I.P. injection of 20 mg of D-galactosamine plus 10 µg of lipopolysaccharide. One hour later, animals are anesthetized and bled by cardiac puncture. Blood plasma is evaluated for TNF levels by an ELISA specific for mouse TNF. Administration of representative compounds of the present invention to mice results in a dose-dependent suppression of plasma TNF levels at one hour in the above assay.

Dosage and Formulation

The compounds of the present invention can be administered orally using any pharmaceutically acceptable dosage form known in the art for such administration. The active ingredient can be supplied in solid dosage forms such as dry powders, granules, tablets or capsules, or in liquid dosage forms, such as syrups or aqueous suspensions. The active ingredient can be administered alone, but is generally administered with a pharmaceutical carrier. A valuable treatise with respect to pharmaceutical dosage forms is Remington's Pharmaceutical Sciences, Mack Publishing.

The compounds of the present invention can be administered in such oral dosage forms as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. Likewise, they may also

be administered in intravenous (bolus or infusion), intraperitoneal, subcutaneous, or intramuscular form, all using dosage forms well known to those of ordinary skill in the pharmaceutical arts. An effective but non-toxic amount of the compound desired can be employed as an antiinflammatory and antiarthritic agent.

The compounds of this invention can be administered by any means that produces contact of the active agent with the agent's site of action, MMP-3, in the body of a mammal. They can be administered by any conventional means available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or in a combination of therapeutic agents. They can be administered alone, but generally administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

The dosage regimen for the compounds of the present invention will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; the route of administration, the renal and hepatic function of the patient, and the effect desired. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug required to prevent, counter, or arrest the progress of the condition.

By way of general guidance, the daily oral dosage of each active ingredient, when used for the indicated effects, will range between about 0.001 to 1000 mg/kg of body weight, preferably between about 0.01 to 100 mg/kg of body weight per day, and most preferably between about 1.0 to 20 mg/kg/day. For a normal male adult human of approximately 70 kg of body weight, this translates into a dosage of 70 to 1400 mg/day. Intravenously, the most preferred doses will range from about 1 to about 10 mg/kg/minute during a constant rate infusion. Advantageously, compounds of the present invention may be

administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily.

5 The compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, 10 of course, be continuous rather than intermittent throughout the dosage regimen.

In the methods of the present invention, the compounds herein described in detail can form the active ingredient, and are typically administered in admixture with suitable 15 pharmaceutical diluents, excipients, or carriers (collectively referred to herein as carrier materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

20 For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium 25 sulfate, mannitol, sorbitol and the like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Moreover, when desired or necessary, suitable binders, 30 lubricants, disintegrating agents, and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth, or sodium alginate, 35 carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrators

include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

Compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacylates, and crosslinked or amphipathic block copolymers of hydrogels.

Dosage forms (pharmaceutical compositions) suitable for administration may contain from about 1 milligram to about 100 milligrams of active ingredient per dosage unit. In these pharmaceutical compositions the active ingredient will ordinarily be present in an amount of about 0.5-95% by weight based on the total weight of the composition.

The active ingredient can be administered orally in solid dosage forms, such as capsules, tablets, and powders, or in liquid dosage forms, such as elixirs, syrups, and suspensions. It can also be administered parenterally, in sterile liquid dosage forms.

Gelatin capsules may contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release

products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective
5 disintegration in the gastrointestinal tract.

Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as

10 propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as
15 sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propyl-paraben, and
20 chlorobutanol.

Suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences, Mack Publishing Company, a standard reference text in this field.

Useful pharmaceutical dosage-forms for administration of the
25 compounds of this invention can be illustrated as follows:

Capsules

Capsules are prepared by conventional procedures so that the dosage unit is 500 milligrams of active ingredient, 100
30 milligrams of cellulose and 10 milligrams of magnesium stearate.

A large number of unit capsules may also prepared by filling standard two-piece hard gelatin capsules each with 100 milligrams of powdered active ingredient, 150 milligrams of
35 lactose, 50 milligrams of cellulose, and 6 milligrams magnesium stearate.

Syrup

	<u>Wt. %</u>
Active Ingredient	10
Liquid Sugar	50
Sorbitol	20
Glycerine	5
Flavor, Colorant and Preservative	as required
Water	as required

The final volume is brought up to 100% by the addition of distilled water.

Aqueous Suspension

	<u>Wt. %</u>
Active Ingredient	10
Sodium Saccharin	0.01
Keltrol® (Food Grade Xanthan Gum)	0.2
Liquid Sugar	5
Flavor, Colorant and Preservative	as required
Water	as required

Xanthan gum is slowly added into distilled water before adding the active ingredient and the rest of the formulation ingredients. The final suspension is passed through a homogenizer to assure the elegance of the final products.

Resuspendable Powder

	<u>Wt. %</u>
Active Ingredient	50.0
Lactose	35.0
Sugar	10.0
Acacia	4.7
Sodium Carboxymethylcellulose	0.3

Each ingredient is finely pulverized and then uniformly mixed together. Alternatively, the powder can be prepared as a suspension and then spray dried.

Semi-Solid Gel

	<u>Wt. %</u>
Active Ingredient	10
Sodium Saccharin	0.02
Gelatin	2
Flavor, Colorant and Preservative	as required
Water	as required

Gelatin is prepared in hot water. The finely pulverized active ingredient is suspended in the gelatin solution and then the rest of the ingredients are mixed in. The suspension is filled

into a suitable packaging container and cooled down to form the gel.

Semi-Solid Paste

5		<u>Wt. %</u>
	Active Ingredient	10
	Gelcarin® (Carrageenin gum)	1
	Sodium Saccharin	0.01
	Gelatin	2
10	Flavor, Colorant and	as required
	Preservative	
	Water	as required

15 Gelcarin® is dissolved in hot water (around 80°C) and then the fine-powder active ingredient is suspended in this solution. Sodium saccharin and the rest of the formulation ingredients are added to the suspension while it is still warm. The
20 suspension is homogenized and then filled into suitable containers.

Emulsifiable Paste

		<u>Wt. %</u>
	Active Ingredient	30
25	Tween® 80 and Span® 80	6
	Keltrol®	0.5
	Mineral Oil	63.5

30 All the ingredients are carefully mixed together to make a homogenous paste.

Soft Gelatin Capsules

A mixture of active ingredient in a digestable oil such as soybean oil, cottonseed oil or olive oil is prepared and
35 injected by means of a positive displacement pump into gelatin to form soft gelatin capsules containing 100 milligrams of the active ingredient. The capsules are washed and dried.

Tablets

40 Tablets may be prepared by conventional procedures so that the dosage unit is 500 milligrams of active ingredient, 150 milligrams of lactose, 50 milligrams of cellulose and 10 milligrams of magnesium stearate.

A large number of tablets may also be prepared by
45 conventional procedures so that the dosage unit was 100 milligrams of active ingredient, 0.2 milligrams of colloidal silicon dioxide, 5 milligrams of magnesium stearate, 275

milligrams of microcrystalline cellulose, 11 milligrams of starch and 98.8 milligrams of lactose. Appropriate coatings may be applied to increase palatability or delay absorption.

5 Injectable

A parenteral composition suitable for administration by injection is prepared by stirring 1.5% by weight of active ingredient in 10% by volume propylene glycol and water. The solution is made isotonic with sodium chloride and sterilized.

10

Suspension

An aqueous suspension is prepared for oral administration so that each 5 mL contain 100 mg of finely divided active ingredient, 200 mg of sodium carboxymethyl cellulose, 5 mg of sodium benzoate, 1.0 g of sorbitol solution, U.S.P., and 0.025 mL of vanillin.

The compounds of the present invention may be administered in combination with a second therapeutic agent, especially non-steroidal anti-inflammatory drugs (NSAID's). The compound of Formula I and such second therapeutic agent can be administered separately or as a physical combination in a single dosage unit, in any dosage form and by various routes of administration, as described above.

The compound of Formula I may be formulated together with the second therapeutic agent in a single dosage unit (that is, combined together in one capsule, tablet, powder, or liquid, etc.). When the compound of Formula I and the second therapeutic agent are not formulated together in a single dosage unit, the compound of Formula I and the second therapeutic agent may be administered essentially at the same time, or in any order; for example the compound of Formula I may be administered first, followed by administration of the second agent. When not administered at the same time, preferably the administration of the compound of Formula I and the second therapeutic agent occurs less than about one hour apart, more preferably less than about 5 to 30 minutes apart.

Preferably the route of administration of the compound of Formula I is oral. Although it is preferable that the

compound of Formula I and the second therapeutic agent are both administered by the same route (that is, for example, both orally), if desired, they may each be administered by different routes and in different dosage forms (that is, for example, one component of the combination product may be administered orally, and another component may be administered intravenously).

The dosage of the compound of Formula I when administered alone or in combination with a second therapeutic agent may vary depending upon various factors such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration, the age, health and weight of the recipient, the nature and extent of the symptoms, the kind of concurrent treatment, the frequency of treatment, and the effect desired, as described above. Particularly when provided as a single dosage unit, the potential exists for a chemical interaction between the combined active ingredients. For this reason, when the compound of Formula I and a second therapeutic agent are combined in a single dosage unit they are formulated such that although the active ingredients are combined in a single dosage unit, the physical contact between the active ingredients is minimized (that is, reduced). For example, one active ingredient may be enteric coated. By enteric coating one of the active ingredients, it is possible not only to minimize the contact between the combined active ingredients, but also, it is possible to control the release of one of these components in the gastrointestinal tract such that one of these components is not released in the stomach but rather is released in the intestines. One of the active ingredients may also be coated with a sustained-release material which effects a sustained-release throughout the gastrointestinal tract and also serves to minimize physical contact between the combined active ingredients. Furthermore, the sustained-released component can be additionally enteric coated such that the release of this component occurs only in the intestine. Still another approach would involve the formulation of a combination product in which the one

component is coated with a sustained and/or enteric release polymer, and the other component is also coated with a polymer such as a lowviscosity grade of hydroxypropyl methylcellulose (HPMC) or other appropriate materials as known in the art, in order to further separate the active components. The polymer coating serves to form an additional barrier to interaction with the other component.

These as well as other ways of minimizing contact between the components of combination products of the present invention, whether administered in a single dosage form or administered in separate forms but at the same time by the same manner, will be readily apparent to those skilled in the art, once armed with the present disclosure.


The present invention also includes pharmaceutical kits useful, for example, in the treatment or prevention of osteoarthritis or rheumatoid arthritis, which comprise one or more containers containing a pharmaceutical composition comprising a therapeutically effective amount of a compound of Formula I. Such kits may further include, if desired, one or more of various conventional pharmaceutical kit components, such as, for example, containers with one or more pharmaceutically acceptable carriers, additional containers, etc., as will be readily apparent to those skilled in the art. Instructions, either as inserts or as labels, indicating quantities of the components to be administered, guidelines for administration, and/or guidelines for mixing the components, may also be included in the kit.

In the present disclosure it should be understood that the specified materials and conditions are important in practicing the invention but that unspecified materials and conditions are not excluded so long as they do not prevent the benefits of the invention from being realized.

Although this invention has been described with respect to specific embodiments, the details of these embodiments are not to be construed as limitations. Various equivalents, changes and modifications may be made without departing from the spirit and scope of this invention, and it is understood that such equivalent embodiments are part of this invention.

WHAT IS CLAIMED AS NEW AND DESIRED TO BE SECURED BY
LETTER PATENT OF UNITED STATES IS:

1. A compound of formula I:



I

or a stereoisomer or pharmaceutically acceptable salt form thereof, wherein;

A is selected from COR^5 , $-\text{CO}_2\text{H}$, $\text{CH}_2\text{CO}_2\text{H}$, $-\text{CO}_2\text{R}^6$, $-\text{CONHOH}$, $-\text{CONHOR}^5$, $-\text{CONHOR}^6$, $-\text{NHR}^a$, $-\text{N}(\text{OH})\text{COR}^5$, $-\text{SH}$, $-\text{CH}_2\text{SH}$, $-\text{SO}_2\text{NHR}^a$, $\text{SN}_2\text{H}_2\text{R}^a$, $\text{PO}(\text{OH})_2$, and $\text{PO}(\text{OH})\text{NHR}^a$;

ring B is a 4-8 membered cyclic amide containing from 0-3 additional heteroatoms selected from O, NR^a , and $\text{S}(\text{O})_p$, 0-1 additional carbonyl groups and 0-1 double bonds;

R^1 is $\text{U-X-Y-Z-U}^a\text{-X}^a\text{-Y}^a\text{-Z}^a$;

U is absent or is selected from: O, NR^a , $\text{C}(\text{O})$, $\text{C}(\text{O})\text{O}$, $\text{OC}(\text{O})$, $\text{C}(\text{O})\text{NR}^a$, $\text{NR}^a\text{C}(\text{O})$, $\text{OC}(\text{O})\text{O}$, $\text{OC}(\text{O})\text{NR}^a$, $\text{NR}^a\text{C}(\text{O})\text{O}$, $\text{NR}^a\text{C}(\text{O})\text{NR}^a$, $\text{S}(\text{O})_p$, $\text{S}(\text{O})_p\text{NR}^a$, $\text{NR}^a\text{S}(\text{O})_p$, and $\text{NR}^a\text{SO}_2\text{NR}^a$;

X is absent or selected from C_{1-10} alkylene, C_{2-10} alkenylene, and C_{2-10} alkynylene;

Y is absent or selected from O, NR^a , $\text{S}(\text{O})_p$, and $\text{C}(\text{O})$;

Z is absent or selected from a C_{3-13} carbocyclic residue substituted with 0-5 R^b and a 5-14 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with 0-5 R^b ;

U^a is absent or is selected from: O, NR^a, C(O), C(O)O, OC(O), C(O)NR^a, NR^aC(O), OC(O)O, OC(O)NR^a, NR^aC(O)O, NR^aC(O)NR^a, S(O)_p, S(O)_pNR^a, NR^aS(O)_p, and NR^aSO₂NR^a;

5 X^a is absent or selected from C₁₋₁₀ alkylene, C₂₋₁₀ alkenylene, C₂₋₁₀ alkynylene;

Y^a is absent or selected from O, NR^a, S(O)_p, and C(O);

10 Z^a is selected from H, a C₃₋₁₃ carbocyclic residue substituted with 0-5 R^c and a 5-14 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with 0-5 R^c;

15 R² is selected from H, Q', C₁₋₁₀ alkylene-Q', C₂₋₁₀ alkenylene-Q', C₂₋₁₀ alkynylene-Q', (CRR')_rO(CRR')_r-Q', (CRR')_rNR^a(CRR')_r-Q', (CRR')_rNR^aC(O)(CRR')_r-Q', (CRR')_rC(O)NR^a(CRR')_r-Q', (CRR')_rC(O)(CRR')_r-Q', (CRR')_rC(O)O(CRR')_r-Q', (CRR')_rS(O)_p(CRR')_r-Q',
 20 (CRR')_rSO₂NR^a(CRR')_r-Q', (CRR')_rNR^aC(O)NR^a(CRR')_r-Q', (CRR')_rOC(O)NR^a(CRR')_r-Q', and (CRR')_rNR^aC(O)O(CRR')_r-Q';

R, at each occurrence, is independently selected from H, CH₃,
 25 CH₂CH₃, CH=CH₂, CH=CHCH₃, and CH₂CH=CH₂;

R', at each occurrence, is independently selected from H, CH₃, CH₂CH₃, and CH(CH₃)₂;

30 alternatively, R¹ and R² combine to form a C₃₋₁₃ carbocyclic residue substituted with R^{1'} and 0-3 R^b or a 5-14 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with R^{1'} and 0-3 R^b;

35

Q' is selected from H, a C₃₋₁₃ carbocyclic residue substituted with 0-5 R^b and a 5-14 membered heterocyclic system

containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with 0-5 R^b;

R^{1'} is U^a-X^a-Y^a-Z^a;

5

R³ is selected from H, Q, C₁₋₁₀ alkylene-Q, C₂₋₁₀ alkenylene-Q, C₂₋₁₀ alkynylene-Q, (CRR')_rO(CRR')_r-Q, (CRR')_rNR^a(CRR')_r-Q, (CRR')_rC(O)(CRR')_r-Q, (CRR')_rC(O)O(CRR')_r-Q, (CRR')_rOC(O)(CRR')_r-Q, (CRR')_rC(O)NR^a(CRR')_r-Q, (CRR')_rNR^aC(O)(CRR')_r-Q, (CRR')_rOC(O)O(CRR')_r-Q, (CRR')_rOC(O)NR^a(CRR')_r-Q, (CRR')_rNR^aC(O)O(CRR')_r-Q, (CRR')_rNR^aC(O)NR^a(CRR')_r-Q, (CRR')_rS(O)_p(CRR')_r-Q, (CRR')_rSO₂NR^a(CRR')_r-Q, (CRR')_rNR^aSO₂(CRR')_r-Q, (CRR')_rNR^aSO₂NR^a(CRR')_r-Q, (CRR')_rNR^aC(O)(CRR')_rNHQ, (CRR')_rNR^aC(O)(CRR')_rNHC(O)OR^a, and (CRR')_rNR^aC(O)(CRR')_rNHC(O)(CRR')_rNHC(O)OR^a,

10

15

Q is selected from H, a C₃₋₁₃ carbocyclic residue substituted with 0-5 R^b and a 5-14 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with 0-5 R^b;

20

R⁴ is selected from H, C₁₋₁₀ alkylene-H, C₂₋₁₀ alkenylene-H, C₂₋₁₀ alkynylene-H, (CRR')_rO(CRR')_r-H, (CRR')_rNR^a(CRR')_r-H, (CRR')_rC(O)(CRR')_r-H, (CRR')_rC(O)O(CRR')_r-H, (CRR')_rOC(O)(CRR')_r-H, (CRR')_rC(O)NR^a(CRR')_r-H, (CRR')_rNR^aC(O)(CRR')_r-H, (CRR')_rOC(O)O(CRR')_r-H, (CRR')_rOC(O)NR^a(CRR')_r-H, (CRR')_rNR^aC(O)O(CRR')_r-H, (CRR')_rNR^aC(O)NR^a(CRR')_r-H, (CRR')_rS(O)_p(CRR')_r-H, (CRR')_rSO₂NR^a(CRR')_r-H, (CRR')_rNR^aSO₂(CRR')_r-H, and (CRR')_rNR^aSO₂NR^a(CRR')_r-H;

25

30

alternatively, R³ and R⁴ combine to form a C₃₋₁₃ carbocyclic residue substituted with R^{1'} and 0-3 R^b or a 5-14 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with R^{1'} and 0-3 R^b;

35

R^a , at each occurrence, is independently selected from H, C_{1-4} alkyl, phenyl and benzyl;

5 $R^{a'}$, at each occurrence, is independently selected from H, C_{1-4} alkyl, phenyl and benzyl;

$R^{a''}$, at each occurrence, is independently selected from H, C_{1-4} alkyl, benzyl, C_{3-7} carbocyclic residue, or a 5 to 6
10 membered heteroaromatic ring containing 1-4 heteroatoms selected from the group consisting of N, O, and S;

alternatively, R^a and $R^{a'}$ taken together with the nitrogen to which they are attached form a 5 or 6 membered ring
15 containing from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

R^b , at each occurrence, is independently selected from C_{1-6} alkyl, OR^a , Cl, F, Br, I, =O, CN, NO_2 , $NR^aR^{a'}$, $C(O)R^{a''}$,
20 $C(O)OR^a$, $C(O)NR^aR^{a'}$, $S(O)_2NR^aR^{a'}$, $S(O)_pR^a$, CF_3 , and CF_2CF_3 ;


R^c , at each occurrence, is independently selected from C_{1-6} alkyl, OR^a , Cl, F, Br, I, =O, CN, NO_2 , $NR^aR^{a'}$, $C(O)R^a$,
 $C(O)OR^a$, $C(O)NR^aR^{a'}$, $NR^aC(O)NR^aR^{a'}$, $S(O)_2NR^aR^{a'}$, $S(O)_pR^a$,
25 CF_3 , CF_2CF_3 , $-CH(=NOH)$, $-C(=NOH)CH_3$, $(CRR')_sO(CRR')_sR^d$,
 $(CRR')_sS(O)_p(CRR')_sR^d$, $(CRR')_sNR^a(CRR')_sR^d$, phenyl, and
a 5-14 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O,
and S;

30 R^5 , at each occurrence, is selected from C_{1-10} alkyl substituted with 0-2 R^b , and C_{1-8} alkyl substituted with 0-2 R^d ;

35 R^d , at each occurrence, is independently selected from phenyl substituted with 0-3 R^b , biphenyl substituted with 0-2 R^b , naphthyl substituted with 0-3 R^b and a 5-10 membered heteroaryl system containing from 1-4 heteroatoms

selected from the group consisting of N, O, and S and substituted with 0-3 R^b;

R⁶, at each occurrence, is selected from phenyl, naphthyl,
 5 C₁₋₁₀ alkyl-phenyl-C₁₋₆ alkyl-, C₃₋₁₁ cycloalkyl, C₁₋₆
 alkylcarbonyloxy-C₁₋₃ alkyl-, C₁₋₆ alkoxy carbonyloxy-C₁₋₃
 alkyl-, C₂₋₁₀ alkoxy carbonyl, C₃₋₆ cycloalkylcarbonyloxy-
 C₁₋₃ alkyl-, C₃₋₆ cycloalkoxy carbonyloxy-C₁₋₃ alkyl-, C₃₋₆
 cycloalkoxy carbonyl, phenoxy carbonyl,
 10 phenyloxy carbonyloxy-C₁₋₃ alkyl-, phenylcarbonyloxy-C₁₋₃
 alkyl-, C₁₋₆ alkoxy-C₁₋₆ alkylcarbonyloxy-C₁₋₃ alkyl-, [5-
 (C₁₋₅ alkyl)-1,3-dioxo-cyclopenten-2-one-yl]methyl, (5-
 aryl-1,3-dioxo-cyclopenten-2-one-yl)methyl, -C₁₋₁₀ alkyl-
 NR⁷R^{7a}, -CH(R⁸)OC(=O)R⁹, -CH(R⁸)OC(=O)OR⁹, and



R⁷ is selected from H and C₁₋₁₀ alkyl, C₂₋₆ alkenyl, C₃₋₆
 cycloalkyl-C₁₋₃ alkyl-, and phenyl-C₁₋₆ alkyl-;

20 R^{7a} is selected from H and C₁₋₁₀ alkyl, C₂₋₆ alkenyl, C₃₋₆
 cycloalkyl-C₁₋₃ alkyl-, and phenyl-C₁₋₆ alkyl-;

R⁸ is selected from H and C₁₋₄ linear alkyl;

25 R⁹ is selected from H, C₁₋₈ alkyl substituted with 1-2 R^e, C₃₋₈
 cycloalkyl substituted with 1-2 R^e, and phenyl
 substituted with 0-2 R^b;

30 R^e, at each occurrence, is selected from C₁₋₄ alkyl, C₃₋₈
 cycloalkyl, C₁₋₅ alkoxy, phenyl substituted with 0-2 R^b;

p, at each occurrence, is selected from 0, 1, and 2;

r, at each occurrence, is selected from 0, 1, 2, 3, 4, and 5;

r', at each occurrence, is selected from 0, 1, 2, 3, 4, and 5;

r'', at each occurrence, is selected from 1, 2, and 3;

5

s, at each occurrence, is selected from 0, 1, 2, and 3; and,

s', at each occurrence, is selected from 0, 1, 2, and 3.

10

2. A compound according to Claim 1, wherein:

A is selected from COR⁵, -CO₂H, CH₂CO₂H, -CONHOH, -CONHOR⁵,
-CONHOR⁶, -N(OH)COR⁵, -SH, and -CH₂SH;

15

ring B is a 4-7 membered cyclic amide containing from 0-2
additional heteroatoms selected from O, NR^a, and S(O)_p,
and 0-1 additional carbonyl groups and 0-1 double bonds;

20 U is absent;

Y is absent;

Z is absent or selected from a C₅₋₁₀ carbocyclic residue
substituted with 0-5 R^b and a 5-10 membered heterocyclic
system containing from 1-4 heteroatoms selected from the
group consisting of N, O, and S and substituted with 0-5
R^b;

30 U^a is absent or is selected from: O, NR^a, C(O), C(O)NR^a,
NR^aC(O), OC(O)NR^a, NR^aC(O)O, NR^aC(O)NR^a, S(O)_pNR^a, and
NR^aS(O)_p;

35 R² is selected from H, Q', C₁₋₅ alkylene-Q', C₂₋₅
alkenylene-Q', C₂₋₅ alkynylene-Q', (CRR')_rO(CRR')_r-Q',
(CRR')_rNR^a(CRR')_r-Q', (CRR')_rNR^aC(O)(CRR')_r-Q',
(CRR')_rC(O)NR^a(CRR')_r-Q', (CRR')_rNR^aC(O)NR^a(CRR')_r-Q',

$(\text{CRR}')_r \cdot \text{C}(\text{O}) (\text{CRR}')_r - \text{Q}'$, $(\text{CRR}')_r \cdot \text{C}(\text{O}) \text{O} (\text{CRR}')_r - \text{Q}'$,
 $(\text{CRR}')_r \cdot \text{S}(\text{O})_p (\text{CRR}')_r - \text{Q}'$, and $(\text{CRR}')_r \cdot \text{SO}_2 \text{NR}^a (\text{CRR}')_r - \text{Q}'$;

Q' is selected from H, phenyl substituted with 0-3 R^b and a
 5 5-6 membered heteroaryl system containing from 1-4
 heteroatoms selected from the group consisting of N, O,
 and S and substituted with 0-3 R^b;

R³ is selected from H, Q, C₁₋₁₀ alkylene-Q, C₂₋₁₀ alkenylene-Q,
 10 C₂₋₁₀ alkynylene-Q, $(\text{CRR}')_r \cdot \text{O} (\text{CRR}')_r - \text{Q}$,
 $(\text{CRR}')_r \cdot \text{NR}^a (\text{CRR}')_r - \text{Q}$, $(\text{CRR}')_r \cdot \text{C}(\text{O}) (\text{CRR}')_r - \text{Q}$,
 $(\text{CRR}')_r \cdot \text{C}(\text{O}) \text{NR}^a (\text{CRR}')_r - \text{Q}$, $(\text{CRR}')_r \cdot \text{NR}^a \text{C}(\text{O}) (\text{CRR}')_r - \text{Q}$,
 $(\text{CRR}')_r \cdot \text{OC}(\text{O}) \text{NR}^a (\text{CRR}')_r - \text{Q}$, $(\text{CRR}')_r \cdot \text{NR}^a \text{C}(\text{O}) \text{O} (\text{CRR}')_r - \text{Q}$,
 $(\text{CRR}')_r \cdot \text{NR}^a \text{C}(\text{O}) \text{NR}^a (\text{CRR}')_r - \text{Q}$, $(\text{CRR}')_r \cdot \text{S}(\text{O})_p (\text{CRR}')_r - \text{Q}$,
 15 $(\text{CRR}')_r \cdot \text{SO}_2 \text{NR}^a (\text{CRR}')_r - \text{Q}$, $(\text{CRR}')_r \cdot \text{NR}^a \text{SO}_2 (\text{CRR}')_r - \text{Q}$, and
 $(\text{CRR}')_r \cdot \text{NR}^a \text{SO}_2 \text{NR}^a (\text{CRR}')_r - \text{Q}$;

R, at each occurrence, is independently selected from H, CH₃,
 and CH₂CH₃;

R', at each occurrence, is independently selected from H and
 CH₃;

Q is selected from H, a C₃₋₁₀ carbocyclic residue substituted
 25 with 0-5 R^b and a 5-10 membered heterocyclic system
 containing from 1-4 heteroatoms selected from the group
 consisting of N, O, and S and substituted with 0-5 R^b;
 and,

R^c, at each occurrence, is independently selected from C₁₋₆
 30 alkyl, OR^a, Cl, F, Br, I, =O, CN, NO₂, NR^aR^{a'}, C(O)R^a,
 C(O)OR^a, C(O)NR^aR^{a'}, S(O)₂NR^aR^{a'}, S(O)_pR^a, CF₃, CF₂CF₃, and
 a 5-10 membered heterocyclic system containing from 1-4
 heteroatoms selected from the group consisting of N, O,
 35 and S.

3. A compound according to Claim 2, wherein:

A is selected from $-\text{CO}_2\text{H}$, $\text{CH}_2\text{CO}_2\text{H}$, $-\text{CONHOH}$, $-\text{CONHOR}^5$, and $-\text{N}(\text{OH})\text{COR}^5$;

- 5 ring B is a 4-6 membered cyclic amide containing from 0-2 additional heteroatoms selected from O, NR^a , and $\text{S}(\text{O})_p$, and 0-1 additional carbonyl groups and 0-1 double bonds;

- 10 Z is absent or selected from a C_{5-6} carbocyclic residue substituted with 0-3 R^b and a 5-9 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with 0-5 R^b ;

- 15 U^a is absent or is selected from: O, NR^a , $\text{C}(\text{O})$, $\text{C}(\text{O})\text{NR}^a$, $\text{NR}^a\text{C}(\text{O})$, and $\text{S}(\text{O})_p\text{NR}^a$;

X^a is absent or C_{1-10} alkylene;

- 20 R^2 is selected from H, C_{1-5} alkylene- Q' , $(\text{CH}_2)_r\text{O}(\text{CH}_2)_r\text{Q}'$, $(\text{CH}_2)_r\text{NR}^a(\text{CH}_2)_r\text{Q}'$, $(\text{CRR}')_r\text{NR}^a\text{C}(\text{O})(\text{CRR}')_r\text{Q}'$, $(\text{CH}_2)_r\text{C}(\text{O})\text{NR}^a(\text{CH}_2)_r\text{Q}'$, $(\text{CRR}')_r\text{NR}^a\text{C}(\text{O})\text{NR}^a(\text{CRR}')_r\text{Q}'$, and $(\text{CH}_2)_r\text{C}(\text{O})(\text{CH}_2)_r\text{Q}'$;

- 25 R^c , at each occurrence, is independently selected from C_{1-6} alkyl, OR^a , Cl, F, Br, I, =O, CN, NO_2 , $\text{NR}^a\text{R}^a'$, $\text{C}(\text{O})\text{R}^a$, $\text{C}(\text{O})\text{OR}^a$, $\text{C}(\text{O})\text{NR}^a\text{R}^a'$, $\text{S}(\text{O})_2\text{NR}^a\text{R}^a'$, $\text{S}(\text{O})_p\text{R}^a$, CF_3 , CF_2CF_3 , and a 5-9 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O,
30 and S; and,

- Q is selected from H, a C_{5-6} carbocyclic residue substituted with 0-5 R^b and a 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group
35 consisting of N, O, and S and substituted with 0-5 R^b .

4. A compound according to Claim 3, wherein:

A is selected from $-\text{CO}_2\text{H}$, $\text{CH}_2\text{CO}_2\text{H}$, $-\text{CONHOH}$, and $-\text{CONHOR}^5$;

ring B is a 4-5 membered cyclic amide containing from 0-2 additional heteroatoms selected from O, NR^a , and $\text{S}(\text{O})_p$, and 0-1 additional carbonyl groups and 0-1 double bonds;

X is absent or selected from C_{1-4} alkylene, C_{2-4} alkenylene, and C_{2-4} alkynylene;

Z is absent or selected from phenyl substituted with 0-3 R^b and a 5-9 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with 0-3 R^b ;

X^a is absent or C_{1-4} alkylene;

Y^a is absent or selected from O and NR^a ;

Z^a is selected from H, a C_{5-10} carbocyclic residue substituted with 0-5 R^c and a 5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with 0-5 R^c ;

R^4 is selected from H, C_{1-4} alkylene-H, $(\text{CH}_2)_r\text{O}(\text{CH}_2)_r\text{-H}$, and $(\text{CH}_2)_r\text{NR}^a(\text{CH}_2)_r\text{-H}$; and,

R^c , at each occurrence, is independently selected from C_{1-6} alkyl, OR^a , Cl, F, Br, I, $=\text{O}$, CN, NO_2 , $\text{NR}^a\text{R}^a'$, $\text{C}(\text{O})\text{R}^a$, $\text{C}(\text{O})\text{OR}^a$, $\text{C}(\text{O})\text{NR}^a\text{R}^a'$, $\text{S}(\text{O})_2\text{NR}^a\text{R}^a'$, $\text{S}(\text{O})_p\text{R}^a$, CF_3 , CF_2CF_3 , and a 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S.

5. A compound according to Claim 1, wherein:

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-(phenylmethoxy)phenyl]-1-pyrrolidineacetamide;

5 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-(4-methoxyphenyl)-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-(1-methylethoxy)phenyl]-2-oxo-1-pyrrolidineacetamide;

10 [1(R)]-3-[4-(1,1-dimethylethoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-(4-(cyclohexyloxy)phenyl)-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

15 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-[4-(1,1-dimethylethyl)phenylmethoxy]phenyl]-1-pyrrolidineacetamide;

20 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-(trans-3-phenyl-2-propenyloxy)phenyl]-1-pyrrolidineacetamide;

[1(R)]-3-[4-[(3-methylphenyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

25 [1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

30 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-(2-propenyloxy)phenyl]-1-pyrrolidineacetamide;

[1(R)]-3-[4-[(3-cyanophenyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

35 [1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-[(2-nitrophenyl)methoxy]phenyl]-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α -3-dimethyl-3-[4-[(3-nitrophenyl)methoxy]phenyl]-2-oxo-1-pyrrolidineacetamide;

5 [1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-[(4-nitrophenyl)methoxy]phenyl]-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-[(1-naphthalenyl)methoxy]phenyl]-2-oxo-1-pyrrolidineacetamide;

10

[1(R)]-N-hydroxy-3-(4-hydroxyphenyl)- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

15 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-[(2-pyridinyl)methoxy]phenyl]-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-[(3-pyridinyl)methoxy]phenyl]-1-pyrrolidineacetamide;

20 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-[(4-pyridinyl)methoxy]phenyl]-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-(2-methylpropyl)phenyl]-2-oxo-1-pyrrolidineacetamide;

25

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-phenyl-1-pyrrolidineacetamide;

N-hydroxy-2-oxo-3-phenyl-1-pyrrolidineacetamide;

30

(+/-)-N-hydroxy-3-methyl-2-oxo-3-phenyl-1-pyrrolidineacetamide;

35 [1(R)]-N-hydroxy- α -methyl-2-oxo-3-phenyl-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy-3-(4-methoxyphenyl)- α -methyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-cyclohexyl-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

5 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-(2-phenylethyl)-1-pyrrolidineacetamide;

[1(R)]-3-(2-cyclohexylethyl)-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

10

[1(R)]-N-hydroxy- α -methyl-2-oxo-3-phenyl-3-(phenylmethyl)-2-oxo-1-pyrrolidineacetamide;

15

[1(R)]-3,4,4',5'-tetrahydro-N-hydroxy- α -methyl-2-oxospiro[naphthalene-2(1H),3'-[3H]pyrrole]-1'(2'H)-acetamide;

20

[1(R)]-3-[4-[(3,5-dibromophenyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

25

[1(R)]-3-[4-[[3,5-bis(trifluoromethyl)phenyl]methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[4-[(3,5-dichlorophenyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

30

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-[(2-methyl-1-naphthalenyl)methoxy]phenyl]-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[4-[(3,5-dimethoxyphenyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

35

[1(R)]-3-[4-[[4-chloro-2-(trifluoromethyl)-6-quinolinyl]methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-[4-(1,2,3-thiadiazol-4-yl)phenyl]methoxy]phenyl]-1-pyrrolidineacetamide;

5 [1(R)]-3-[4-([1,1'-biphenyl]-2-ylmethoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide ;

10

[1(R)]-3-[4-(1H-benzotriazol-1-ylmethoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

15

[1(R)]-3-[4-[(4,6-dimethyl-2-pyrimidinyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide ;

[1(R)]-3-[4-(1,3-benzodioxol-5-ylmethoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

20

[1(R)]-3-[4-[(2-chloro-6-ethoxy-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide;

25

[1(R)]-3-[4-[(4,5-dimethyl-2-thiazolyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

30

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-[(3-methyl-5-nitrophenyl)methoxy]phenyl]-2-oxo-1-pyrrolidineacetamide;

35

[1(R)]-3-[4-[(3-amino-5-methylphenyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[4-[[3-(acetylamino)-5-methylphenyl]methoxy]phenyl]-
N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

5 [1(R)]-1,1-dimethylethyl [2-[[3-[[4-[1-[2-(hydroxyamino)-1-
methyl-2-oxoethyl]-3-methyl-2-oxo-3-
pyrrolidinyl]phenoxy]methyl]-5-methylphenyl]amino]-2-
oxoethyl]carbamate;

10 [1(R)]-3-[4-[[3-[(aminoacetyl)amino]-5-
methylphenyl]methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-
oxo-1-pyrrolidineacetamide;

15 [1(R)]-1,1-dimethylethyl [2-[[2-[[3-[[4-[1-[2-(hydroxyamino)-
1-methyl-2-oxoethyl]-3-methyl-2-oxo-3-
pyrrolidinyl]phenoxy]methyl]-5-methylphenyl]amino]-2-
oxoethyl]amino]-2-oxoethyl]carbamate;

20 [1(R)]-3-[4-[[3-[[[(aminoacetyl)amino]acetyl]amino]-5-
methylphenyl]methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-
oxo-1-pyrrolidineacetamide;

25 [1(R)]-N-[3-[[4-[1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-3-
methyl-2-oxo-3-pyrrolidinyl]phenoxy]methyl]-5-
methylphenyl]-4-morpholinecarboxamide;

3- [4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-
 α , α ,3-trimethyl-2-oxo-1-pyrrolidineacetamide;

30 [1(R)]-3-[1,1'-biphenyl]-4-yl-N-hydroxy- α ,3-dimethyl-2-oxo-1-
pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-3-(2'-methyl[1,1'-biphenyl]-4-
yl)-2-oxo-1-pyrrolidineacetamide;

35 [1(R)]-N-hydroxy- α ,3-dimethyl-3-(4'-methyl[1,1'-biphenyl]-4-
yl)-2-oxo-1-pyrrolidineacetamide;

[1(R)-3-(3',4'-dimethoxy[1,1'-biphenyl]-4-yl)-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

5 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[2'-(trifluoromethyl)[1,1'-biphenyl]-4-yl]-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-(4-methylphenoxy)phenyl]-2-oxo-1-pyrrolidineacetamide;

10 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-(4-phenoxyphenyl)-1-pyrrolidineacetamide;

15 [1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-(2-methylphenoxy)phenyl]-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[4-(3,5-dichlorophenoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

20 [1(R)]-3-[4-(3,4-dimethoxyphenoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[4-(1,3-benzodioxol-5-yloxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

25 [1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-[3-(1-methylethyl)phenoxy]phenyl]-2-oxo-1-pyrrolidineacetamide;

30 [1(R)]-N-hydroxy-3-[4-(3-methoxyphenoxy)phenyl]- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-(3-thienyloxy)phenyl]-1-pyrrolidineacetamide;

35 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-(3,4,5-trimethoxyphenoxy)phenyl]-1-pyrrolidineacetamide;

[1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-(1-naphthalenyloxy)phenyl]-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy-3-[4-[3-[(hydroxyimino)methyl]phenoxy]phenyl]- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy-3-[4-[4-[1-(hydroxyimino)ethyl]phenoxy]phenyl]- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[4-([1,1'-biphenyl]-4-yloxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[4-(3,5-dibromophenoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[4-[3-(acetylamino)phenoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-(4-nitrophenoxy)phenyl]-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-3-(4-methylphenyl)-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)oxy]methyl]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide ;

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-[(4-quinolinyl)oxy]methyl]phenyl]-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-3-(4-nitrophenyl)-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-
[(phenylcarbonyl)amino]phenyl]-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-
5 [(phenylsulfonyl)amino]phenyl]-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-
[[(phenylamino) carbonyl] amino]phenyl]-1-
pyrrolidineacetamide;

10

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[4-[(1-
naphthalenylmethyl)amino]phenyl]-2-oxo-1-
pyrrolidineacetamide;

15 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[4-[(4-
quinolinylmethyl)amino]phenyl]-1-pyrrolidineacetamide;

[1(R)]-3-[4-[[(3,5-dimethoxyphenyl)methyl] amino]phenyl]-N-
hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

20

3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy-3-methyl-
2-oxo-1-pyrrolidineacetamide;

3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-
25 methyl-2-oxo-1-pyrrolidineacetamide;

3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-
methyl-2-oxo-1-pyrrolidineacetamide;

30 [1(R)]-N-hydroxy-3-methyl- α -(1-methylethyl)-2-oxo-3-[4-(4-
quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy-3-methyl- α -(1-methylethyl)-2-oxo-3-[4-
(phenylmethoxy)phenyl]-1-pyrrolidineacetamide;

35

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-
hydroxy-3-methyl- α -(1-methylethyl)-2-oxo-1-
pyrrolidineacetamide;

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

5

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

10 [1(R)]-3-[4-[[3,5-bis(trifluoromethyl)phenyl]methoxy]phenyl]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

15 [1(R)]-3-[4-[(3,5-dichlorophenyl)methoxy]phenyl]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-3-[3-(phenylmethoxy)propyl]-1-pyrrolidineacetamide;

20 [1(R)]-N-hydroxy-3-methyl-3-[2-methyl-4-(phenylmethoxy)phenyl]- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

25 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]-2-methylphenyl]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

30 [1(R)]-N-hydroxy-3-methyl-3-[2-methyl-4-(2-naphthalenylmethoxy)phenyl]- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy-3-methyl- α -(2-methylpropyl)-3-[2-methyl-4-(4-pyridinylmethoxy)phenyl]-2-oxo-1-pyrrolidineacetamide;

35 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]-2-methylphenyl]-N-hydroxy-3-methyl- α -(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy-3-methyl- α -[2-(methylthio)ethyl]-2-oxo-3-[4-(phenylmethoxy)phenyl]-1-pyrrolidineacetamide;

5 [1(R)]-3-[4-(3,5-dibromophenoxy)phenyl]-3-methyl- α -[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetic acid;

[1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-N-hydroxy-3-methyl- α -[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide;

10

[1(R)]-3-[4-(3,5-dibromophenoxy)phenyl]-N-hydroxy-3-methyl- α -[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide;

15

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl- α -[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide;

20

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl- α -[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide ;

[1(R)]-N-hydroxy-3-methyl- α -[2-(methylsulfonyl)ethyl]-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide;

25

N-hydroxy-1-[3-methyl-2-oxo-3-[4-(phenylmethoxy)phenyl]-1-pyrrolidinyl]cyclopropanecarboxamide;

[1(R)]-N-hydroxy- α -[(4-hydroxyphenyl)methyl]-3-methyl-2-oxo-3-[4-(phenylmethoxy)phenyl]-1-pyrrolidineacetamide;

30

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α -(2-hydroxyethyl)-3-methyl-2-oxo-1-pyrrolidineacetamide;

35

[1(R)]-1,1-dimethylethyl [5-[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]carbamate;

[1(R)]- α -(4-aminobutyl)-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide;

5 [1(R)]- α -[4-(acetylamino)butyl]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide;

10 [1(R)]-N-[5-[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]-3-pyridineacetamide;

15 [1(R)]-N-[5-[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]-4-morpholinecarboxamide;

20 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl- α -[4-[(methylsulfonyl)amino]butyl]-2-oxo-1-pyrrolidineacetamide;

[1(R)]- α -[4-(acetylamino)butyl]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide;

25 [1(R)]-1,1-dimethylethyl [5-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]carbamate;

30 [1(R)]- α -(4-aminobutyl)-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide;

35 [1(R)]- α -[4-[(aminoacetyl)amino]butyl]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]- α -[4-(acetylamino)butyl]-3-[4-[[3,5-bis(trifluoromethyl)phenyl]methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide;

5 [1(R)]-1,1-dimethylethyl [5-[3-[4-(3,5-dibromophenoxy)phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]carbamate;

10 [1(R)]- α -(4-aminobutyl)-3-[4-(3,5-dibromophenoxy)phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide;

15 [1(R)]-1,1-dimethylethyl [3-[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-4-(hydroxyamino)-4-oxobutyl]carbamate;

[1(R)]- α -(2-aminoethyl)-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide;

20 [1(R)]- α -[2-(acetylamino)ethyl]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide;

25 [1(R)]-1,1-dimethylethyl [3-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-4-(hydroxyamino)-4-oxobutyl]carbamate;

30 [1(R)]- α -(2-aminoethyl)-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide;

N-[3-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-4-(hydroxyamino)-4-oxobutyl]-3-pyridinecarboxamide;

35 [1(R)]-N-[3-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-4-(hydroxyamino)-4-oxobutyl]-4-morpholinecarboxamide;

[1(R)]-1,1-dimethylethyl [2-[[3-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-4-(hydroxyamino)-4-oxobutyl]amino]-2-oxoethyl]carbamate;

5

[1(R)]- α -[2-[(aminoacetyl)amino]ethyl]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide;

10 [1(R)]-1,1-dimethylethyl [2-[[2-[[3-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-4-(hydroxyamino)-4-oxobutyl]amino]-2-oxoethyl]amino]-2-oxoethyl]carbamate;

15 [1(R)]- α -[2-[[[(aminoacetyl)amino]acetyl]amino]ethyl]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-methyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy-3-methyl-2-oxo- α -[(phenylmethoxy)methyl]-3-[4-(phenylmethoxy)phenyl]-1-pyrrolidineacetamide;

20

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α -(hydroxymethyl)-3-methyl-2-oxo-1-pyrrolidineacetamide;

25

[1(R)]-1,1-dimethylethyl 4-[2-(hydroxyamino)-1-[3-methyl-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-2-oxoethyl]-1-piperidinecarboxylate;

30 [1(R)]-N-hydroxy- α -[3-methyl-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-4-piperidineacetamide;

[1(R)]-N-hydroxy- α -[3-methyl-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-1-(methylsulfonyl)-4-piperidineacetamide;

35

[1(R)]-1-(2-furanylcarbonyl)-N-hydroxy- α -[3-methyl-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidinyl]-4-piperidineacetamide;

5 [1(R)]-1,1-dimethylethyl 4-[1-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-2-(hydroxyamino)-2-oxoethyl]-1-piperidinecarboxylate;

10 [1(R)]- α -[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide;

15 [1(R)]-methyl 4-[1-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-2-(hydroxyamino)-2-oxoethyl]-1-piperidinecarboxylate;

20 [1(R)]- α -[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-N-hydroxy-1-(methylsulfonyl)-4-piperidineacetamide;

[1(R)]-1-acetyl- α -[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide;

25 [1(R)]-1-(2,2-dimethyl-1-oxopropyl)- α -[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-N-hydroxy-4-piperidineacetamide;

30 [1(R)]- α -[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-N-hydroxy-1-methyl-4-piperidineacetamide;

35 [1(R)]- α -[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-3-methyl-2-oxo-1-pyrrolidinyl]-N-hydroxy-1-(1-methylethyl)-4-piperidineacetamide;

[1(R)]-3-amino-N-hydroxy- α -(2-methylpropyl)-2-oxo-3-[4-(2-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide;

[1(R)]-3-amino-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-1-pyrrolidineacetamide;

5 [1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-3-
[[(ethylamino) carbonyl] amino]-N-hydroxy-alpha-methyl-2-
oxo-1-pyrrolidineacetamide;

10 [1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy-
alpha-methyl-3-[(methylsulfonyl) amino]-2-oxo-1-
pyrrolidineacetamide;

15 [1(R)]-N-[3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-1-[2-
(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-
pyrrolidinyl]-3-pyridineacetamide;

20 [1(R)]-N-[3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-1-[2-
(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-
pyrrolidinyl]-4-pyridinecarboxamide;

[1(R)]-3-amino-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]
phenyl]-N-hydroxy-alpha-methyl-2-oxo-1-
pyrrolidineacetamide;

25 [1(R)]-N-[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-1-[2-
(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-
pyrrolidinyl]-4-pyridinecarboxamide;

30 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-
[[(ethylamino) carbonyl] amino]-N-hydroxy-alpha-methyl-2-
oxo-1-pyrrolidineacetamide;

35 [1(R)]-1,1-dimethylethyl [2-[[3-[4-[(2,6-dichloro-4-
pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-
oxoethyl]-2-oxo-3-pyrrolidinyl] amino]-2-
oxoethyl] carbamate;

[1(R)]-3-[(aminoacetyl)amino]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-1-pyrrolidineacetamide;

5 [1(R)]-N-[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]-3-pyridineacetamide;

10 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-3-[[[(phenylmethyl)amino]carbonyl]amino]-1-pyrrolidineacetamide;

15 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-[[[(2,4-dimethoxyphenyl)amino]carbonyl]amino]-N-hydroxy-alpha-methyl-2-oxo-1-pyrrolidineacetamide;

20 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-3-[[[(phenylamino)carbonyl]amino]-1-pyrrolidineacetamide;

25 [1(R)]-1,1-dimethylethyl [3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]carbamate;

30 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-3-[[[2-(4-morpholinyl)ethyl]amino]carbonyl]amino]-2-oxo-1-pyrrolidineacetamide;

[1(R)]-1,1-dimethylethyl N-[[[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]amino]carbonyl]glycine;

35 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-3-[[2-thiazolylamino]carbonyl]amino]-1-pyrrolidineacetamide;

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-3-[[4-pyridinylamino)carbonyl]amino]-1-pyrrolidineacetamide;

5 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-[[[(3-hydroxyphenyl)amino]carbonyl]amino]-alpha-methyl-2-oxo-1-pyrrolidineacetamide;

10 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-3-[[[(2,3-dihydro-2-oxo-1H-benzimidazol-5-yl)amino]carbonyl]amino]-N-hydroxy-alpha-methyl-2-oxo-1-pyrrolidineacetamide;

15 [1(R)]-3-amino-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide;

20 [1(R)]-3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidine acetamide;

25 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-3-[[2-thiazolylamino)carbonyl]amino]-1-pyrrolidineacetamide;

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-3-[[2-thiazolylamino)carbonyl]amino]-1-pyrrolidineacetamide;

30 [5(R)]-2-propenyl [5-[3-amino-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]carbamate;

35 [5(R)]-2-propenyl [5-[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]carbamate;

[1(R)]-3-amino-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

5 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[[2-thiazolylamino)carbonyl]amino]-1-pyrrolidineacetamide;

10 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[[2-thiazolylamino)carbonyl]amino]-1-pyrrolidineacetamide;

15 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[[2-pyridinylamino)carbonyl]amino]-1-pyrrolidineacetamide;

20 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[(trifluoroacetyl)amino]-1-pyrrolidineacetamide;

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[[2-pyridinylamino)carbonyl]amino]-1-pyrrolidineacetamide;

25 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[[[(phenylsulfonyl)amino]carbonyl]amino]-1-pyrrolidineacetamide;

30 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[[[(phenylsulfonyl)amino]carbonyl]amino]-1-pyrrolidineacetamide;

35 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-[[[(3-methyl-5-isothiazolyl)amino]carbonyl]amino]-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[[(1H-benzimidazol-2-ylamino) carbonyl] amino]-3-[4-
[(2,6-dichloro-4-pyridinyl) methoxy] phenyl]-N-hydroxy-
alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

5

[1(R)]-3-[[(1H-benzimidazol-2-ylamino) carbonyl] amino]-3-[4-
[(2,6-dimethyl-4-pyridinyl) methoxy] phenyl]-N-hydroxy-
alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

10 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl) methoxy] phenyl]-N-
hydroxy-alpha-(2-methylpropyl)-2-oxo-3-
[[(phenylamino) carbonyl] amino]-1-pyrrolidineacetamide;

15 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl) methoxy] phenyl]-N-
hydroxy-alpha-(2-methylpropyl)-2-oxo-3-
[[(phenylamino) carbonyl] amino]-1-pyrrolidineacetamide;

20 [1(R)]-1-[1-[(hydroxyamino) carbonyl]-3-methylbutyl]-N,N,N-
trimethyl-2-oxo-3-[4-(phenylmethoxy) phenyl]-1-
pyrrolidinemethanaminium;

[1(R)]-3-amino-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[4-(4-
quinolinylmethoxy) phenyl]-1-pyrrolidineacetamide;

25 [1(R)]-3-amino-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[4-(2-
oxo-2-phenylethoxy) phenyl]-1-pyrrolidineacetamide;

30 [1(R)]-3-amino-3-[4-[(3,5-dimethyl-4-
isoxazolyl) methoxy] phenyl]-N-hydroxy-alpha-(2-
methylpropyl)-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-amino-3-[4-[(2,6-dimethyl-4-
pyridinyl) methoxy] phenyl]-N-hydroxy-alpha-(2-
methylpropyl)-2-oxo-1-pyrrolidineacetamide;

35

[1(R)]-3-amino-3-[4-[2-(2-benzothiazolylamino)-2-
oxoethoxy] phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-
1-pyrrolidineacetamide;

[1(R)]-3-amino-N-hydroxy-3-[4-[(2-methoxy-4-quinolinyl)methoxy]phenyl]-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

5

[1(R)]-3-amino-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[4-[(2-phenyl-4-quinolinyl)methoxy]phenyl]-1-pyrrolidineacetamide;

10

[1(R)]-3-amino-3-[4-[(2,6-dimethyl-4-quinolinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

15

[1(R)]-3-amino-3-[4-[(2-chloro-4-quinolinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

20

[1(R)]-3-amino-3-[4-[2-(2,5-dimethoxyphenyl)-2-(hydroxyimino)ethoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

25

[1(R)]-3-amino-N-hydroxy-3-[4-[(2-methylimidazo[1,2-a]pyridin-3-yl)methoxy]phenyl]-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

30

[1(R)]-3-amino-3-[4-[[1,4-dimethyl-2-(methylthio)-1H-imidazol-5-yl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

35

[1(R)]-3-amino-3-[4-[(2,4-dimethyl-5-thiazolyl)methoxy]phenyl]-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-amino-N-hydroxy-alpha-(2-methylpropyl)-3-[4-[(2-methyl-4-quinolinyl)methoxy]phenyl]-2-oxo-1-pyrrolidineacetamide;

5 [1(R)]-3-amino-3-[4-[(2-chloro-4-quinolinyl)methoxy]phenyl]-N-hydroxy-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide;

10 [1(R)]-3-amino-N-hydroxy-3-[4-[(2-methyl-4-quinolinyl)methoxy]phenyl]-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide;

15 [1(R)]-3-amino-3-[4-[(3,5-dimethoxyphenyl)methoxy]phenyl]-N-hydroxy-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide;

20 [1(R)]-3-amino-N-hydroxy-3-[4-[(2-methoxy-4-quinolinyl)methoxy]phenyl]-alpha-[2-(methylsulfonyl)ethyl]-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-amino-3-[4-[(3,5-dimethoxyphenyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

25 [1(R)]-3-(aminomethyl)-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

30 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-(2-methylpropyl)-2-oxo-3-[[[(2-thiazolylamino)carbonyl]amino]methyl]-1-pyrrolidineacetamide;

35 [1(R)]-3-(aminomethyl)-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2-oxo-3-[[[(2-thiazolylamino)carbonyl]amino]methyl]-1-pyrrolidineacetamide;

5

[1(R)]-4-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy-alpha,4-dimethyl-5-oxo-1-imidazolidineacetamide;

[1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy-3-(hydroxymethyl)-alpha-methyl-2-oxo-1-pyrrolidineacetamide;

10

[1(R)]-[3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]methyl ethylcarbamate;

15

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-(hydroxymethyl)-alpha-methyl-2-oxo-1-pyrrolidineacetamide;

20

[1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy-alpha,3-dimethyl-2-oxo-1-azetidineacetamide;

[1(R)]-3-[5-[(3,5-dimethylphenoxy)methyl]-2-thiazolyl]-N-hydroxy-alpha,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

25

[1(R)]-4-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy-alpha-methyl-2,5-dioxo-4-(2-propenyl)-1-imidazolidineacetamide;

30

[1(R)]-N-hydroxy-alpha,3-dimethyl-2-oxo-3-[[4-(phenylmethoxy)phenyl]methyl]-1-pyrrolidineacetamide;

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-(methylamino)-alpha-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

35

[1(R)]-N-hydroxy-3-(methylamino)-alpha-(2-methylpropyl)-3-[4-
[(2-methyl-4-quinolinyl)methoxy]phenyl]-2-oxo-1-
pyrrolidineacetamide;

5 [1(R)]-alpha,3-dimethyl-N-hydroxy-2-oxo-3-[4-
(phenylmethoxy)phenyl]-1-piperidineacetamide ;

[1(R)]-alpha-[3-amino-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-
pyrrolidinyl]-N-hydroxy-4-piperidineacetamide;

10

[1(R)]-alpha-[3-amino-3-[4-[(2,6-dichloro-4-
pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-
hydroxy-4-piperidineacetamide;

15 [1(R)]-1,1-dimethylethyl 4-[1-[3-[(1,1-
dimethylethoxy)carbonyl]amino]-3-[4-[(1,1-dimethyl-4-
pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-2-
(hydroxyamino)-2-oxoethyl]-1-piperidinecarboxylate;

20 [1(R)]-alpha-[3-amino-3-[4-[(2,6-dimethyl-4-
pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-
hydroxy-4-piperidineacetamide;

[1(R)]-alpha-[3-amino-3-[4-[(2,6-dimethyl-4-
25 pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-
hydroxy-1-(methylsulfonyl)-4-piperidineacetamide;

[1(R)]-1-acetyl-alpha-[3-amino-3-[4-[(2,6-dimethyl-4-
pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-
30 hydroxy-4-piperidineacetamide;

[1(R)]-alpha-[3-amino-3-[4-[(2,6-dimethyl-4-
pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-1-(2,2-
dimethyl-1-oxopropyl)-N-hydroxy-4-piperidineacetamide;

35

[1(R)]-1,1-dimethylethyl 4-[1-[3-amino-3-[4-[(2,6-dimethyl-4-
pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-2-
(hydroxyamino)-2-oxoethyl]-1-piperidinecarboxylate;

[1(R)]-methyl 4-[1-[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-2-(hydroxyamino)-2-oxoethyl]-1-piperidinecarboxylate;

5

[1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-N-hydroxy-1-methyl-4-piperidineacetamide;

10

[1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-1-dimethylcarbonyl-N-hydroxy-4-piperidineacetamide ;

15

[1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-1-cyclopropanecarbonyl-N-hydroxy-4-piperidineacetamide ;

20

[1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide;

[1(R)]-3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α -(1-methylethyl)-2-oxo-1-pyrrolidineacetamide;

25

[1(R)]-3-amino- α -cyclohexyl-N-hydroxy-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide;

30

[1(R)]-3-amino- α -cyclohexyl-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-2-oxo-1-pyrrolidineacetamide;

35

3-amino- α -(1,1-dimethylethyl)-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-amino- α -(1,1-dimethylethyl)-N-hydroxy-2-oxo-3-[4-(4-quinolinylmethoxy)phenyl]-1-pyrrolidineacetamide;

[1(R)]-3-amino- α -(1,1-dimethylethyl)-N-hydroxy-2-oxo-3-[4-[(2-methyl-4-quinolinyl)methoxy]phenyl]-1-pyrrolidineacetamide;

5 [1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-[(2-methyl-4-quinolinyl)methoxy]phenyl]-1-pyrrolidineacetamide;

10 [1(R)]-3-amino-N-hydroxy- α -(1-methylethyl)-2-oxo-3-[4-[(2,6-dimethyl-4-quinolinyl)methoxy]phenyl]-1-pyrrolidineacetamide;

15 [1(R)]-N-[4-[1-[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-2-(hydroxyamino)-2-oxoethyl]-1-piperidine]-4-morpholinecarboxamide;

20 [1(R)]- α -[3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-1-pyrrolidinyl]-1-(2-methyl-1-oxopropyl)-N-hydroxy-4-piperidineacetamide ;

25 [1(R)]-3-amino-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α -(4-methoxycyclohexyl)-2-oxo-1-pyrrolidineacetamide;

[1'(R)]-N-hydroxy-1,2-dihydro- α -(1-methylethyl)-2,2'-dioxo-6-(phenylmethoxy)spiro[3H-indole-3,3'-pyrrolidine]-1'-acetamide;

30 [1(R)]-N-hydroxy- α ,3-dimethyl-2-oxo-3-[3-(phenylmethoxy)phenyl]-1-pyrrolidineacetamide;

[1(R)]-3-[3-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

35 [1(R)]-N-hydroxy- α ,3-dimethyl-3-[3-[(3-methylphenyl)methoxy]phenyl]-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy- α ,3-dimethyl-3-[3-(1-methylethoxy)phenyl]-2-oxo-1-pyrrolidineacetamide;

5 [1(R)]-3-[3-(heptyloxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-1,3,4-thiadiazol-2-yl-1,3-pyrrolidinediacetamide;

10

[1(R)]-1,1-dimethylethyl 1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-[4-(phenylmethoxy)phenyl]-3-pyrrolidineacetate;

15

[1(R)]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-[4-(phenylmethoxy)phenyl]-3-pyrrolidineacetic acid;

[1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-N3-[2-(methylamino)-2-oxoethyl]-2-oxo-1,3-pyrrolidinediacetamide;

20

[1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide;

25

[1(R)]-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N-hydroxy- α -methyl-3-[2-(4-morpholinyl)-2-oxoethyl]-2-oxo-1-pyrrolidineacetamide;

30

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide;

35 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-[2-(4-morpholinyl)ethyl]-1,3-pyrrolidinediacetamide;

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide;

5

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide;

10 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-(3-pyridinylmethyl)-1,3-pyrrolidinediacetamide;

15 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-(2-pyridinylmethyl)-1,3-pyrrolidinediacetamide;

20 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-4-pyridinyl-1,3-pyrrolidinediacetamide;

25 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-N3-(3-methyl-5-isothiazolyl)-2-oxo-1,3-pyrrolidinediacetamide;

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N3-[5-(1,1-dimethylethyl)-1,3,4-thiadizol-2-yl]-N1-hydroxy- α 1-methyl-2-oxo-1,3-pyrrolidinediacetamide;

30 [1(R)]-1,1-dimethylethyl 2-[[[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]acetyl]amino]-4-thiazoleacetate;

35 [1(R)]-2-[[[3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-1-[2-(hydroxyamino)-1-methyl-2-oxoethyl]-2-oxo-3-pyrrolidinyl]acetyl]amino]-4-thiazoleacetic acid;

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-methyl-N3-[4-[2-(methylamino)-2-oxoethyl]-2-thiazolyl]-2-oxo-1,3-pyrrolidinediacetamide;

5

[1(R)]-3-(1H-benzimidazol-2-ylmethyl)-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy- α -methyl-2-oxo-1-pyrrolidineacetamide;

10 [1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N-hydroxy-3-(3H-imidazo(4,5-c)pyridin-2-ylmethyl)- α -methyl-2-oxo-1-pyrrolidineacetamide;

15 [1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide;

20 [1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-N1-hydroxy- α 1-methyl-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide;

25 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-(1-methylethyl)-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide;

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-(1-methylethyl)-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide;

30 [1(R)]- α 1-(cyclohexylmethyl)-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide;

35 [1(R)]- α 1-(cyclohexylmethyl)-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide;

[1(R)]-1,1-dimethylethyl [5-[3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-2-oxo-3-[2-oxo-2-[(4-pyridinylmethyl)amino]ethyl]-1-pyrrolidinyl]-6-(hydroxyamino)-6-oxohexyl]carbamate;

5

[1(R)]- α 1-(4-aminobutyl)-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide;

10 [1(R)]-3-[3-(1H-benzotriazol-1-ylmethoxy)phenyl]-N-hydroxy- α ,3-dimethyl-2-oxo-1-pyrrolidineacetamide;

[1(R)]-N-hydroxy-3,4,4-trimethyl- α -[3-methyl-2-oxo-3[4-(phenylmethoxy)phenyl]-1-pyrrolidinyl]-2,5-dioxo-1-
15 imidazolidinepropanamide ;

[1(R)]-1,1-dimethylethyl 1-[(hydroxyamino)carbonyl]-3-methylbutyl]-2-oxo-3-[4-(phenyl)-3-pyrrolidineacetate;

20 [1(R)]-N1-hydroxy-3-[4-[(3,5-dimethylphenyl)methoxy]phenyl]-N3-[2-(methylamino)-2-oxoethyl]- α -(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide;

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-N3-[2-(methylamino)-2-oxoethyl]- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide;

25

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide;

30

[1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-N1-hydroxy-N3-[2-(methylamino)-2-oxoethyl]- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide;

35

[1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-N3-(4-pyridinylmethyl)-1,3-pyrrolidinediacetamide;

[1(R)]-3-[4-[(2,6-dichloro-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-N3-phenyl-1,3-pyrrolidinediacetamide;

5

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-N3-methyl- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide;

10 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-N3-[2-(1H-imidazol-4-yl)ethyl]- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide;

15 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-N3-[1-(phenylmethyl)-4-piperidinyl]-1,3-pyrrolidinediacetamide;

20 [1(R)]-N3-[2-(dimethylamino)ethyl]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide;

25 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-N3-(4-hydroxyphenyl)- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide;

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N3-hydroxy- α 1-(2-methylpropyl)-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide;

30 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N3-hydroxy-3-(2-hydroxyethyl)- α 1-(2-methylpropyl)-2-oxo-1-pyrrolidineacetamide;

35 [1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N3-(4,5-dimethyl-2-thiazolyl)-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide;

[1(R)]-3-[4-[(2,6-dimethyl-4-pyridinyl)methoxy]phenyl]-N1-hydroxy-N3-1H-indazol-5-yl- α 1-(2-methylpropyl)-2-oxo-1,3-pyrrolidinediacetamide; and,

5 [1(R)]-3-[4-[3,5-bis(trifluoromethyl)phenoxy]phenyl]-N1-hydroxy- α 1-(2-methylpropyl)-2-oxo-N3-2-thiazolyl-1,3-pyrrolidinediacetamide;

or a pharmaceutically acceptable salt form thereof.

10

6. A compound according to Claim 1, wherein:

A is selected from COR⁵, -CO₂H, CH₂CO₂H, -CONHOH, -CONHOR⁵,
15 -CONHOR⁶, -N(OH)COR⁵, -SH, and -CH₂SH;

ring B is a 4-7 membered cyclic amide containing from 0-3 additional heteroatoms selected from O, NR^a, and S(O)_p, and 0-1 additional carbonyl groups and 0-1 double bonds;

20

R¹ and R² combine to form a C₅₋₁₄ carbocyclic residue substituted with R^{1'} and 0-3 R^b or a 5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with R^{1'} and 0-3 R^b;

25

Z^a is selected from H, a C₅₋₁₀ carbocyclic residue substituted with 0-5 R^c and a 5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with 0-5 R^c;

30

R³ is selected from H, Q, C₁₋₁₀ alkylene-Q, C₂₋₁₀ alkenylene-Q, C₂₋₁₀ alkynylene-Q, (CRR')_rO(CRR')_r-Q, (CRR')_rNR^a(CRR')_r-Q, (CRR')_rC(O)(CRR')_r-Q, (CRR')_rC(O)NR^a(CRR')_r-Q, (CRR')_rNR^aC(O)(CRR')_r-Q, (CRR')_rOC(O)NR^a(CRR')_r-Q, (CRR')_rNR^aC(O)O(CRR')_r-Q, (CRR')_rNR^aC(O)NR^a(CRR')_r-Q, (CRR')_rS(O)_p(CRR')_r-Q,

35

$(\text{CRR}')_r \cdot \text{SO}_2 \text{NR}^a (\text{CRR}')_r - \text{Q}$, $(\text{CRR}')_r \cdot \text{NR}^a \text{SO}_2 (\text{CRR}')_r - \text{Q}$, and
 $(\text{CRR}')_r \cdot \text{NR}^a \text{SO}_2 \text{NR}^a (\text{CRR}')_r - \text{Q}$;

R, at each occurrence, is independently selected from H, CH₃,
 5 CH₂CH₃, CH=CH₂, CH=CHCH₃, and CH₂CH=CH₂;


R', at each occurrence, is independently selected from H, CH₃,
 CH₂CH₃, and CH(CH₃)₂;

10 Q is selected from H, a C₃₋₁₀ carbocyclic residue substituted
 with 0-5 R^b and a 5-10 membered heterocyclic system
 containing from 1-4 heteroatoms selected from the group
 consisting of N, O, and S and substituted with 0-5 R^b;

15 R⁴ is selected from H;

R^c, at each occurrence, is independently selected from C₁₋₆
 alkyl, OR^a, Cl, F, Br, I, =O, CN, NO₂, NR^aRA', C(O)R^a,
 C(O)OR^a, C(O)NR^aRA', S(O)₂NR^aRA', S(O)_pRA', CF₃, CF₂CF₃, and
 20 a 5-10 membered heterocyclic system containing from 1-4
 heteroatoms selected from the group consisting of N, O,
 and S.

25 7. A compound according to Claim 6, wherein the compound
 is of formula II:



30 wherein, A is selected from -CO₂H, CH₂CO₂H, -CONHOH, and
 -CONHOR⁵;

ring C is fused to ring G and is a phenyl ring or 5-6 membered
 aromatic heterocycle containing from 1-4 heteroatoms
 35 selected from O, N, and S(O)_p, and ring C is substituted
 with 1 R^{1'};

ring G is a 4-8 membered carbocyclic ring substituted with 0-1 carbonyl groups

5 alternatively, ring G is a 4-8 membered heterocyclic ring containing from 1-2 heteroatoms selected from O and NR^a and substituted with 0-2 carbonyl groups and 0-1 double bonds;

10 U^a is absent or is selected from: O, NR^a, C(O), C(O)NR^a, NR^aC(O), and S(O)_pNR^a;

X^a is absent or C₁₋₄ alkylene;

15 Y^a is absent or selected from O and NR^a;

Z^a is selected from H, phenyl substituted with 0-5 R^c and a 5-9 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with 0-5 R^c;

20 Q is selected from H, a C₅₋₆ carbocyclic residue substituted with 0-5 R^b and a 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S and substituted with 0-5 R^b; and,

25 R^c, at each occurrence, is independently selected from C₁₋₆ alkyl, OR^a, Cl, F, Br, I, =O, CN, NO₂, NR^aR^{a'}, C(O)R^a, C(O)OR^a, C(O)NR^aR^{a'}, S(O)₂NR^aR^{a'}, S(O)_pR^a, CF₃, CF₂CF₃, and a 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S.

35

8. A pharmaceutical composition, comprising: a pharmaceutically acceptable carrier and a therapeutically

effective amount of a compound of one of Claims 1-7 or a pharmaceutically acceptable salt form thereof.

5 9. A method for treating or preventing an inflammatory disorder, comprising: administering to a patient in need thereof a therapeutically effective amount of a compound of one of Claims 1-7 or a pharmaceutically acceptable salt form thereof.

10

 10. A method of treating a condition or disease mediated by MMPs, TNF, aggrecanase, or a combination thereof in a mammal, comprising: administering to the mammal in need of such treatment a therapeutically effective amount of a
15 compound of one of Claims 1-7 or a pharmaceutically acceptable salt form thereof.

 11. A method of treating a condition or disease wherein
20 the disease or condition is referred to as rheumatoid arthritis, osteoarthritis, periodontitis, gingivitis, corneal ulceration, solid tumor growth and tumor invasion by secondary metastases, neovascular glaucoma, multiple sclerosis, or psoriasis in a mammal, comprising: administering to the
25 mammal in need of such treatment a therapeutically effective amount of a compound of one of Claims 1-7 or a pharmaceutically acceptable salt form thereof.

30 12. A method of treating a condition or disease wherein the disease or condition is referred to as fever, cardiovascular effects, hemorrhage, coagulation, cachexia, anorexia, alcoholism, acute phase response, acute infection, shock, graft versus host reaction, autoimmune disease or HIV
35 infection in a mammal comprising administering to the mammal in need of such treatment a therapeutically effective amount of a compound of one of Claims 1-7 or a pharmaceutically acceptable salt form thereof.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/21037

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D207/27 A61K31/40 C07D401/12 C07D417/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97 32846 A (UPJOHN CO ; JACOBSEN E JON (US)) 12 September 1997 see the whole document	1-8
Y	WO 97 16425 A (JONES A BRIAN ; ADAMS ALAN D (US); MERCK & CO INC (US); ACTON JOHN) 9 May 1997 see the whole document	1-8
Y	WO 96 29313 A (PROCTER & GAMBLE) 26 September 1996 see claim 1; examples	1-8

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

26 January 1999

Date of mailing of the international search report

02/02/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lauro, P

INTERNATIONAL SEARCH REPORT
Information

H REPORT
ent family members

Int. Publication No
PCT/US 98/21037

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9732846	A	12-09-1997	AU	2052597 A	22-09-1997
WO 9716425	A	09-05-1997	AU	7604196 A	22-05-1997
WO 9629313	A	26-09-1996	US	5672598 A	30-09-1997
			AU	5255896 A	08-10-1996
			BR	9607772 A	07-07-1998
			CA	2216129 A	26-09-1996
			CZ	9702951 A	13-05-1998
			EP	0815084 A	07-01-1998
			HU	9800813 A	28-07-1998
			NO	974335 A	21-11-1997

LINE 1000-0000
MARK (USPTO)